



University of Kentucky  
UKnowledge

---

Theses and Dissertations--Biosystems and  
Agricultural Engineering

Biosystems and Agricultural Engineering

---

2016

## LONG-TERM EFFECTS OF FORESTRY BEST MANAGEMENT PRACTICES ON HYDROLOGY AND WATER CHEMISTRY IN THREE APPALCHIAN HEADWATER CATCHMENTS

Kameryn I. Wright

University of Kentucky, kamerynwright25@gmail.com

Digital Object Identifier: <http://dx.doi.org/10.13023/ETD.2016.371>

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

---

### Recommended Citation

Wright, Kameryn I., "LONG-TERM EFFECTS OF FORESTRY BEST MANAGEMENT PRACTICES ON HYDROLOGY AND WATER CHEMISTRY IN THREE APPALCHIAN HEADWATER CATCHMENTS" (2016). *Theses and Dissertations--Biosystems and Agricultural Engineering*. 45.  
[https://uknowledge.uky.edu/bae\\_etds/45](https://uknowledge.uky.edu/bae_etds/45)

This Master's Thesis is brought to you for free and open access by the Biosystems and Agricultural Engineering at UKnowledge. It has been accepted for inclusion in Theses and Dissertations--Biosystems and Agricultural Engineering by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## **STUDENT AGREEMENT:**

I represent that my thesis or dissertation and abstract are my original work. Proper attribution has been given to all outside sources. I understand that I am solely responsible for obtaining any needed copyright permissions. I have obtained needed written permission statement(s) from the owner(s) of each third-party copyrighted matter to be included in my work, allowing electronic distribution (if such use is not permitted by the fair use doctrine) which will be submitted to UKnowledge as Additional File.

I hereby grant to The University of Kentucky and its agents the irrevocable, non-exclusive, and royalty-free license to archive and make accessible my work in whole or in part in all forms of media, now or hereafter known. I agree that the document mentioned above may be made available immediately for worldwide access unless an embargo applies.

I retain all other ownership rights to the copyright of my work. I also retain the right to use in future works (such as articles or books) all or part of my work. I understand that I am free to register the copyright to my work.

## **REVIEW, APPROVAL AND ACCEPTANCE**

The document mentioned above has been reviewed and accepted by the student's advisor, on behalf of the advisory committee, and by the Director of Graduate Studies (DGS), on behalf of the program; we verify that this is the final, approved version of the student's thesis including all changes required by the advisory committee. The undersigned agree to abide by the statements above.

Kameryn I. Wright, Student

Dr. Carmen T. Agouridis, Major Professor

Dr. Donald G. Colliver, Director of Graduate Studies

LONG-TERM EFFECTS OF FORESTRY BEST MANAGEMENT PRACTICES  
ON HYDROLOGY AND WATER CHEMISTRY IN THREE APPALACHIAN  
HEADWATER CATCHMENTS

---

THESIS

---

A thesis submitted in partial fulfillment of the  
requirements for the degree of Master of Science in Biosystems and Agricultural  
Engineering in the College of Engineering at the University of Kentucky

By

Kameryn Isaiah Wright

Lexington, Kentucky

Director: Dr. Carmen T. Agouridis, Associate Professor of Biosystems and  
Agricultural Engineering

Lexington, Kentucky

2016

Copyright © Kameryn Isaiah Wright 2016

## ABSTRACT OF THESIS

### LONG-TERM EFFECTS OF FORESTRY BEST MANAGEMENT PRACTICES ON HYDROLOGY AND WATER CHEMISTRY IN THREE APPALACHIAN HEADWATER CATCHMENTS

In 1982, a study was initiated in the Field Branch watershed, in the University of Kentucky's Robinson Forest, to evaluate forestry best management practice (BMP) effectiveness after intensive harvesting. The study utilized a paired watershed approach on three adjacent Field Branch subcatchments. One subcatchment was left as the control, one had BMPs implemented (including a 50-ft undisturbed buffer along the stream), and one was clear-cut to the stream's banks without the use of BMPs (i.e. logger's choice). Prior research has shown that logging can negatively impact watershed functions by altering stream hydrology, geomorphology, water quality, and instream habitat. Thus, the goal of forestry BMPs is to mitigate these impacts; however, information on their effectiveness, especially on the long-term, is limited. Monitoring of three streams, one in each subcatchment, has continued since 1982. In 1985, two years after harvest, results indicated a significant increase in stormflow, baseflow, storm volume as a percentage of rainfall, and curve number in the BMP implemented subcatchment. Conversely, in the clear-cut subcatchment these same parameters were significantly increased in 1984, 1985, 2006, and 2007. Water quality results were mixed and forestry BMPs seemingly added little to no benefit over the clear-cut subcatchment for most monitored constituents.

**KEYWORDS:** Storm hydrology, best management practices, water quality, headwater streams, forest harvesting.

---

Kameryn Isaiah Wright

---

July 29, 2016

LONG-TERM EFFECTS OF FORESTRY BEST MANAGEMENT PRACTICES  
ON HYDROLOGY AND WATER CHEMISTRY IN THREE APPALACHIAN  
HEADWATER CATCHMENTS

By

Kameryn Isaiah Wright

---

Carmen T. Agouridis  
Director of Thesis

---

Donald G. Colliver  
Director of Graduate Studies

---

July 29, 2016

---

For my children Elyja William, Elyse Mae, and Hadley Jane

## **ACKNOWLEDGMENTS**

I would like to begin by thanking my beautiful wife who has supported me every step of the way in this endeavor; I couldn't have done this without her help. Next I would like to thank the two professors who made this possible, Dr. Carmen Agouridis and Dr. Christopher Barton. They were pivotal to the completion of this research and imparted invaluable knowledge and experience. Another thanks to my fellow peers who have made these last few years an enjoyable learning experience and to Tyler Sanderson for the long hours he put in collecting data with me. Also, big thanks to Kenton Sena in the Department of Forestry for his aid in the statistical analysis and to everyone at Robinson Forest who aided in the collection and analysis of the paired watershed data for the past 30+ years. I would also like to thank committee member Dr. Richard Warner and all of the faculty and staff who have imparted knowledge along the way and made this journey possible.

Finally, I would like to thank my parents Kent and Trina Wright for their unwavering support, and to my mother and father-in-law Lisa and Dennis Barker for letting me stay with them while I spent many long hours writing this.

## TABLE OF CONTENTS

|  |     |
|--|-----|
| ACKNOWLEDGMENTS .....  | III |
| TABLE OF CONTENTS .....  | IV  |
| LIST OF FIGURES .....  | VI  |
| LIST OF TABLES .....   | XIV |
| CHAPTER 1: INTRODUCTION .....  | 1   |
| 1.1 INTRODUCTION.....  | 1   |
| 1.2 OBJECTIVES.....  | 2   |
| 1.3 ORGANIZATION OF THESIS .....   | 3   |
| CHAPTER 2: EFFECT OF BEST MANAGEMENT PRACTICES ON THE<br>HYDROLOGY OF THREE APPALACHIAN HEADWATER CATCHMENTS .....                 | 4   |
| 2.1 INTRODUCTION.....  | 4   |
| 2.2 METHODS .....  | 9   |
| 2.2.1 Study Site.....  | 9   |
| 2.2.2 Precipitation Data.....  | 18  |
| 2.2.3 Hydrograph Data.....   | 21  |
| 2.2.3.1 Curve Number Method.....   | 24  |
| 2.2.4 Forest Stand Analysis .....  | 25  |
| 2.2.5 Statistical Analysis .....   | 26  |
| 2.3 RESULTS AND DISCUSSION.....  | 28  |
| 2.3.1 Precipitation Characteristics.....   | 28  |
| 2.3.2 Hydrograph Characteristics.....  | 45  |
| 2.3.2.1 Baseflow, Storm Flow and Total Flow Volumes .....  | 62  |
| 2.3.2.2 Peak Discharge.....  | 71  |
| 2.3.2.3 Stormflow Volume as a Percentage of Rainfall .....   | 76  |
| 2.3.2.4 Time to Peak.....  | 83  |
| 2.3.2.5 Curve Number.....  | 87  |
| 2.3.3 Forest Stand Characteristics .....   | 103 |
| 2.4 CONCLUSIONS.....   | 119 |
| CHAPTER 3: LONG-TERM EFFECTS OF BEST MANAGEMENT PRACTICES<br>ON WATER CHEMISTRY FOR THREE APPALACHIAN HEADWATER<br>CATCHMENTS..... | 121 |
| 3.1 INTRODUCTION.....  | 121 |
| 3.2 METHODS .....  | 125 |
| 3.2.1 Water Quality Data.....  | 125 |
| 3.2.2 Statistical Analysis .....   | 126 |
| 3.3 RESULTS AND DISCUSSION.....  | 127 |
| 3.3.1 Time and Treatment Effects.....  | 127 |
| 3.4 CONCLUSIONS.....   | 157 |



|  |     |
|--|-----|
| CHAPTER 4: CONCLUSIONS.....                            | 158 |
| CHAPTER 5: FUTURE WORK .....                           | 160 |
| APPENDIX A: INDIVIDUAL HYDROGRAPH CHARACTERISTICS..... | 161 |
| APPENDIX B: FWMC TABLES .....                          | 197 |
| REFERENCES .....                                       | 216 |
| VITA .....   | 229 |

## LIST OF FIGURES

|   |    |
|---|----|
| Figure 2.1: Location of University of Kentucky's Robinson Forest labeled in green. ....   | 10 |
| Figure 2.2: Mowbray-Robinson logging operations at Robinson station in Quicksand Kentucky. ....   | 12 |
| Figure 2.3: Robinson Forest hydrologic features and topography. The study area is N. Field Branch (WSA); W. Field Branch (WSB); and S. Field Branch (WSC). ....   | 13 |
| Figure 2.4: Digital elevation model of study site with stream network shown in yellow, WSA shown in blue, WSB in red, and WSC in green. ....  | 14 |
| Figure 2.5: Study site post-harvest (1984) with logging roads shown in brown: WSA is on the far left, WSB is in the middle along with the riparian buffer that was left along the stream, and WSC is on the right. .... | 16 |
| Figure 2.6: Study site 22 years after harvest (2005) with WSA on the far left, WSB in the middle, and WSC on the right. ....  | 17 |
| Figure 2.7: Example of baseflow separation and a suitable (single peak) hydrograph for analysis .....   | 20 |
| Figure 2.8: Example of an unsuitable hydrograph (multiple peaks). ....  | 20 |
| Figure 2.9: Yearly precipitation sums and 27-year precipitation average. Data were taken from nearby Jackson County Airport for 1999 and 2000. ....   | 41 |
| Figure 2.10: Pre-harvest hydrographs for WSA, WSB, and WSC for a storm event (38.1 mm) on August 5, 1982 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....   | 45 |
| Figure 2.11: Pre-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.9 mm) on May 22, 1983 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....   | 46 |
| Figure 2.12: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.4 mm) on February 27, 1984 (NG). WSA=control, WSB=BMP, and WSC=no BMPs. ....  | 46 |

|  |    |
|--|----|
| Figure 2.13: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (44.7 mm) on February 11, 1985 (NG). WSA=control, WSB=BMP, and WSC=no BMPs. ....   | 47 |
| Figure 2.14: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (48.0 mm) on February 2, 1986 (NG). WSA=control, WSB=BMP, and WSC=no BMPs. ....    | 47 |
| Figure 2.15: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.1 mm) on February 26, 1987 (NG). WSA=control, WSB=BMP, and WSC=no BMPs. ....   | 48 |
| Figure 2.16: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (44.7 mm) on August 23, 1988 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....      | 48 |
| Figure 2.17: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.8 mm) on September 30, 1989 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....   | 49 |
| Figure 2.18: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.1 mm) on February 3, 1990 (NG). WSA=control, WSB=BMP, and WSC=no BMPs. ....    | 49 |
| Figure 2.19: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (35.6 mm) on June 25, 1991 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....        | 50 |
| Figure 2.20: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (49.8 mm) on August 27, 1992 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....      | 50 |
| Figure 2.21: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.6 mm) on October 18, 1993 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....     | 51 |
| Figure 2.22: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (42.2 mm, 7.2 hr) on May 13, 2002 (G). WSA=control, WSB=BMP, and WSC=no BMPs. .... | 51 |

|  |    |
|--|----|
| Figure 2.23: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (46.2 mm, 18.2 hr) on November 18, 2003 (NG). WSA=control, WSB=BMP, and WSC=no BMPs..... | 52 |
| Figure 2.24: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (37.3 mm, 3 hr) on June 15, 2004 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....        | 52 |
| Figure 2.25: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (30.5 mm, 6.7 hr) on May 19, 2005 (G). WSA=control, WSB=BMP, and WSC=no BMPs. ....       | 53 |
| Figure 2.26: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (31.5 mm, 10.8 hr) on January 23, 2006 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.....  | 53 |
| Figure 2.27: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (33.3 mm, 9.2 hr) on November 14, 2007 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.....  | 54 |
| Figure 2.28: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (26.4 mm, 9.5 hr) on January 29, 2008 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.....   | 54 |
| Figure 2.29: Yearly baseflow and stormflow WSA, WSB, and WSC (1982-1993). ....   | 64 |
| Figure 2.30: Yearly baseflow and stormflow WSA, WSB, and WSC (2002-2008). ....   | 64 |
| Figure 2.31: Growing season stormflow and baseflow WSA, WSB, and WSC (1982-1993).....  | 66 |
| Figure 2.32: Growing Season baseflow and stormflow WSA, WSB, and WSC (2002-2008).....  | 66 |
| Figure 2.33: Non-growing season baseflow and stormflow WSA, WSB, and WSC (1982-1993).....  | 68 |
| Figure 2.34: Non-growing season baseflow and stormflow WSA, WSB, and WSC (2002-2008).....  | 68 |
| Figure 2.35: Average Storm Peak Flow WSA, WSB, WSC (1982-1993).....  | 73 |

|  |    |
|--|----|
| Figure 2.36: Average Storm Peak Flow WSA, WSB, WSC (2002-2008).....  | 73 |
| Figure 2.37: Growing Season Average Storm Peak Flow WSA, WSB, WSC<br>(1982-1993).....                              | 74 |
| Figure 2.38: Growing Season Average Storm Peak Flow WSA, WSB, WSC<br>(2002-2008).....                              | 74 |
| Figure 2.39: Non-Growing Season Average Storm Peak Flow WSA, WSB, WSC<br>(1982-1993).....                          | 75 |
| Figure 2.40: Non-Growing Season Average Storm Peak Flow WSA, WSB, WSC<br>(2002-2008).....                          | 75 |
| Figure 2.41: Average storm volume as a percentage of rainfall WSA, WSB, WSC<br>(1982-1993).....                    | 78 |
| Figure 2.42: Average storm volume as a percentage of rainfall WSA, WSB, WSC<br>(2002-2008).....                    | 78 |
| Figure 2.43: Growing season average storm volume as a percentage of rainfall<br>WSA, WSB, WSC (1982-1993). ....    | 80 |
| Figure 2.44: Growing season average storm volume as a percentage of rainfall<br>WSA, WSB, WSC (2002-2008). ....    | 80 |
| Figure 2.45: Non-growing season average storm volume as a percentage of<br>rainfall WSA, WSB, WSC (1982-1993)..... | 82 |
| Figure 2.46: Non-growing season average storm volume as a percentage of<br>rainfall WSA, WSB, WSC (2002-2008)..... | 82 |
| Figure 2.47: Average Time to Peak WSA, WSB, WSC (2002-2008).....   | 85 |
| Figure 2.48: Growing season average Time to Peak WSA, WSB, WSC (2002-<br>2008). ....                               | 85 |
| Figure 2.49: Non-growing season average Time to Peak WSA, WSB, WSC<br>(2002-2008).....                             | 86 |
| Figure 2.50: Average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993)....                                  | 89 |
| Figure 2.51: Average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008)....                                  | 89 |
| Figure 2.52: Average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993)...                                  | 90 |
| Figure 2.53: Average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008)...                                  | 90 |

|   |     |
|---|-----|
| Figure 2.54: Growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993).....   | 92  |
| Figure 2.55: Growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008).....   | 92  |
| Figure 2.56: Growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993).....  | 93  |
| Figure 2.57: Growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008).....  | 93  |
| Figure 2.58: Non-growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993).....   | 95  |
| Figure 2.59: Non-growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008).....   | 95  |
| Figure 2.60: Non-growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993).....  | 96  |
| Figure 2.61: Non-growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008).....  | 96  |
| Figure 2.62: Average curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. ....           | 98  |
| Figure 2.63: Average curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. ....          | 98  |
| Figure 2.64: Growing season curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. ....    | 100 |
| Figure 2.65: Growing season curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. ....   | 100 |
| Figure 2.66: Non-growing season curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008..... | 102 |

|  |     |
|--|-----|
| Figure 2.67: Non-growing season curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008..... | 102 |
| Figure 2.68 Percent stand composition in WSA, WSB, and WSC in 2010. ....   | 104 |
| Figure 2.69: Leaf of <i>Liriodendron tulipifera</i> L.-Yellow Poplar (Mohlenbrock, 1995).<br>.....   | 105 |
| Figure 2.70: Leaf of <i>Quercus rubra</i> -Red Oak (Seiberling et al., 2006). ....   | 105 |
| Figure 2.71: Leaf of <i>Acer rubrum</i> -Red Maple (Seiberling et al., 2006). ....   | 106 |
| Figure 2.72: Leaf of <i>Magnolia accuminata</i> -Cucumber Tree (Chamuris, 2016). ....  | 106 |
| Figure 2.73: Leaf of <i>Tilia americana</i> -Basswood (Herman, D.E., et al., 1996)... ..   | 107 |
| Figure 2.74: Leaf of <i>Q. montana</i> -Chestnut Oak (Mohlenbrock, 1995).....  | 107 |
| Figure 2.75: Leaves of <i>Magnolia macrophylla</i> -Bigleaf Magnolia (Profant, 2010).<br>.....   | 108 |
| Figure 2.76: Invasive grass <i>Microstegium</i> sp.-Nepalese browntop (Mohlenbrock, 1991). ....  | 110 |
| Figure 2.77: Invasive <i>Lonicera maackii</i> -Honeysuckle (Herman, D.E., et al., 1996).<br>.....  | 110 |
| Figure 2.78: <i>Ailanthus altissima</i> -Tree of heaven (Goldman, 2012).....   | 111 |
| Figure 2.79: Vegetation height collected by LiDAR in 2013 for WSA, WSB, and WSC. ....  | 113 |
| Figure 2.80: Vegetation height distributions in 2013 in WSA, WSB, and WSC. ....  | 114 |
| Figure 2.81: Stand density in WSA (2015). ....   | 116 |
| Figure 2.82: Stand density and regrowth in WSB (2015).....   | 117 |
| Figure 2.83: Stand density and regrowth in WSC (2015). ....  | 118 |
| Figure 3.1: Sediment and debris buildup behind WSA flume (2016). ....  | 129 |
| Figure 3.2: Sediment and debris buildup behind WSB flume (2016). ....  | 130 |
| Figure 3.3: Sediment and debris buildup behind WSC flume (2016). ....  | 131 |
| Figure 3.4: Yearly calcium FWMC (1982-1993) and (2002-2008). ....  | 141 |
| Figure 3.5: Growing season calcium FWMC (1982-1993) and (2002-2008). ...   | 141 |
| Figure 3.6: Non-growing season calcium FWMC (1982-1993) and (2002-2008).<br>.....  | 141 |

|  |     |
|--|-----|
| Figure 3.7: Yearly potassium FWMC (1982-1993) and (2002-2008). .....               | 142 |
| Figure 3.8: Growing season potassium FWMC (1982-1993) and (2002-2008). .....       | 142 |
| Figure 3.9: Non-growing season potassium FWMC (1982-1993) and (2002-2008). .....   | 142 |
| Figure 3.10: Yearly magnesium FWMC (1982-1993) and (2002-2008). .....              | 144 |
| Figure 3.11: Growing season magnesium FWMC (1982-1993) and (2002-2008). .....      | 144 |
| Figure 3.12: Non-growing season magnesium FWMC (1982-1993) and (2002-2008). .....  | 144 |
| Figure 3.13: Yearly sodium FWMC (1982-1993) and (2002-2008). .....                 | 145 |
| Figure 3.14: Growing season sodium FWMC (1982-1993) and (2002-2008). .....         | 145 |
| Figure 3.15: Non-growing season sodium FWMC (1982-1993) and (2002-2008). .....     | 145 |
| Figure 3.16: Yearly sulfate FWMC (1982-1993) and (2002-2008). .....                | 147 |
| Figure 3.17: Growing season sulfate FWMC (1982-1993) and (2002-2008). .....        | 147 |
| Figure 3.18: Non-growing season sulfate FWMC (1982-1993) and (2002-2008). .....    | 147 |
| Figure 3.19: Yearly nitrate FWMC (1982-1990) and (2002-2008). .....                | 148 |
| Figure 3.20: Growing season nitrate FWMC (1982-1990) and (2002-2008). .....        | 148 |
| Figure 3.21: Non-growing season nitrate FWMC (1982-1990) and (2002-2008). .....    | 148 |
| Figure 3.22: Yearly ammonium FWMC (2002-2008). .....                               | 149 |
| Figure 3.23: Growing season ammonium FWMC (2002-2008). .....                       | 149 |
| Figure 3.24: Non-growing season ammonium FWMC (2002-2008). .....                   | 149 |
| Figure 3.25: Yearly alkalinity FWMC (1982-1993) and (2002-2008). .....             | 151 |
| Figure 3.26: Growing season alkalinity FWMC (1982-1993) and (2002-2008). .....     | 151 |
| Figure 3.27: Non-growing season alkalinity FWMC (1982-1993) and (2002-2008). ..... | 151 |
| Figure 3.28: Yearly phosphate FWMC (1982-1993). .....                              | 153 |
| Figure 3.29: Growing season phosphate FWMC (1982-1993). .....                      | 153 |
| Figure 3.30: Non-growing season phosphate FWMC (1982-1993). .....                  | 153 |



|   |     |
|---|-----|
| Figure 3.31: Yearly chloride FWMC (2002-2008). .....                      | 155 |
| Figure 3.32: Growing season chloride FWMC (2002-2008). .....              | 155 |
| Figure 3.33: Non-growing season chloride FWMC (2002-2008).....            | 155 |
| Figure 3.34: Yearly total organic carbon FWMC (2002-2008). .....          | 156 |
| Figure 3.35: Growing season total organic carbon FWMC (2002-2008).....    | 156 |
| Figure 3.36: Non-growing season total organic carbon FWMC (2002-2008).... | 156 |

## LIST OF TABLES

|  |    |
|--|----|
| Table 2.1: Total, growing, and non-growing season suitable hydrographs 1982-1993 and 2002-2008. ....   | 23 |
| Table 2.2: Total, growing, and non-growing season suitable hydrographs with a cumulative storm depth equal to or greater than 25.4 mm 1982-1993 and 2002-2008. ....  | 25 |
| Table 2.3: Storm event characteristics 1982-1993. ....   | 29 |
| Table 2.4: Storm Event Characteristics 2002-2008. ....   | 35 |
| Table 2.5: Average storm depth $\pm$ Std. dev. (mm), average 5-day antecedent depth $\pm$ std. dev. (mm), maximum storm depth (mm), cumulative storm depth (mm), and sample size 1982-1993 and 2002-2008. .... | 43 |
| Table 2.6: Average storm intensity $\pm$ standard deviation (mm/hr), average maximum intensity $\pm$ standard deviation (mm/hr), and sample size 2002-2008. ....   | 44 |
| Table 2.7: ANCOVA baseflow volume and stormflow volume results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup> ....  | 56 |
| Table 2.8: ANCOVA peak flow ( $Q_p$ ) ( $m^3 s^{-1} ha^{-1} \times 10^{-2}$ ) results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup> ....   | 57 |
| Table 2.9: ANCOVA storm volume as a (%) of rainfall results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup> ....   | 58 |
| Table 2.10: ANCOVA time to peak ( $T_p$ ) results for WSA, WSB, and WSC 2002-2008. <sup>1</sup> ....   | 60 |
| Table 2.11: ANCOVA CN ( $\lambda=0.2$ & $\lambda=0.05$ ) results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup> ....  | 61 |
| Table 2.12: Yearly WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area <sup>1</sup> 1982-1993 and 2002-2008.....   | 63 |
| Table 2.13: Growing season WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area <sup>1</sup> 1982-1993 and 2002-2008.....   | 65 |
| Table 2.14: Non-growing season WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area <sup>1</sup> 1982-1993 and 2002-2008.....   | 67 |

|   |     |
|---|-----|
| Table 2.15: WSA, WSB, and WSC average yearly, growing, and non-growing season peak flow ( $Q_p$ ) $\pm$ standard deviation ( $m^3 ha^{-1} s^{-1} \times 10^{-2}$ ) normalized by drainage area <sup>1</sup> 1982-1993 and 2002-2008. .... | 72  |
| Table 2.16: Yearly average storm volume as a (%) of rainfall $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.....  | 77  |
| Table 2.17: Growing season average storm volume as a (%) of rainfall $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008. ....   | 79  |
| Table 2.18: Non-growing season average storm volume as a (%) of rainfall $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008. ....   | 81  |
| Table 2.19: WSA, WSB, and WSC average time to peak ( $T_p$ ) $\pm$ standard deviation (hr) 2002-2008. ....  | 84  |
| Table 2.20: Yearly curve numbers (CNs) $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008. ....   | 88  |
| Table 2.21: Growing season curve numbers (CNs) $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.....  | 91  |
| Table 2.22: Non-growing season curve numbers (CNs) $\pm$ standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.....  | 94  |
| Table 2.23: Average curve numbers (CNs) $\pm$ standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.....             | 97  |
| Table 2.24: Growing season curve numbers (CNs) $\pm$ standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. ....     | 99  |
| Table 2.25: Non-growing season curve numbers (CNs) $\pm$ standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008. .... | 101 |
| Table 2.26: Vegetation height distribution in WSA, WSB, and WSC. ....   | 115 |
| Table 3.1: ANOVA time interaction results, pre- and post-harvest concentration ( $mg L^{-1}$ ) plus sample size (pre)x(post). <sup>1</sup> .....  | 133 |

|  |     |
|--|-----|
| Table 3.2: ANOVA treatment interaction results, concentration ( $\text{mg L}^{-1}$ ) plus sample. <sup>1</sup> size (pre)x(post).....                            | 135 |
| Table 3.3: ANOVA time*treatment interaction results, pre and post-harvest concentration ( $\text{mg L}^{-1}$ ) plus sample size (pre)x(post). <sup>1</sup> ..... | 137 |
| Table 3.4: Water quality regression analysis, slope ( $\text{mg L}^{-1} \text{ d}^{-1} \times 10^{-4}$ ), <i>p</i> -value, and sample size. <sup>1</sup> .....   | 139 |

## **CHAPTER 1: INTRODUCTION**

### **1.1 INTRODUCTION**

Riparian zones, also known as streamside zones, provide a buffer between upland management scenarios and the stream. The importance of riparian zones in maintaining hydrologic function, filtering upland derived sediments, utilizing upland derived nutrients, maintaining in-stream and near-stream temperature profiles, and providing habitat and corridors for aquatic and terrestrial fauna has been identified but not well quantified, particularly for headwater stream systems (NCASI, 1999a). Because of their small size, headwater streams are quite sensitive to anthropogenic disturbances such as harvesting which can cause larger runoff volumes, higher peak flows, and decreased water quality (Richardson and Danehy, 2007; Reeves, 2012). Most forestry best management practices (BMPs) are designed to decrease sediment transport resulting from soil erosion. Soil erosion and subsequent suspended sediment in surface waters was considered by many as one of the largest environmental concerns in the U.S. during the 1980's and is still a major concern today (USEPA, 1992). Erosion of organic and nutrient rich surface soils also decreases forest productivity (Pritchett and Fisher, 1987). Transport of sediment to streams and subsequent sedimentation leads to loss of stream habitat and altered stream hydrology (NCASI, 1999a; 1999b). Although sediment transport is a main concern associated with forest harvesting, nutrient transport and impacts on stream temperature and carbon distribution are also important (Hornbeck and Edwards, 1990; Arthur et al., 1998). While research has examined the short term effectiveness of forestry BMPs (Arthur et al., 1998; McClure et al., 2004; Witt et al., 2011), research on the long-term effects on these BMPs is quite limited (Burt et al., 2015). As such, a thorough investigation into the long-term effectiveness of forestry BMP use is needed, particularly with respect to riparian buffers, to provide a framework of understanding of how such BMPs can protect headwater streams systems thus leading to better forestry management practices.

In 1982, a study was initiated to evaluate BMP effectiveness on three headwater catchments following logging. The headwater catchments are located in the Field Branch watershed in the University of Kentucky's Robinson Forest, which is a teaching, research and extension experimental forest located in the Cumberland Plateau of eastern Kentucky. Treatments included a control or reference catchment (WSA), a catchment with BMPs implemented including a 15.2 m buffer along the perennial stream segment and others pertaining to road and trail development and retirement (WSB), and a catchment that was harvested completely to the stream's banks without the use of BMPs (i.e. logger's choice) (WSC). An overview of initial results by Arthur et al. (1998) concluded that the harvested sites recovered within approximately four years after harvest and that WSB performed marginally better than WSC at reducing water yield. With regards to water quality, the streamside buffer in WSB limited  $\text{NO}_3$  export but the benefits of forestry BMPs at limiting the other nutrient fluxes in WSB was marginal. The results regarding sediment loading that were found in the Arthur et al. (1998) study concluded that WSB and WSC were significantly higher than WSA from 1984-1986, WSC was significantly higher than WSA in 1988, and in 1990 all three watersheds were significantly different from each other with WSC being the highest and WSA being the lowest. Although the preliminary conclusions were intriguing from a BMP effectiveness standpoint, little is known about the long-term significance that BMP implementation may have on water yield and quality.

## **1.2 OBJECTIVES**

The main goal of this project was to evaluate the effectiveness of forestry BMPs on long-term hydrology and water quality characteristics by comparing three forested Appalachian headwater catchments before and after logging. Three treatments were implemented and include: control (WSA), BMPs implemented during and after harvest (WSB), and clear-cut to the stream (i.e., loggers' choice) with no BMPs (WSC). The objectives of the study were:

1. Evaluate yearly, growing, and non-growing season baseflow volume, stormflow volume, peak flow, stormflow volume as a percentage of rainfall, time to peak, and curve number on the paired watersheds to determine significant pairwise differences and long-term trends for the monitored time periods (Chapter 2).
2. Analyze the monitored pollutants on the paired watersheds by comparing yearly, growing, and non-growing season concentrations to determine significant pairwise differences and long-term trends for the monitored time periods (Chapter 3).

### **1.3 ORGANIZATION OF THESIS**

Chapter 1 contains an overview of the research problems and objectives. Chapters 2 and 3 present detailed descriptions of the work performed to satisfy the research objectives. Chapter 4 discusses the conclusions of the research while Chapter 5 looks into opportunities for future work.

## **CHAPTER 2: EFFECT OF BEST MANAGEMENT PRACTICES ON THE HYDROLOGY OF THREE APPALACHIAN HEADWATER CATCHMENTS**

### **2.1 INTRODUCTION**

Harvesting has the potential to significantly alter the hydrologic response of a watershed. Reduced vegetative cover often means reduced evapotranspiration (ET) rates, higher peak flows, quicker response times, and greater storm runoff volumes. A review of 94 catchment experiments found that every experiment except one observed significant increases in water yield and decreases in ET rates with decreased vegetative cover (Bosch and Hewlett 1982). An experiment detailing the hydrologic changes following clear-cutting on a southern Appalachian catchment noted similar results as well as a significant increase in stormflow volumes and higher peak flows (Swank et al., 2001). One means of reducing the impacts of harvesting on watershed hydrology is through the use of best management practices (BMPs).

Forestry BMPs are employed primarily to reduce erosion and subsequent suspended sediment from reaching streams but other goals may include: preserving wildlife and stand characteristics, aesthetics, recreation, and promoting water quality. Although the primary goal of forestry BMPs is to reduce erosion, studies have shown that altered forest hydrology due to harvesting may play a significant role in determining sediment transport rates (Troendle and Olsen, 1993; Arthur et al., 1998). The USDA Forest Service has a complete guide on designing and implementing BMPs on national forested lands while Kentucky's Division of Forestry has their own region specific guide (Stringer, 2001; USDA Forest Service, 2012). Common forestry BMPs that are detailed in these guides include: the use of streamside management zones (SMZs), access road construction and subsequent seeding, stream crossing improvements, and reestablishing vegetation on disturbed sites.

A number of studies throughout the United States have examined the role of BMPs in mitigating the negative impacts of harvesting a forested watershed. However, much of this work has focused on water chemistry and less on the



effectiveness of forestry BMPs for minimizing changes in watershed hydrology. A review of 81 different BMP effectiveness studies revealed that only five examined impacts to watershed hydrology (Patric, 1980; Keppeler and Ziemer, 1990; Lynch and Corbett, 1990; Arthur et al., 1998; Keppeler et al., 2008; Stednic, 2008; Cristan et al., 2016). Of those five studies, only two, Patric (1980) and Arthur et al. (1998) are located in the Appalachian Plateaus physiographic province.

In an eastern Kentucky study, Arthur et al. (1998) concluded that forestry BMPs provided some marginal benefit over an adjacent clear-cut watershed in reducing water yield post-harvest, but neither watershed had returned to pre-harvest conditions at the conclusion of the study eight years post-harvest. In a West Virginia study, results indicated that the use of a 20 m SMZ was effective at limiting significant increases in water yield following clear-cutting and that water yield nearly returned to pre-harvest conditions five years post-harvest (Patric, 1980). One additional study, located in the Allegheny Mountains, found that annual peak flows and total annual stormflow volumes were unchanged from pre-harvest to post-harvest with BMP implementation, but significant increases in peak flow and stormflow volumes were observed during the growing season for six years post-harvest (Kochenderfer et al., 1997). Barring the initial study by Arthur et al. (1998), neither Patric (1980) nor Kochenderfer et al. (1997) utilized a paired watershed approach to determine the performance of BMPs on watershed hydrology as compared to a clear-cut watershed and a control. This is important because a secondary conclusion that was reached by Bosch and Hewlett (1982) was that the most reliable results from catchment experiments were developed on studies that utilized the paired watershed approach.

The Wagon Wheel Gap Study, initiated in Colorado in 1909, is widely recognized as the first paired watershed study in the U.S. where separate treatment and control watersheds were used to examine the hydrologic effects of forest cutting (Hewlett et al., 1969; Bosch and Hewlett, 1982; Stednick, 1996; Ice and Stednick, 2004). The paired watershed approach to hydrologic study involves the use of a control watershed and one or more treatment watersheds (Cherry, 2006; Brooks et al., 2012; Witt, 2012). Initially, before treatments are

implemented, each watershed is monitored for a specific period of time during which data for all desired parameters are collected. This is referred to as the calibration period. These data are then used to develop a regression relationship between the control watershed and each individual treatment watershed through a mass balance approach. Once a suitable relationship has been developed, treatments are carried out and monitoring continues. Data from this post treatment period are then used to develop post treatment regression relationships among control and treatment watersheds. Treatment effects and their magnitude are detected by examining differences in slopes and y-intercepts between calibration and post treatment regressions. In order for this method to be effective, it is important that control and treatment watersheds are as similar as possible in terms of watershed area, location, aspect, vegetative cover, soil types, geologic composition and topography, and that there are no deep seepages into or out of any of the watersheds so that differences detected throughout the post-treatment period are confidently attributed to the treatments alone (Borman and Likens, 1979; Cherry, 2006; Witt, 2012).

As the relationship between forest cutting, water yield, and evapotranspiration became accepted amongst forest scientists, it became apparent that large variability in results from hydrologic studies were observed from one site to another (Hibbert, 1967; Bosch and Hewlett, 1982; Witt, 2012). While it was known that forest removal could generate increases in water yield, it was unclear as to what the threshold level of removal was. Stednick (1996) reported that the threshold of harvested area needed to generate a streamflow response in various regions of the United States was between 15 – 50% (Stednick, 1996; Wei and Zhang, 2010). In the Appalachian Mountains, the minimum harvested area necessary to produce a measureable streamflow response was found to be 20% (Stednick, 1996). However, these suggested levels were shown not to hold true in all instances. For example, Adams et al. (2004) reported that at least 25% of stand basal area must be removed in order to elicit a hydrologic response at the Fernow Experimental Forest in West Virginia. Moreover, the minimum threshold has been reported to be as low as

10% at the Coweeta Hydrologic Laboratory in North Carolina (Stednick, 1996; Swank and Crossley, 2012).

This level of variability within the Appalachian region is primarily a reflection of the complexity of climatic, geographic and biotic factors it presents. The mixed mesophytic forest that covers much of Appalachia and most of eastern Kentucky is a complex mixture of different forest types and is considered by many to be one of the most diverse (in terms of flora) ecosystems in the United States (Moore et al., 2005). This mosaic of forest types is controlled by numerous factors including elevation, aspect, geology, land use history, and the species composition of the surroundings. The effect that forest management/manipulation may have on hydrology is directly tied to interactions among all of these factors. For example, Douglas (1983) showed that first year hydrologic response to forest cutting in Appalachian forests is determined mainly by the amount of basal area removed and the amount of solar radiation received at the site, which is a function of catchment aspect. Hewlett and Hibbert (1967) observed an almost three-fold difference in water yield between north and south facing catchments after harvest at the Coweeta Hydrologic Laboratory, with north facing catchments being the greater of the two (Douglas, 1983). Further, it has been shown that management actions that result in changes in species composition and tree ecophysiology may alter hydrologic processes and evapotranspiration through changes in canopy interception and transpiration (Stoy et al., 2006; Ford et al., 2011; Vose et al., 2011). Lu et al. (2003) found that the spatial variability of regional actual evapotranspiration in the southeastern U.S. was best explained by a multivariate linear regression model using precipitation, latitude, elevation, and percentage of forest cover as independent variables. However, over the long term, Zhang et al. (2004) found that under the same climatic conditions average annual evapotranspiration was determined mainly by the vegetative characteristics of a watershed and how the species present use available soil water.

Another important aspect of forest hydrology is recovery time. That is, the time needed for a watershed to resume a hydrologic regime which is statistically

similar to that observed prior to disturbance (Hibbert and Gottfried, 1987; Stednick and Kern, 1992; Hornbeck et al., 1993; Stednick, 1996; Stednick, 2008). In a meta-analysis of various hydrologic databases from throughout the southeastern U.S., Sun et al. (2005) found that forest removal in the southeast will increase water yield, with the greatest increases occurring in areas with greater precipitation. Furthermore, Appalachian watersheds could experience the greatest water yield increases after harvest due to their high precipitation input, low temperatures, and forest cover characteristics. Sun et al. (2005) also stated that hydrologic recovery time for hilly upland systems is expected to be significantly longer than for other systems in the Southeast. Long-term data from Coweeta Hydrologic Laboratory in North Carolina showed that the longest recovery times were observed in high elevation and north facing watersheds, where either stand species conversion or coppicing treatments have been repeatedly implemented (Ford et al., 2011). As of 2008, neither of two species conversion watersheds or the coppicing watershed had fully recovered since their latest treatment in 1956-57 and 1962, respectively. Ford et al. (2011) proposed that the unexpected level of streamflow increase from the coppicing treatment at Coweeta may be due to increased competition among individual stems reducing leaf area index, thereby reducing evapotranspiration.

Species conversions resulting in changes to leaf area index can significantly increase or decrease streamflow over the long term, depending on the direction of change in leaf area index (Ford et al., 2011). Higher leaf areas indices are associated with decreased streamflow while lower leaf areas indices are associated with increased streamflow. Wullschleger et al. (2001) found that diffuse porous hardwood species generally contain a larger sapwood area within their stems than ring porous hardwoods of similar stem diameter. A number of studies have observed that differences in stomatal conductance between xylem functional groups (i.e. ring porous vs. diffuse porous) in Appalachian forests may be great (Wallace, 1988; Wullschleger et al., 2001; Vose, 2007; Ford et al., 2010; Ford et al., 2011). The implications of these findings were stated eloquently by Wullschleger et al. (2001):

*“Whenever a forest is composed of both ring porous and diffuse porous species, total transpiration is likely to be dominated not by the species with the largest basal area, nor by the species present in the greatest number, but rather by the species with the largest sapwood area.”*

The hydrologic recovery time of disturbed watersheds in Appalachia is dictated by numerous parameters. Douglas and Swank (1972) produced a regression model relating first year water yield increases to percentage of basal area cut using data from all known forest harvesting experiments in the Appalachian Highland Physiographic Division (Douglas, 1983). The results of this regression model indicated that hydrologic recovery time in this region is 1.57 years for each inch of water yield recorded in the first year after harvest. In light of the limited amount of research on the long-term effects of forestry BMPs on watershed hydrology, the objective of this study was to evaluate yearly, growing, and non-growing season baseflow volume, stormflow volume, peak flow, stormflow volume as a percentage of rainfall, time to peak, and curve number on the paired watersheds to examine significant pairwise differences and long-term trends for the monitored time periods

## **2.2 METHODS**

### **2.2.1 Study Site**

The research was conducted at the University of Kentucky’s Robinson Forest, which is located in southeastern Kentucky in the Cumberland Plateau section of the Appalachian Plateaus province of the Appalachian Highlands (latitude 37° 27.01 N; longitude 83° 11.43 W) (Figure 2.1). Robinson Forest is approximately 6,000 ha in size with eight discontinuous properties: the main block where this research takes place is 4,200 ha. Robinson Forest is situated in the Appalachian mixed mesophytic forest region, which is characterized by high hills and low valleys with elevations ranging from 385 to 610 m (1,270 to 2,000 ft).




Figure 2.1: Location of University of Kentucky's Robinson Forest labeled in green.

These mixed mesophytic forests occur on moist and topographically protected areas within highly dissected hills and mountains. These forests are a part of a transition zone from oak-hickory forest to the northern hardwood forest and are among the most diverse in the United States with more than 30 canopy tree species (Moore et al., 2005). Robinson Forest is characterized by steep side-slopes ( $\mu=45\%$ ) and a hydrologically restrictive geologic substrate consisting of interbedded sandstone, siltstone, shale, and coal (McDowell et al., 1981). Robinson Forest was clear-cut in the early 1900's then donated by the Mowbray-Robinson lumber company to the University of Kentucky as a teaching, research, and extension experimental forest (Figure 2.2).

The study site where this research takes place is just a small portion of Robinson Forest and consists of three adjacent subcatchments (WSA, WSB, and WSC) all located in the Field Branch watershed (Figures 2.3 and 2.4). The drainage areas of WSA, WSB and WSC are 16.1, 11.2, and 10.5 ha, respectively. At the start of the study in 1982, the overstory of WSB and WSC was dominated by oaks (*Quercus spp.*; 39%), hickories (*Carya spp.*; 17%), and yellow poplar (*Liriodendron tulipifera L.*; 15%) while WSA's overstory consisted predominately of yellow poplar (Overstreet, 1984).



Figure 2.2: Mowbray-Robinson logging operations at Robinson station in Quicksand Kentucky.






Figure 2.3: Robinson Forest hydrologic features and topography. The study area is N. Field Branch (WSA); W. Field Branch (WSB); and S. Field Branch (WSC).




Figure 2.4: Digital elevation model of study site with stream network shown in yellow, WSA shown in blue, WSB in red, and WSC in green.

For this study, a paired watershed approach was used with WSA serving as the uncut control, WSB was clear-cut with BMP implementation, and WSC was clear-cut to the stream with no BMP implementation.

The BMPs that were implemented in WSB included:

1. 15.2 m (50 ft) undisturbed riparian buffer on both sides of the perennial stream
2. Logging roads seeded with fescue after logging was complete
3. Logging roads constructed on less than 10% grade
4. Log skidder kept on roads
5. Broad-based dips used as water control structures on logging roads
6. Logging debris was kept out of the stream

The management strategies that were implemented at WSC include:

1. No intact riparian buffer on either side of the perennial stream
2. Roads left bare after logging was complete
3. Logging roads constructed on more than 10% grade
4. Logs repeatedly skidded downhill
5. No water control structures were used on logging roads
6. Trees felled into and across the stream

Hydrologic monitoring for the three subcatchments began in February of 1982, and following a 19-month calibration period, WSB and WSC were harvested (Figure 2.5). Harvesting at both sites resulted in a complete silvicultural clear-cut with commercial timbers (> 35.5 cm diameter at breast height (dbh)) being removed from the site and all stems < 5 cm dbh cut and left on site (Arthur et al., 1998). Since harvest, the subcatchments have experienced no further harvesting or modifications and visibly larger trees surround the riparian buffer in WSB compared to WSC in 2005, 22 years post-harvest (Figure 2.6). Hydrologic monitoring has continued post-harvest until the present day.




Figure 2.5: Study site post-harvest (1984) with logging roads shown in brown: WSA is on the far left, WSB is in the middle along with the riparian buffer that was left along the stream, and WSC is on the right.






Figure 2.6: Study site 22 years after harvest (2005) with WSA on the far left, WSB in the middle, and WSC on the right.

### 2.2.2 Precipitation Data

Precipitation data (P) were collected by weather stations located in Little Millseat and at Camp (Figure 2.3). The Little Millseat weather station is located below the confluence of the three watersheds (latitude 37° 28.53 N; longitude 83° 09.63 W) and the Camp weather station is located in a nearby hollow (latitude 37° 27.01 N; longitude 83° 11.43 W). Daily cumulative precipitation data at the weather stations were collected using tipping bucket recording gauges and weighing gauges. At the Camp site, a Campbell Scientific weather station was installed in 2000 which allowed for the collection of 15-minute cumulative precipitation data as well as relative humidity, temperature, wind speed and direction, and solar radiation. During the first half of the study (1982-1993), precipitation data from the Little Millseat weather station were used in the hydrologic analyses. The Camp weather station data was used during the second half of the study (2002-2008) because of its ability to record 15-min precipitation data. The Camp weather station is located about 3 km south of the Little Millseat weather station. Data from both weather stations were unavailable in 1999 and 2000 and were substituted with data from nearby ( $\approx 20$  Km) Jackson County Airport (latitude 37° 59.36 N; longitude 83° 31.73 W). These data were only used to help determine the 27-year study period average annual precipitation characteristics.

Storms events were characterized by cumulative precipitation (rainfall) depths greater than or equal to 11 mm. The 11 mm value was chosen in accordance with the Revised Universal Soil Loss Equation (RUSLE), which specifies that erosion does not occur during rainfall events less than 12.7 mm ( $\frac{1}{2}$  in). Reducing this value to 11 mm increased overall sample size and was utilized in previous storm characteristic and hydrograph studies in eastern Kentucky (Warner et al., 2010b; Blackburn-Lynch, 2015). Storm events were considered single if the time gap between recorded rainfall exceeded three hours (Warner et al., 2010b; Blackburn-Lynch, 2015). For instances when only daily cumulative precipitation data were available, unit runoff depths and durations were computed for each hydrograph and compared to daily precipitation depths to

determine if a storm event occurred over multiple days. Consecutive days with precipitation that fell during the course of the hydrograph response were included in the total storm depth. The only time this was not the case was if a spike in daily precipitation occurred past the time of peak hydrograph flow. Hydrograph response would dictate whether this spike in precipitation was two separate storm events (multiple peaks) or a single storm event (single peak) (Figures 2.7 and 2.8). If possible, hydrographs were separated into their respective storm events; otherwise, they were placed into the unsuitable hydrograph category (Figure 2.8). Each storm depth was compared against each respective unit hydrograph depth to confirm that flow out was not greater than flow in and to ensure all attributable precipitation was recorded with storm depth.




Figure 2.7: Example of baseflow separation and a suitable (single peak) hydrograph for analysis




Figure 2.8: Example of an unsuitable hydrograph (multiple peaks).



### 2.2.3 Hydrograph Data

Stage data for WSA, WSB and WSC were each recorded using an H-flume equipped with a stage-height recorder. WSA was equipped with a .91 m (3 ft) tall H-flume while WSB and WSC were both equipped with a .76 m (2.5 ft) tall H-flume. Equations were then formulated to detail the stage-discharge relationship for the flumes. To ensure accuracy for all flow regimes, the stage-discharge relationships for the flumes were subdivided into a low-flow regime and a high-flow regime. Low-flows were defined as follows: flows that occurred during a stage less than 0.28 m in the 0.91 m tall flume and 0.20 m in the 0.76 m tall flume. High-flows were defined as everything above the low-flow stages up to the tops of the flumes. The resulting stage-discharge equations are as follows (stage denoted as  $h$  (m), discharge denoted as  $Q$  ( $\text{m}^3 \text{s}^{-1}$ )):

$$Q = 0.6h^2 + 0.02h \quad \text{Equation 2.1}$$

$$Q = 1.16h^2 - 0.14h \quad \text{Equation 2.2}$$

$$Q = 0.6h^2 + 0.01h \quad \text{Equation 2.3}$$

$$Q = 1.06h^2 - 0.1h \quad \text{Equation 2.4}$$

The low-flow and high-flow stage-discharge relationships for WSA are shown in equations 2.1 and 2.2, respectively. The low-flow and high-flow stage-discharge relationships for both WSB and WSC are shown in equations 2.3 and 2.4, respectively. The coefficients of determination ( $R^2$ ) for equations 2.1-2.4 were all  $R^2=0.99$  or greater, indicating a very good fit between stage and discharge. From the stage-discharge relationship, it was determined that the maximum flow that the flume in WSA could accommodate was  $0.83 \text{ m}^3 \text{s}^{-1}$  ( $29.4 \text{ ft}^3 \text{s}^{-1}$ ) and in WSB and WSC it was  $0.52 \text{ m}^3 \text{s}^{-1}$  ( $18.5 \text{ ft}^3 \text{s}^{-1}$ ). During the analyzation periods, the respective flume stage-discharge relationships were used to check that overtopping of the flumes did not occur. Data were collected on a flow-weighted basis with a minimum recording increment of 15 minutes during rapidly varied flow and a maximum recording increment of 1 day during minimally varied flow. Periodic equipment failure at the three flumes resulted in data gaps from 1994-

2001 for WSA and WSB and from 1994-2004 for WSC. Equipment failure was largely an effect of periodic flooding, which resulted in broken stilling wells and displaced and damaged stream-gauging equipment. High sediment loads associated with the floods also periodically filled the flumes and, on occasion, cracked flume foundations resulting in leakage (Figures 3.2 and 3.3).

Storm hydrographs were separated into base flow and storm flow using Purdue's Web-based Hydrograph Analysis Tool (WHAT) (Figure 2.7). WHAT is a recursive digital filter which uses hydrogeological specific constants and a filter algorithm to partition streamflow into baseflow and stormflow (Eckhardt, 2005). As noted by Eckhardt (2005), this type of filter helps reduce subjectivity by limiting the influence of human judgment, is easily reproducible, and can be automated for large data sets. The one subjective element in this technique is the maximum baseflow index ( $BFI_{max}$ ) parameter, which is the ratio of baseflow to total flow used by the algorithm. The  $BFI_{max}$  parameter varies by hydrogeologic region. A study of the United States by Santhi et al. (2008) recommended a baseflow index of 0.2-0.4 for the hydrogeologic region in which this study is located. WHAT recommended a  $BFI_{max}$  of 0.25 for catchments with a perennial stream and hard rock aquifers, which fits the study area, and thus this was the value used. Currently, the only way that the  $BFI_{max}$  parameter can be accurately selected is through tracer-based experiments. However, a sensitivity analysis was conducted on the  $BFI_{max}$  parameter, and it was found that an incorrectly selected  $BFI_{max}$  value led to little relative error as long as the hydrogeologic condition was accurately selected (Eckhardt, 2012).

Over the study period (1982-1993 and 2002-2008), approximately 555 hydrographs from each subcatchment (WSA, WSB, and WSC) were recorded. All hydrographs, including multiple peak hydrographs, were used to calculate annual, annual growing, and annual non-growing season baseflow volumes; storm flow volumes; and total flow (baseflow + stormflow) volumes (Figure 2.8). The growing season dates were defined by the Natural Resource Conservation Service (NRCS) soil survey for Breathitt county and ran from April 20-October 26 (Hayes, 1998). For suitable (e.g. single peak) hydrographs (Figure 2.7), the

following characteristics were computed and statistically analyzed: stormflow volume, baseflow volume, peak flow ( $Q_p$ ), stormflow volume as a percentage of rainfall ( $[S/P] \%$ ), time to peak ( $T_p$ ) (only if 15-min rainfall data were available), and curve number (CN). Time to peak was defined as the time difference between peak hydrograph discharge and the start of rainfall. Subcatchment hydrograph characteristics that were suitable for analysis across all watersheds, 190 hydrographs from the first time period and 104 hydrographs from the second, were evaluated on an annual, annual growing season, and annual non-growing season basis (Table 2.1).

Table 2.1: Total, growing, and non-growing season suitable hydrographs 1982-1993 and 2002-2008.

| Period       | Year              | Total | Growing | Non-growing |
|--------------|-------------------|-------|---------|-------------|
| Pre-harvest  | 1982              | 15    | 7       | 8           |
| Pre-harvest  | 1983              | 16    | 11      | 5           |
| Post-harvest | 1984              | 19    | 9       | 10          |
| Post-harvest | 1985              | 15    | 11      | 4           |
| Post-harvest | 1986              | 11    | 6       | 5           |
| Post-harvest | 1987              | 17    | 8       | 9           |
| Post-harvest | 1988              | 11    | 5       | 6           |
| Post-harvest | 1989              | 18    | 11      | 7           |
| Post-harvest | 1990              | 20    | 10      | 10          |
| Post-harvest | 1991              | 22    | 11      | 11          |
| Post-harvest | 1992              | 15    | 12      | 3           |
| Post-harvest | 1993              | 11    | 10      | 1           |
| Post-harvest | 2002 <sup>1</sup> | 15    | 8       | 7           |
| Post-harvest | 2003 <sup>1</sup> | 16    | 8       | 8           |
| Post-harvest | 2004 <sup>1</sup> | 26    | 16      | 10          |
| Post-harvest | 2005              | 15    | 13      | 2           |
| Post-harvest | 2006              | 11    | 8       | 3           |
| Post-harvest | 2007              | 12    | 6       | 6           |
| Post-harvest | 2008              | 9     | 3       | 6           |

<sup>1</sup> WSC was excluded from analysis from 2002-2004

### 2.2.3.1 Curve Number Method

The CN method was developed by the United States Soil Conservation Service (SCS), presently NRCS, to determine runoff depth and peak flow for different size storm events (USDA-SCS, 1972; USDA-NRCS 2004). The CN method is primarily a planning and design tool that enables professionals to determine peak flows and runoff depths for selected storm sizes at a specific site. In this capacity, CN is selected based on land-use, hydrologic condition, and hydrologic soil group, and ranges from 30 to 100. Higher CNs indicates higher runoff potential while lower CNs indicates lower runoff potential. In this study, however, runoff depth along with precipitation depth were known allowing for direct calculation of CN through equation 2.8 and 2.9 (Hawkins, 1973). Results from Hawkins (1993) found a CN of 85 for Little Millseat watershed in Robinson forest; Little Millseat is located directly north and adjacent to WSA (Figure 2.3).

$$Q = \frac{(P - I_a)^2}{(P - I_a) + S} \quad \text{Equation 2.5}$$

$$I_a = \lambda S \quad \text{Equation 2.6}$$

$$Q = \frac{(P - 0.2S)^2}{(P + 0.8S)} \quad \text{Equation 2.7}$$

$$S = \frac{1000}{CN} - 10 \quad \text{Equation 2.8}$$

$$S = 5[P + 2Q - (4Q^2 + 5PQ)^{0.5}] \quad \text{Equation 2.9}$$

The variable Q is the runoff depth (in), P is the precipitation depth (in), S is the potential maximum retention after runoff begins,  $\lambda$  is the initial abstraction ratio, and  $I_a$  is the initial abstraction. Initial abstraction represents all the losses (i.e. water retained by surface depressions, interception, evaporation, and infiltration) before runoff occurs. The initial abstraction ratio is commonly chosen to be  $\lambda=0.2$  although Hawkins et al. (2002) has shown that this provides an over-approximation of the CN and that  $\lambda=0.05$  is more suitable. CNs in this study were computed using initial abstraction ratios of  $\lambda=0.2$  and  $\lambda=0.05$ . Rainfall events with a cumulative depth less than 25.4 mm (1 in) have also been shown to over

approximate the CN (Hawkins et al., 2002; Warner et al., 2010b). Because of this, CN analysis was separated into all suitable hydrograph events (Table 2.1), and all suitable hydrograph events with a cumulative depth equal to or greater than 25.4 mm (Table 2.2). For the latter, analysis was broken into a pre-harvest period (1982 and 1983) and four post-harvest periods (1984-1988, 1989-1993, 2002-2004, and 2005-2008). This was due to the limited number of suitable events associated with a cumulative depth equal to or greater than 25.4 mm. Statistical analysis of CN was only completed for all suitable events (Table 2.1) to allow for larger yearly sample sizes and to provide a more robust analysis.

Table 2.2: Total, growing, and non-growing season suitable hydrographs with a cumulative storm depth equal to or greater than 25.4 mm 1982-1993 and 2002-2008.

| Period       | Year                   | Total | Growing | Non-growing |
|--------------|------------------------|-------|---------|-------------|
| Pre-harvest  | 1982-1983              | 16    | 11      | 5           |
| Post-harvest | 1984-1988              | 48    | 14      | 24          |
| Post-harvest | 1989-1993              | 55    | 26      | 29          |
| Post-harvest | 2002-2004 <sup>1</sup> | 26    | 15      | 11          |
| Post-harvest | 2005-2008              | 12    | 7       | 5           |

<sup>1</sup> WSC was excluded from analysis from 2002-2004

#### 2.2.4 Forest Stand Analysis

A stand survey was performed in the fall of 2010 to determine the species composition and stand density of the study area. Using ArcMap® software, a systematic random plot distribution was created which resulted in 25 plots per watershed all of which were greater than 20 m apart. Inventory plots were 10 m x 10 m with one ground cover plot in each plot corner measuring 1 m x 1 m. In each inventory plot, all trees with dbh  $\geq$  5 cm were recorded by species, diameter and total height. Tree heights were measured using a Suunto® clinometer. Tree diameters were measured using a Forestry Suppliers diameter measuring tape. Each ground cover plot was assigned a ground cover

percentage using a 1 m x 1 m grid placed within each corner of inventory plots. Throughout the inventory process, any invasive species were noted.

Light detection and ranging (LiDAR) technology was used to further evaluate stand structure. Three LiDAR datasets were used in the analysis. One dataset was low density ( $\sim 1 \text{ pt m}^{-2}$ ) collected in the spring of 2013 during the leaf-off season for the purpose of acquiring terrain information, as a part of a statewide elevation data acquiring program from the Kentucky Division of Geographic Information. The second dataset was high density ( $\sim 40 \text{ pt m}^{-2}$ ) collected in the summer of 2013 during the leaf-on season for the purpose of collecting detailed vegetation information. Raw LiDAR datasets were processed using the TerraScan software (Terrasolid Ltd., 2012) to classify LiDAR points into ground and non-ground points. A third dataset was also created by combining both low-density and high-density points. For each of the three LiDAR datasets (low-density, high-density, and combined), the “Create LAS Dataset” tool in ArcMap 10.2 was used to create a LAS dataset file. The LAS dataset was then filtered to include ground points only, and the “LAS dataset to Raster” tool in ArcMap was used to create a 1 m resolution DEM using the natural neighbor as a fill void method. Four DEMs for each dataset were created considering different interpolation methods: average, inverse distance weighted, minimum, and nearest neighbor. As a result, a total of 12 DEMs covering the study areas were created: three LiDAR datasets and four interpolation methods.

### **2.2.5 Statistical Analysis**

Due to periodic data gaps related to equipment failure, the study’s post-harvest phase was separated into two different time periods: 1984-1993 and 2002-2008. In the second time period, WSC was analyzed from (2005-2008) due to complications with the stage-height recorder from 2002-2004. Massive flooding in May of 2009 compromised flume integrity at all three subcatchments, and in 2010 pressure transducers were installed all three site thereby replacing the stage-height recorders. Large inaccuracies were observed in the data recorded by the pressure transducers from 2010-present and inclusion of these data in this

study were unfeasible. The reason for these inaccuracies was thought to be a result of sediment accumulating in the stilling wells.

One-way analyses of variance (ANOVAs) were used to evaluate annual, pre-harvest and post-harvest storm characteristic differences. Differences were assessed on an annual basis so comparisons could be made between yearly storm characteristics and hydrograph characteristics. Differences in storm characteristics were assessed on a pre-harvest and post-harvest basis to test if statistically significant differences between the two periods existed. Significant differences ( $p \leq 0.05$ ) between annual storm characteristics and the study period average storm characteristic were evaluated using Dunn's multiple comparison tests. Significant differences ( $p \leq 0.05$ ) between pre-harvest and post-harvest storm characteristics were also evaluated using Dunn's multiple comparison tests. Storm characteristics evaluated were storm depth (mm), 5-day antecedent moisture (mm) (1982-1993 and 2002-2008), and in the second time period average storm intensity ( $\text{mm hr}^{-1}$ ), and max storm intensity ( $\text{mm hr}^{-1}$ ) (2002-2008).

One-way analyses of covariance (ANCOVAs) were used to evaluate differences in baseflow volume, stormflow volume, peak flow, stormflow volume as a percentage of rainfall, time to peak, and curve number between the subcatchments. Precipitation depth and growing season served as covariates. Differences were assessed on an annual basis to limit the temporal variability of subcatchment hydrology due to regrowth and to determine the year when treatment watersheds reached pre-harvest conditions. Additionally, a three-year post-harvest period was assessed to determine differences in treatment effects directly after harvest. Significant pairwise differences ( $p \leq 0.05$ ) between subcatchments were evaluated using Holm-Sidak multiple comparison tests. In one instance, the underlying assumptions of an ANCOVA were not met (e.g. linearity of regression, homogeneity of error variances, independence of error terms, normality of error terms, and homogeneity of regression slopes). For this instance, a one-way analysis of variance (ANOVA) was used and significant pairwise differences ( $p \leq 0.05$ ) were evaluated using Tukey comparison tests.

ANCOVA was preferred to ANOVA because the inclusion of the covariates, precipitation depth and growing season, reduced the amount of unexplained variance in the hydrograph parameters allowing for a more accurate assessment of treatment effects. Hydrograph parameters that were statistically analyzed included baseflow volume ( $\text{m}^3 \text{ ha}^{-1}$ ), stormflow volume ( $\text{m}^3 \text{ ha}^{-1}$ ), peak flow ( $\text{m}^3 \text{ s}^{-1} \text{ ha}^{-1}$ ), stormflow volume as a percentage of rainfall (%), time to peak (hr), and curve number (CN). Baseflow volume, stormflow volume, and peak flow were normalized by drainage area. This was achieved by dividing baseflow volume, stormflow volume, and peak flow by the respective drainage areas in each subcatchment. By design, stormflow volume as a percentage of rainfall is already normalized by drainage area. These measures were taken to reduce the differences in evaluated characteristics due to differences in subcatchment size. Statistical analyses were performed in SigmaPlot 13 (Systat Software, Inc., 2015).

## **2.3 RESULTS AND DISCUSSION**

### **2.3.1 Precipitation Characteristics**

Table 2.3 contains storm event characteristics for the first time period (1982-1993) while Table 2.4 contains storm event characteristics for the second time period (2002-2008). The pre-harvest period had an average of 33 storms per year, comparatively; post-harvest averaged 34 storms per year. Average storm events for the entire analyzation period (1982-1993 and 2002-2008) equaled 34. The yearly cumulative precipitation for the study site from 1982-2008 is shown in Figure 2.9; the average yearly cumulative precipitation for this time period was 1120 mm. The year 2004 had the highest cumulative rainfall at 1490 mm while 2007 had the lowest at 850 mm. The maximum and minimum storm depths in the first study period were 80.0 mm and 11.2 mm, respectively, and in the second time period they were 105.7 mm and 10.4 mm, respectively. Additionally, the maximum and minimum storm depths during pre-harvest were 48.5 mm and 11.2 mm, respectively while post-harvest values were 105.7 mm and 10.4 mm, respectively. The average storm depth in the first time period was



Table 2.3: Storm event characteristics 1982-1993.

| Date                         | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|------------------------------|-----------------------|---|---------------------|
| February 8, 1982             | 49.3                  | 7.4                                     | NG                  |
| February 16, 1982            | 29.5                  | 12.7                                    | NG                  |
| March 15, 1982               | 36.1                  | 22.4                                    | NG                  |
| March 31, 1982               | 23.4                  | 1.0                                     | NG                  |
| May 21, 1982                 | 31.0                  | 11.4                                    | G                   |
| May 28, 1982                 | 16.5                  | 9.1                                     | G                   |
| June 4, 1982                 | 16.3                  | 20.3                                    | G                   |
| July 28, 1982                | 36.8                  | 3.0                                     | G                   |
| August 5, 1982               | 38.1                  | 4.3                                     | G                   |
| September 13, 1982           | 62.2                  | 0.0                                     | G                   |
| September 25, 1982           | 15.7                  | 10.9                                    | G                   |
| November 20, 1982            | 21.3                  | 3.0                                     | NG                  |
| November 30, 1982            | 13.2                  | 24.6                                    | NG                  |
| December 5, 1982             | 15.2                  | 13.2                                    | NG                  |
| December 15, 1982            | 14.5                  | 21.6                                    | NG                  |
| January 21, 1983             | 26.9                  | 1.0                                     | NG                  |
| February 10, 1983            | 22.1                  | 9.1                                     | NG                  |
| April 14, 1983               | 23.6                  | 7.4                                     | NG                  |
| May 3, 1983                  | 34.3                  | 12.2                                    | G                   |
| May 13, 1983                 | 93.5                  | 17.3                                    | G                   |
| May 22, 1983                 | 41.9                  | 19.8                                    | G                   |
| June 3, 1983                 | 27.9                  | 2.3                                     | G                   |
| June 4, 1983                 | 15.0                  | 29.5                                    | G                   |
| July 3, 1983                 | 33.3                  | 0.0                                     | G                   |
| July 5, 1983                 | 13.2                  | 33.3                                    | G                   |
| July 18, 1983                | 12.2                  | 8.9                                     | G                   |
| August 2, 1983               | 25.4                  | 0.0                                     | G                   |
| August 11, 1983              | 32.5                  | 0.5                                     | G                   |
| August 27, 1983 <sup>2</sup> | 14.0                  | 0.0                                     | G                   |
| November 14, 1983            | 30.2                  | 12.7                                    | NG                  |
| December 27, 1983            | 16.3                  | 5.8                                     | NG                  |
| February 27, 1984            | 41.4                  | 5.6                                     | NG                  |
| March 20, 1984               | 27.9                  | 16.0                                    | NG                  |
| March 28, 1984               | 24.1                  | 2.8                                     | NG                  |
| April 4, 1984                | 27.7                  | 1.0                                     | NG                  |
| April 9, 1984                | 41.4                  | 27.7                                    | NG                  |

Table 2.3: Continued

| Date               | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|--------------------|-----------------------|---|---------------------|
| April 21, 1984     | 44.7                  | 18.3                                    | G                   |
| May 4, 1984        | 23.6                  | 23.6                                    | G                   |
| May 6, 1984        | 136.1                 | 42.4                                    | G                   |
| May 23, 1984       | 11.4                  | 0.0                                     | G                   |
| May 28, 1984       | 22.1                  | 21.6                                    | G                   |
| July 26, 1984      | 21.3                  | 17.3                                    | G                   |
| July 27, 1984      | 16.3                  | 33.5                                    | G                   |
| July 30, 1984      | 11.4                  | 49.8                                    | G                   |
| August 22, 1984    | 38.9                  | 7.1                                     | G                   |
| November 4, 1984   | 20.1                  | 9.4                                     | NG                  |
| November 18, 1984  | 64.8                  | 3.3                                     | NG                  |
| November 28, 1984  | 27.4                  | 0.0                                     | NG                  |
| December 20, 1984  | 22.1                  | 12.2                                    | NG                  |
| December 24, 1984  | 21.8                  | 31.0                                    | NG                  |
| January 3, 1985    | 27.4                  | 26.2                                    | NG                  |
| February 11, 1985  | 44.7                  | 0.0                                     | NG                  |
| May 15, 1985       | 18.5                  | 1.0                                     | G                   |
| June 5, 1985       | 18.3                  | 13.0                                    | G                   |
| June 10, 1985      | 22.6                  | 23.1                                    | G                   |
| June 11, 1985      | 28.7                  | 27.4                                    | G                   |
| July 10, 1985      | 27.9                  | 0.0                                     | G                   |
| July 30, 1985      | 27.7                  | 2.5                                     | G                   |
| August 1, 1985     | 38.1                  | 27.7                                    | G                   |
| August 17, 1985    | 14.5                  | 25.4                                    | G                   |
| August 25, 1985    | 13.2                  | 10.9                                    | G                   |
| August 30, 1985    | 39.4                  | 13.2                                    | G                   |
| September 26, 1985 | 14.7                  | 1.8                                     | G                   |
| November 2, 1985   | 62.5                  | 5.8                                     | NG                  |
| December 12, 1985  | 27.9                  | 0.0                                     | NG                  |
| February 2, 1986   | 48.0                  | 2.0                                     | NG                  |
| April 28, 1986     | 13.7                  | 0.0                                     | G                   |
| May 11, 1986       | 26.2                  | 4.1                                     | G                   |
| July 2, 1986       | 19.1                  | 8.9                                     | G                   |
| July 20, 1986      | 16.5                  | 0.0                                     | G                   |
| July 26, 1986      | 18.3                  | 5.1                                     | G                   |
| October 1, 1986    | 16.5                  | 0.8                                     | G                   |
| November 5, 1986   | 30.0                  | 3.3                                     | NG                  |

Table 2.3: Continued

| Date               | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|--------------------|-----------------------|---|---------------------|
| November 8, 1986   | 82.0                  | 40.9                                    | NG                  |
| November 10, 1986  | 23.4                  | 122.9                                   | NG                  |
| December 8, 1986   | 33.8                  | 0.8                                     | NG                  |
| January 18, 1987   | 35.8                  | 5.1                                     | NG                  |
| February 22, 1987  | 23.6                  | 5.6                                     | NG                  |
| February 26, 1987  | 41.1                  | 23.6                                    | NG                  |
| March 18, 1987     | 12.7                  | 13.0                                    | NG                  |
| March 30, 1987     | 72.4                  | 0.0                                     | NG                  |
| May 12, 1987       | 14.0                  | 0.0                                     | G                   |
| May 21, 1987       | 18.5                  | 4.3                                     | G                   |
| May 25, 1987       | 15.2                  | 19.3                                    | G                   |
| June 16, 1987      | 14.0                  | 5.1                                     | G                   |
| June 25, 1987      | 37.1                  | 16.8                                    | G                   |
| July 11, 1987      | 28.7                  | 7.6                                     | G                   |
| August 22, 1987    | 19.1                  | 16.5                                    | G                   |
| September 12, 1987 | 17.0                  | 13.2                                    | G                   |
| November 9, 1987   | 39.9                  | 0.0                                     | NG                  |
| November 16, 1987  | 15.7                  | 0.0                                     | NG                  |
| December 14, 1987  | 13.7                  | 0.0                                     | NG                  |
| December 24, 1987  | 79.0                  | 7.1                                     | NG                  |
| January 18, 1988   | 37.8                  | 6.4                                     | NG                  |
| April 6, 1988      | 36.1                  | 5.8                                     | NG                  |
| May 4, 1988        | 50.3                  | 0.0                                     | G                   |
| June 2, 1988       | 12.2                  | 0.8                                     | G                   |
| June 9, 1988       | 13.2                  | 0.0                                     | G                   |
| August 23, 1988    | 44.7                  | 5.1                                     | G                   |
| September 16, 1988 | 50.8                  | 0.0                                     | G                   |
| November 4, 1988   | 33.0                  | 0.0                                     | NG                  |
| November 27, 1988  | 15.2                  | 2.5                                     | NG                  |
| December 21, 1988  | 30.0                  | 0.0                                     | NG                  |
| December 24, 1988  | 32.0                  | 38.6                                    | NG                  |
| January 11, 1989   | 51.3                  | 20.3                                    | NG                  |
| February 3, 1989   | 31.0                  | 9.1                                     | NG                  |
| February 13, 1989  | 60.7                  | 0.0                                     | NG                  |
| February 20, 1989  | 50.5                  | 30.0                                    | NG                  |
| March 20, 1989     | 43.4                  | 1.3                                     | NG                  |

Table 2.3: Continued

| Date               | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|--------------------|-----------------------|---|---------------------|
| April 3, 1989      | 24.4                  | 18.5                                    | NG                  |
| May 19, 1989       | 14.5                  | 9.4                                     | G                   |
| June 12, 1989      | 104.6                 | 8.9                                     | G                   |
| June 22, 1989      | 14.7                  | 10.7                                    | G                   |
| July 23, 1989      | 25.1                  | 26.9                                    | G                   |
| July 27, 1989      | 20.3                  | 26.9                                    | G                   |
| July 31, 1989      | 26.2                  | 29.0                                    | G                   |
| August 5, 1989     | 31.0                  | 26.2                                    | G                   |
| August 18, 1989    | 37.3                  | 6.1                                     | G                   |
| September 22, 1989 | 56.4                  | 0.0                                     | G                   |
| September 30, 1989 | 36.8                  | 11.4                                    | G                   |
| October 16, 1989   | 102.4                 | 0.0                                     | G                   |
| November 14, 1989  | 35.6                  | 1.8                                     | NG                  |
| January 29, 1990   | 26.2                  | 4.8                                     | NG                  |
| February 3, 1990   | 36.1                  | 26.2                                    | NG                  |
| February 9, 1990   | 35.3                  | 2.5                                     | NG                  |
| February 15, 1990  | 31.0                  | 13.7                                    | NG                  |
| March 16, 1990     | 38.6                  | 0.0                                     | NG                  |
| April 6, 1990      | 25.9                  | 0.0                                     | NG                  |
| May 26, 1990       | 14.5                  | 8.1                                     | G                   |
| May 28, 1990       | 29.2                  | 15.7                                    | G                   |
| June 2, 1990       | 33.8                  | 29.2                                    | G                   |
| June 21, 1990      | 19.8                  | 4.3                                     | G                   |
| July 30, 1990      | 11.4                  | 1.3                                     | G                   |
| August 8, 1990     | 19.1                  | 15.2                                    | G                   |
| August 29, 1990    | 16.8                  | 0.0                                     | G                   |
| September 9, 1990  | 16.5                  | 0.0                                     | G                   |
| September 12, 1990 | 11.4                  | 16.5                                    | G                   |
| October 4, 1990    | 27.9                  | 0.0                                     | G                   |
| December 2, 1990   | 33.8                  | 4.6                                     | NG                  |
| December 20, 1990  | 51.1                  | 61.7                                    | NG                  |
| December 27, 1990  | 38.1                  | 38.4                                    | NG                  |
| December 30, 1990  | 31.0                  | 38.1                                    | NG                  |
| January 6, 1991    | 32.3                  | 0.0                                     | NG                  |
| February 6, 1991   | 23.6                  | 1.3                                     | NG                  |
| February 13, 1991  | 29.7                  | 0.0                                     | NG                  |

Table 2.3: Continued

| Date               | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|--------------------|-----------------------|---|---------------------|
| February 17, 1991  | 62.2                  | 31.0                                    | NG                  |
| March 22, 1991     | 34.0                  | 12.7                                    | NG                  |
| March 29, 1991     | 34.3                  | 9.4                                     | NG                  |
| April 15, 1991     | 26.7                  | 11.4                                    | NG                  |
| April 19, 1991     | 29.2                  | 26.7                                    | NG                  |
| May 9, 1991        | 17.8                  | 6.9                                     | G                   |
| May 18, 1991       | 18.5                  | 0.0                                     | G                   |
| May 27, 1991       | 16.5                  | 0.0                                     | G                   |
| May 29, 1991       | 29.2                  | 16.5                                    | G                   |
| June 22, 1991      | 30.5                  | 32.3                                    | G                   |
| June 25, 1991      | 35.6                  | 39.9                                    | G                   |
| July 10, 1991      | 17.3                  | 7.6                                     | G                   |
| July 12, 1991      | 30.5                  | 23.6                                    | G                   |
| August 7, 1991     | 31.0                  | 0.0                                     | G                   |
| October 5, 1991    | 38.1                  | 0.0                                     | G                   |
| October 15, 1991   | 21.6                  | 0.0                                     | G                   |
| November 21, 1991  | 59.7                  | 2.0                                     | NG                  |
| December 13, 1991  | 24.1                  | 20.8                                    | NG                  |
| December 28, 1991  | 40.9                  | 10.2                                    | NG                  |
| February 24, 1992  | 40.1                  | 7.4                                     | NG                  |
| March 30, 1992     | 34.8                  | 5.6                                     | NG                  |
| May 28, 1992       | 38.9                  | 0.8                                     | G                   |
| June 17, 1992      | 15.2                  | 7.4                                     | G                   |
| June 24, 1992      | 28.4                  | 0.8                                     | G                   |
| July 1, 1992       | 80.8                  | 14.5                                    | G                   |
| July 14, 1992      | 24.6                  | 0.3                                     | G                   |
| July 17, 1992      | 11.2                  | 24.6                                    | G                   |
| July 22, 1992      | 17.5                  | 12.4                                    | G                   |
| July 24, 1992      | 57.9                  | 18.8                                    | G                   |
| August 8, 1992     | 22.9                  | 6.4                                     | G                   |
| August 27, 1992    | 49.8                  | 5.3                                     | G                   |
| September 4, 1992  | 11.7                  | 0.0                                     | G                   |
| September 18, 1992 | 25.1                  | 0.0                                     | G                   |
| December 20, 1992  | 29.2                  | 22.1                                    | NG                  |
| March 29, 1993     | 40.6                  | 15.0                                    | NG                  |

Table 2.3: Continued

| Date               | Precipitation<br>(mm) | Total 5-day antecedent<br>rainfall (mm) | Season <sup>1</sup> |
|--------------------|-----------------------|---|---------------------|
| April 25, 1993     | 21.6                  | 17.3                                    | G                   |
| May 9, 1993        | 23.9                  | 0.0                                     | G                   |
| June 9, 1993       | 21.6                  | 11.4                                    | G                   |
| June 21, 1993      | 21.6                  | 0.0                                     | G                   |
| July 13, 1993      | 58.4                  | 25.4                                    | G                   |
| July 15, 1993      | 17.0                  | 75.4                                    | G                   |
| July 26, 1993      | 36.8                  | 0.0                                     | G                   |
| September 2, 1993  | 30.7                  | 4.1                                     | G                   |
| September 15, 1993 | 57.9                  | 0.0                                     | G                   |
| October 18, 1993   | 36.6                  | 9.1                                     | G                   |

Hydrograph was plotted for storm event.

<sup>1</sup>G-Growing season runs from April 20 - October 26.

NG-Non-Growing season runs from October 27 - April 19.

<sup>2</sup>Final storm before harvest.

Table 2.4: Storm Event Characteristics 2002-2008.

| Start Date     | End Date       | Duration<br>(hr) | Depth<br>(mm) | 5 Day<br>Prior (mm) | Avg. Int.<br>(mm/hr) | Max Int.<br>(mm/hr) | Season <sup>1</sup> |
|----------------|----------------|------------------|---------------|---------------------|----------------------|---------------------|---------------------|
| 1/9/02 13:00   | 1/9/02 17:45   | 4.7              | 11.2          | 3.3                 | 2.6                  | 6.1                 | NG                  |
| 1/22/02 22:15  | 1/23/02 8:15   | 10.0             | 14.0          | 21.8                | 2.2                  | 6.1                 | NG                  |
| 1/24/02 5:45   | 1/24/02 19:15  | 13.5             | 22.9          | 31.0                | 2.4                  | 6.1                 | NG                  |
| 2/7/02 11:30   | 2/7/02 16:00   | 4.5              | 11.9          | 0.3                 | 2.6                  | 4.1                 | NG                  |
| 3/16/02 3:45   | 3/16/02 8:15   | 4.5              | 16.8          | 2.5                 | 5.6                  | 22.4                | NG                  |
| 3/17/02 19:15  | 3/18/02 6:15   | 11.0             | 23.4          | 27.2                | 2.2                  | 6.1                 | NG                  |
| 3/19/02 6:15   | 3/19/02 14:00  | 7.7              | 13.2          | 50.0                | 1.9                  | 5.1                 | NG                  |
| 3/20/02 7:15   | 3/20/02 12:15  | 5.0              | 19.1          | 64.8                | 3.6                  | 7.1                 | NG                  |
| 3/26/02 3:45   | 3/26/02 13:30  | 9.7              | 16.3          | 0.0                 | 3.0                  | 15.2                | NG                  |
| 3/29/02 19:15  | 3/30/02 3:00   | 7.7              | 16.5          | 16.5                | 3.9                  | 16.3                | NG                  |
| 3/31/02 1:45   | 3/31/02 15:45  | 14.0             | 23.1          | 36.8                | 2.3                  | 8.1                 | NG                  |
| 4/14/02 14:45  | 4/14/02 18:30  | 3.7              | 14.7          | 7.9                 | 8.4                  | 30.5                | NG                  |
| 4/21/02 23:00  | 4/25/02 1:15   | 74.2             | 50.8          | 5.3                 | 7.5                  | 25.4                | G                   |
| 4/28/02 6:30   | 4/28/02 9:45   | 3.2              | 15.5          | 31.2                | 5.2                  | 16.3                | G                   |
| 5/2/02 12:00   | 5/2/02 22:00   | 10.0             | 59.4          | 27.7                | 9.5                  | 41.7                | G                   |
| 5/8/02 11:00   | 5/9/02 13:15   | 26.2             | 11.7          | 18.0                | 5.2                  | 18.3                | G                   |
| 5/13/02 3:30   | 5/13/02 10:45  | 7.2              | 42.2          | 12.7                | 6.0                  | 30.5                | G                   |
| 5/17/02 19:30  | 5/18/02 6:30   | 11.0             | 29.5          | 43.4                | 4.7                  | 23.4                | G                   |
| 6/6/02 2:30    | 6/6/02 14:30   | 12.0             | 27.7          | 12.4                | 4.8                  | 19.3                | G                   |
| 6/12/02 19:45  | 6/12/02 23:15  | 3.5              | 13.7          | 3.8                 | 4.6                  | 18.3                | G                   |
| 7/12/02 18:30  | 7/13/02 17:30  | 23.0             | 61.2          | 0.3                 | 4.5                  | 24.4                | G                   |
| 7/19/02 14:00  | 7/20/02 5:00   | 15.0             | 18.3          | 13.0                | 8.1                  | 32.5                | G                   |
| 8/15/02 16:30  | 8/15/02 17:00  | 0.5              | 18.5          | 1.8                 | 24.7                 | 43.7                | G                   |
| 9/21/02 1:00   | 9/21/02 12:45  | 11.7             | 17.5          | 0.5                 | 3.0                  | 12.2                | G                   |
| 9/22/02 12:45  | 9/22/02 15:15  | 2.5              | 15.2          | 29.0                | 6.1                  | 34.5                | G                   |
| 9/25/02 18:15  | 9/26/02 23:45  | 29.5             | 37.3          | 43.7                | 1.8                  | 5.1                 | G                   |
| 10/10/02 1:15  | 10/11/02 5:45  | 28.5             | 29.7          | 4.6                 | 1.9                  | 6.1                 | G                   |
| 10/15/02 17:15 | 10/16/02 10:45 | 17.5             | 15.7          | 15.2                | 1.4                  | 3.0                 | G                   |
| 10/28/02 6:15  | 10/28/02 11:45 | 5.5              | 22.1          | 6.6                 | 3.8                  | 10.2                | NG                  |
| 10/29/02 4:30  | 10/29/02 9:30  | 5.0              | 14.2          | 28.7                | 3.8                  | 14.2                | NG                  |
| 11/5/02 9:00   | 11/6/02 2:00   | 17.0             | 17.8          | 4.8                 | 1.7                  | 5.1                 | NG                  |
| 11/10/02 23:30 | 11/11/02 4:15  | 4.7              | 16.0          | 2.8                 | 3.8                  | 15.2                | NG                  |
| 11/15/02 13:30 | 11/16/02 0:30  | 11.0             | 11.7          | 16.0                | 1.4                  | 2.0                 | NG                  |
| 12/5/02 1:15   | 12/5/02 6:45   | 5.5              | 11.2          | 0.8                 | 2.2                  | 12.2                | NG                  |
| 12/10/02 20:15 | 12/11/02 8:15  | 12.0             | 15.7          | 12.7                | 1.7                  | 4.1                 | NG                  |
| 12/13/02 8:30  | 12/14/02 9:00  | 24.5             | 25.9          | 27.9                | 2.3                  | 10.2                | NG                  |
| 12/19/02 19:45 | 12/20/02 4:00  | 8.2              | 14.7          | 0.5                 | 2.8                  | 8.1                 | NG                  |
| 1/1/03 6:30    | 1/1/03 9:00    | 2.5              | 11.2          | 3.3                 | 4.5                  | 9.1                 | NG                  |
| 1/29/03 0:45   | 1/29/03 16:15  | 15.5             | 15.2          | 3.3                 | 1.5                  | 4.1                 | NG                  |

Table 2.4: Continued

| Start Date     | End Date       | Duration<br>(hr) | Depth<br>(mm) | 5 Day<br>Prior (mm) | Avg. Int.<br>(mm/hr) | Max Int.<br>(mm/hr) | Season <sup>1</sup> |
|----------------|----------------|------------------|---------------|---------------------|----------------------|---------------------|---------------------|
| 2/3/03 21:15   | 2/4/03 4:00    | 6.7              | 14.0          | 2.3                 | 2.8                  | 10.2                | NG                  |
| 2/15/03 10:30  | 2/18/03 5:30   | 67.0             | 105.7         | 4.8                 | 2.2                  | 10.2                | NG                  |
| 2/23/03 1:45   | 2/23/03 8:30   | 6.7              | 25.7          | 7.4                 | 4.3                  | 10.2                | NG                  |
| 2/23/03 16:45  | 2/23/03 19:30  | 2.7              | 16.3          | 31.2                | 8.1                  | 20.3                | NG                  |
| 3/30/03 6:15   | 3/30/03 10:30  | 4.2              | 13.0          | 5.6                 | 3.0                  | 8.1                 | NG                  |
| 4/7/03 16:00   | 4/8/03 6:15    | 14.2             | 27.9          | 11.9                | 2.1                  | 5.1                 | NG                  |
| 4/9/03 11:30   | 4/9/03 20:45   | 9.2              | 19.8          | 40.4                | 2.9                  | 9.1                 | NG                  |
| 4/10/03 0:45   | 4/10/03 22:00  | 21.2             | 17.0          | 60.2                | 1.8                  | 8.1                 | NG                  |
| 4/17/03 13:30  | 4/18/03 2:00   | 12.5             | 25.7          | 0.0                 | 2.6                  | 8.1                 | NG                  |
| 5/8/03 23:15   | 5/9/03 2:15    | 3.0              | 20.1          | 14.5                | 10.0                 | 51.8                | G                   |
| 5/15/03 16:15  | 5/15/03 18:00  | 1.7              | 13.0          | 16.8                | 8.6                  | 42.7                | G                   |
| 5/29/03 8:30   | 5/29/03 16:15  | 7.7              | 11.4          | 7.1                 | 2.5                  | 9.1                 | G                   |
| 6/6/03 20:30   | 6/7/03 9:30    | 13.0             | 30.2          | 14.2                | 4.2                  | 17.3                | G                   |
| 6/11/03 13:45  | 6/11/03 18:30  | 4.7              | 12.2          | 33.8                | 5.4                  | 33.5                | G                   |
| 6/16/03 15:00  | 6/16/03 17:30  | 2.5              | 13.2          | 38.6                | 4.8                  | 15.2                | G                   |
| 7/10/03 17:15  | 7/10/03 21:15  | 4.0              | 22.6          | 15.7                | 6.5                  | 25.4                | G                   |
| 7/28/03 9:45   | 7/28/03 15:45  | 6.0              | 25.1          | 3.0                 | 9.1                  | 36.6                | G                   |
| 8/3/03 21:45   | 8/3/03 22:30   | 0.7              | 16.0          | 21.3                | 16.0                 | 26.4                | G                   |
| 8/4/03 6:15    | 8/4/03 8:45    | 2.5              | 12.7          | 37.3                | 4.6                  | 18.3                | G                   |
| 8/17/03 15:30  | 8/17/03 17:30  | 2.0              | 16.5          | 0.0                 | 13.2                 | 30.5                | G                   |
| 9/3/03 21:30   | 9/4/03 8:00    | 10.5             | 63.8          | 36.8                | 10.5                 | 34.5                | G                   |
| 9/22/03 3:15   | 9/22/03 18:00  | 14.7             | 31.0          | 0.0                 | 3.2                  | 13.2                | G                   |
| 9/27/03 9:00   | 9/27/03 15:15  | 6.2              | 11.4          | 26.4                | 4.2                  | 10.2                | G                   |
| 10/14/03 12:00 | 10/14/03 16:00 | 4.0              | 23.6          | 0.3                 | 5.6                  | 17.3                | G                   |
| 10/26/03 11:45 | 10/27/03 11:00 | 23.2             | 21.1          | 0.0                 | 2.1                  | 7.1                 | G                   |
| 11/5/03 11:15  | 11/5/03 17:30  | 6.2              | 11.4          | 10.4                | 3.0                  | 13.2                | NG                  |
| 11/12/03 17:30 | 11/12/03 19:00 | 1.5              | 15.2          | 2.3                 | 8.7                  | 45.7                | NG                  |
| 11/18/03 18:45 | 11/19/03 13:00 | 18.2             | 46.2          | 14.2                | 3.3                  | 13.2                | NG                  |
| 11/28/03 5:45  | 11/28/03 15:45 | 10.0             | 15.5          | 12.7                | 2.1                  | 16.3                | NG                  |
| 12/10/03 12:00 | 12/10/03 21:45 | 9.7              | 14.0          | 6.9                 | 3.1                  | 17.3                | NG                  |
| 12/14/03 3:45  | 12/14/03 18:30 | 14.7             | 13.0          | 16.3                | 1.3                  | 3.0                 | NG                  |
| 12/16/03 20:00 | 12/17/03 4:15  | 8.2              | 11.2          | 13.2                | 1.9                  | 3.0                 | NG                  |
| 12/29/03 21:00 | 12/30/03 2:45  | 5.7              | 12.2          | 0.0                 | 2.6                  | 15.2                | NG                  |
| 1/1/04 23:45   | 1/2/04 16:15   | 16.5             | 45.5          | 12.4                | 3.4                  | 23.4                | NG                  |
| 1/5/04 1:00    | 1/5/04 8:30    | 7.5              | 20.1          | 46.7                | 3.0                  | 13.2                | NG                  |
| 1/17/04 17:45  | 1/18/04 13:30  | 19.7             | 27.2          | 2.5                 | 1.9                  | 5.1                 | NG                  |
| 2/2/04 18:00   | 2/3/04 10:15   | 16.2             | 15.5          | 6.9                 | 2.1                  | 6.1                 | NG                  |
| 2/5/04 11:45   | 2/5/04 17:45   | 6.0              | 34.0          | 22.4                | 5.4                  | 10.2                | NG                  |
| 2/5/04 20:30   | 2/6/04 7:30    | 11.0             | 31.2          | 55.1                | 2.8                  | 5.1                 | NG                  |



Table 2.4: Continued

| Start Date     | End Date       | Duration<br>(hr) | Depth<br>(mm) | 5 Day<br>Prior (mm) | Avg. Int.<br>(mm/hr) | Max Int.<br>(mm/hr) | Season <sup>1</sup> |
|----------------|----------------|------------------|---------------|---------------------|----------------------|---------------------|---------------------|
| 3/5/04 19:45   | 3/6/04 3:30    | 7.8              | 45.0          | 6.1                 | 5.6                  | 18.3                | NG                  |
| 3/16/04 1:30   | 3/16/04 9:15   | 7.7              | 16.0          | 0.5                 | 2.6                  | 7.1                 | NG                  |
| 3/30/04 1:45   | 3/30/04 9:00   | 7.3              | 16.0          | 0.5                 | 2.6                  | 9.1                 | NG                  |
| 4/12/04 7:15   | 4/12/04 19:30  | 12.3             | 27.2          | 2.5                 | 2.5                  | 9.1                 | NG                  |
| 4/13/04 1:45   | 4/13/04 10:15  | 8.5              | 28.4          | 29.7                | 4.1                  | 11.2                | NG                  |
| 5/2/04 5:30    | 5/2/04 13:15   | 7.7              | 14.7          | 0.5                 | 2.5                  | 9.1                 | G                   |
| 5/16/04 5:15   | 5/16/04 10:30  | 5.3              | 14.0          | 5.1                 | 3.5                  | 8.1                 | G                   |
| 5/24/04 21:15  | 5/24/04 23:15  | 2.0              | 24.9          | 0.0                 | 11.1                 | 26.4                | G                   |
| 5/26/04 23:00  | 5/27/04 6:00   | 7.0              | 38.6          | 40.4                | 7.0                  | 40.6                | G                   |
| 5/27/04 23:45  | 5/28/04 1:15   | 1.5              | 19.3          | 79.0                | 12.9                 | 39.6                | G                   |
| 5/28/04 2:45   | 5/28/04 6:30   | 3.8              | 11.9          | 98.3                | 4.3                  | 23.4                | G                   |
| 5/30/04 11:00  | 5/30/04 15:15  | 4.2              | 49.5          | 90.2                | 19.8                 | 64.0                | G                   |
| 5/31/04 1:30   | 5/31/04 5:00   | 3.5              | 16.5          | 140.7               | 5.5                  | 44.7                | G                   |
| 6/4/04 12:45   | 6/4/04 16:30   | 3.8              | 14.0          | 37.6                | 4.0                  | 10.2                | G                   |
| 6/11/04 17:45  | 6/11/04 18:15  | 0.5              | 11.4          | 2.8                 | 15.2                 | 43.7                | G                   |
| 6/15/04 20:15  | 6/15/04 23:15  | 3.0              | 37.3          | 26.4                | 12.4                 | 25.4                | G                   |
| 6/22/04 18:45  | 6/22/04 20:00  | 1.3              | 10.7          | 3.0                 | 7.1                  | 17.3                | G                   |
| 6/25/04 7:30   | 6/25/04 18:30  | 11.0             | 23.9          | 17.3                | 2.7                  | 10.2                | G                   |
| 7/6/04 21:15   | 7/6/04 22:45   | 1.5              | 13.7          | 18.3                | 7.8                  | 26.4                | G                   |
| 7/22/04 7:45   | 7/22/04 10:30  | 2.8              | 13.0          | 4.3                 | 5.2                  | 18.3                | G                   |
| 7/26/04 15:45  | 7/26/04 22:15  | 6.5              | 25.7          | 16.8                | 6.0                  | 28.4                | G                   |
| 7/27/04 2:00   | 7/27/04 5:00   | 3.0              | 13.0          | 42.4                | 4.3                  | 16.3                | G                   |
| 7/31/04 8:30   | 7/31/04 14:15  | 5.7              | 13.0          | 39.9                | 3.2                  | 22.4                | G                   |
| 8/5/04 4:30    | 8/5/04 8:30    | 4.0              | 22.4          | 19.6                | 6.0                  | 29.5                | G                   |
| 8/12/04 7:15   | 8/12/04 13:15  | 6.0              | 20.3          | 1.8                 | 3.4                  | 11.2                | G                   |
| 9/7/04 9:45    | 9/9/04 1:15    | 39.5             | 92.7          | 0.8                 | 3.1                  | 10.2                | G                   |
| 9/16/04 18:15  | 9/17/04 17:15  | 23.0             | 78.7          | 3.0                 | 4.3                  | 17.3                | G                   |
| 10/2/04 7:15   | 10/2/04 12:00  | 4.8              | 10.4          | 0.0                 | 2.3                  | 7.1                 | G                   |
| 10/13/04 9:15  | 10/13/04 10:30 | 1.3              | 14.2          | 7.9                 | 9.5                  | 38.6                | G                   |
| 10/18/04 22:15 | 10/19/04 4:00  | 5.8              | 34.0          | 12.4                | 6.8                  | 19.3                | G                   |
| 10/27/04 1:30  | 10/27/04 7:30  | 6.0              | 30.7          | 8.6                 | 5.9                  | 15.2                | NG                  |
| 11/4/04 1:30   | 11/4/04 10:00  | 8.5              | 38.4          | 0.5                 | 6.7                  | 32.5                | NG                  |
| 11/11/04 23:00 | 11/12/04 7:15  | 8.2              | 15.7          | 0.0                 | 2.4                  | 6.1                 | NG                  |
| 11/30/04 6:30  | 12/1/04 3:15   | 20.7             | 42.4          | 5.3                 | 2.5                  | 11.2                | NG                  |
| 12/6/04 0:30   | 12/6/04 4:45   | 4.2              | 10.7          | 7.1                 | 3.6                  | 13.2                | NG                  |
| 12/6/04 21:00  | 12/7/04 9:00   | 12.0             | 13.0          | 11.2                | 1.6                  | 4.1                 | NG                  |
| 12/9/04 5:30   | 12/9/04 16:45  | 11.2             | 32.8          | 24.1                | 3.5                  | 8.1                 | NG                  |
| 12/23/04 3:30  | 12/23/04 8:15  | 4.7              | 14.0          | 1.3                 | 3.1                  | 7.1                 | NG                  |
| 1/4/05 13:45   | 1/4/05 21:45   | 8.0              | 12.2          | 10.2                | 2.0                  | 4.1                 | NG                  |

Table 2.4: Continued

| Start Date     | End Date       | Duration<br>(hr) | Depth<br>(mm) | 5 Day<br>Prior (mm) | Avg. Int.<br>(mm/hr) | Max Int.<br>(mm/hr) | Season <sup>1</sup> |
|----------------|----------------|------------------|---------------|---------------------|----------------------|---------------------|---------------------|
| 1/7/05 14:30   | 1/8/05 2:15    | 11.7             | 25.9          | 26.4                | 3.0                  | 10.2                | NG                  |
| 1/13/05 17:30  | 1/14/05 0:30   | 7.0              | 13.2          | 5.1                 | 1.9                  | 4.1                 | NG                  |
| 1/29/05 6:15   | 1/30/05 0:30   | 18.3             | 22.9          | 3.3                 | 1.8                  | 7.1                 | NG                  |
| 2/20/05 11:15  | 2/20/05 23:00  | 11.8             | 17.5          | 4.3                 | 2.3                  | 14.2                | NG                  |
| 2/28/05 0:30   | 2/28/05 8:00   | 7.5              | 14.2          | 0.8                 | 2.0                  | 4.1                 | NG                  |
| 3/7/05 21:15   | 3/8/05 4:15    | 7.0              | 13.5          | 6.6                 | 1.9                  | 6.1                 | NG                  |
| 3/28/05 2:30   | 3/28/05 6:15   | 3.7              | 18.5          | 10.7                | 4.9                  | 12.2                | NG                  |
| 4/1/05 19:15   | 4/2/05 3:15    | 8.0              | 18.5          | 36.1                | 3.7                  | 15.2                | NG                  |
| 4/2/05 13:00   | 4/2/05 19:45   | 6.8              | 12.2          | 43.4                | 2.1                  | 7.1                 | NG                  |
| 4/29/05 16:15  | 4/30/05 5:30   | 13.3             | 49.8          | 19.3                | 5.4                  | 14.2                | G                   |
| 5/19/05 18:30  | 5/20/05 1:15   | 6.7              | 30.5          | 3.0                 | 5.8                  | 28.4                | G                   |
| 5/22/05 23:30  | 5/23/05 1:00   | 1.5              | 13.5          | 31.8                | 7.7                  | 17.3                | G                   |
| 6/3/05 13:45   | 6/3/05 16:00   | 2.3              | 13.5          | 7.6                 | 6.7                  | 29.5                | G                   |
| 6/20/05 18:15  | 6/20/05 20:45  | 2.5              | 30.2          | 0.0                 | 11.0                 | 37.6                | G                   |
| 7/1/05 14:15   | 7/1/05 18:00   | 3.8              | 23.9          | 12.4                | 9.6                  | 19.3                | G                   |
| 7/7/05 7:15    | 7/7/05 16:00   | 8.8              | 14.5          | 0.8                 | 2.1                  | 5.1                 | G                   |
| 7/13/05 5:15   | 7/13/05 13:30  | 8.3              | 14.5          | 2.8                 | 2.1                  | 6.1                 | G                   |
| 7/27/05 14:30  | 7/27/05 17:30  | 3.0              | 13.7          | 0.5                 | 7.8                  | 42.7                | G                   |
| 8/5/05 18:30   | 8/5/05 19:45   | 1.2              | 18.5          | 0.0                 | 12.4                 | 47.8                | G                   |
| 8/16/05 13:30  | 8/16/05 14:30  | 1.0              | 14.7          | 0.0                 | 11.8                 | 26.4                | G                   |
| 8/19/05 19:45  | 8/19/05 20:30  | 0.8              | 16.5          | 22.4                | 16.5                 | 52.8                | G                   |
| 9/26/05 3:30   | 9/26/05 10:30  | 7.0              | 11.9          | 0.0                 | 1.8                  | 4.1                 | G                   |
| 11/16/05 0:00  | 11/16/05 5:45  | 5.8              | 27.9          | 14.5                | 5.1                  | 25.4                | NG                  |
| 11/28/05 23:30 | 11/29/05 7:30  | 8.0              | 21.6          | 0.0                 | 3.3                  | 12.2                | NG                  |
| 12/3/05 21:45  | 12/4/05 2:15   | 4.5              | 11.9          | 24.4                | 3.4                  | 7.1                 | NG                  |
| 12/8/05 14:45  | 12/9/05 0:00   | 9.2              | 12.7          | 14.5                | 1.7                  | 4.1                 | NG                  |
| 12/15/05 1:15  | 12/15/05 15:15 | 14.0             | 20.3          | 3.3                 | 1.8                  | 4.1                 | NG                  |
| 1/17/06 14:15  | 1/18/06 1:30   | 11.3             | 25.1          | 18.3                | 2.6                  | 7.1                 | NG                  |
| 1/23/06 0:15   | 1/23/06 11:00  | 10.8             | 31.5          | 17.3                | 3.8                  | 12.2                | NG                  |
| 3/13/06 16:45  | 3/13/06 21:45  | 5.0              | 17.5          | 15.5                | 5.4                  | 23.4                | NG                  |
| 4/2/06 18:45   | 4/3/06 5:30    | 10.8             | 17.3          | 7.9                 | 3.8                  | 15.2                | NG                  |
| 4/7/06 16:15   | 4/7/06 17:45   | 1.5              | 18.5          | 22.1                | 10.6                 | 43.7                | NG                  |
| 4/19/06 6:00   | 4/19/06 9:00   | 3.0              | 13.0          | 13.2                | 4.0                  | 20.3                | NG                  |
| 4/21/06 20:00  | 4/22/06 5:15   | 9.2              | 17.3          | 27.9                | 2.6                  | 9.1                 | G                   |
| 5/2/06 17:30   | 5/2/06 18:15   | 0.7              | 12.7          | 5.1                 | 12.7                 | 25.4                | G                   |
| 5/25/06 22:45  | 5/26/06 3:00   | 4.3              | 22.9          | 2.5                 | 5.7                  | 26.4                | G                   |
| 5/31/06 15:45  | 5/31/06 19:30  | 3.8              | 17.8          | 0.0                 | 7.9                  | 43.7                | G                   |
| 6/11/06 0:15   | 6/11/06 3:15   | 3.0              | 12.7          | 0.0                 | 5.6                  | 23.4                | G                   |
| 6/23/06 1:45   | 6/23/06 6:30   | 4.8              | 25.1          | 5.6                 | 5.3                  | 35.6                | G                   |

Table 2.4: Continued

| Start Date     | End Date       | Duration<br>(hr) | Depth<br>(mm) | 5 Day<br>Prior (mm) | Avg. Int.<br>(mm/hr) | Max Int.<br>(mm/hr) | Season <sup>1</sup> |
|----------------|----------------|------------------|---------------|---------------------|----------------------|---------------------|---------------------|
| 6/25/06 12:15  | 6/25/06 19:00  | 6.8              | 14.0          | 37.3                | 3.7                  | 15.2                | G                   |
| 7/4/06 18:15   | 7/4/06 19:30   | 1.3              | 13.2          | 1.5                 | 10.6                 | 25.4                | G                   |
| 7/5/06 10:45   | 7/5/06 17:15   | 6.5              | 30.5          | 23.4                | 7.6                  | 28.4                | G                   |
| 7/20/06 16:00  | 7/20/06 16:15  | 0.2              | 11.4          | 0.0                 | 22.9                 | 44.7                | G                   |
| 8/11/06 17:45  | 8/11/06 20:00  | 2.3              | 13.2          | 8.6                 | 7.5                  | 29.5                | G                   |
| 8/28/06 13:45  | 8/28/06 15:30  | 1.8              | 14.5          | 0.3                 | 8.3                  | 17.3                | G                   |
| 9/18/06 18:00  | 9/18/06 19:30  | 1.5              | 12.4          | 0.0                 | 7.1                  | 16.3                | G                   |
| 9/23/06 5:30   | 9/23/06 10:15  | 4.7              | 24.1          | 19.6                | 4.8                  | 15.2                | G                   |
| 9/23/06 19:00  | 9/23/06 22:30  | 3.5              | 13.0          | 34.5                | 4.0                  | 21.3                | G                   |
| 9/30/06 16:45  | 9/30/06 23:00  | 6.3              | 12.4          | 14.2                | 2.8                  | 14.2                | G                   |
| 10/16/06 17:30 | 10/17/06 5:00  | 11.5             | 21.3          | 4.3                 | 2.1                  | 6.1                 | G                   |
| 10/26/06 21:30 | 10/27/06 23:00 | 25.5             | 61.7          | 0.3                 | 3.1                  | 15.2                | G                   |
| 11/1/06 18:00  | 11/1/06 21:30  | 3.5              | 14.7          | 16.5                | 5.4                  | 19.3                | NG                  |
| 11/7/06 17:45  | 11/8/06 12:45  | 19.0             | 17.5          | 2.8                 | 1.5                  | 3.0                 | NG                  |
| 11/16/06 3:30  | 11/16/06 6:15  | 2.7              | 12.7          | 16.3                | 4.2                  | 12.2                | NG                  |
| 12/22/06 6:45  | 12/22/06 12:30 | 5.8              | 18.8          | 9.1                 | 3.3                  | 11.2                | NG                  |
| 1/4/07 22:30   | 1/5/07 6:00    | 7.5              | 16.0          | 12.2                | 2.2                  | 10.2                | NG                  |
| 1/7/07 11:45   | 1/7/07 19:45   | 8.0              | 16.8          | 16.5                | 2.5                  | 9.1                 | NG                  |
| 1/21/07 7:45   | 1/21/07 14:45  | 7.0              | 17.8          | 0.0                 | 2.5                  | 6.1                 | NG                  |
| 3/1/07 9:45    | 3/1/07 23:15   | 13.5             | 35.6          | 7.4                 | 3.2                  | 7.1                 | NG                  |
| 3/28/07 10:30  | 3/28/07 13:00  | 2.5              | 12.7          | 0.0                 | 5.1                  | 11.2                | NG                  |
| 4/3/07 20:30   | 4/4/07 0:45    | 4.2              | 15.5          | 1.8                 | 4.8                  | 14.2                | NG                  |
| 4/11/07 12:15  | 4/11/07 23:00  | 10.7             | 20.8          | 1.0                 | 3.3                  | 14.2                | NG                  |
| 4/14/07 0:30   | 4/15/07 4:00   | 27.5             | 39.9          | 20.8                | 2.2                  | 15.2                | NG                  |
| 5/16/07 11:15  | 5/16/07 14:00  | 2.7              | 19.1          | 0.0                 | 6.9                  | 30.5                | G                   |
| 6/1/07 18:15   | 6/1/07 19:15   | 1.0              | 33.3          | 0.0                 | 26.6                 | 99.6                | G                   |
| 6/5/07 13:15   | 6/5/07 21:00   | 7.7              | 20.1          | 40.9                | 5.7                  | 30.5                | G                   |
| 6/18/07 18:00  | 6/18/07 18:30  | 0.5              | 17.8          | 0.0                 | 23.7                 | 36.6                | G                   |
| 6/25/07 17:45  | 6/25/07 19:00  | 1.2              | 11.9          | 6.1                 | 15.9                 | 39.6                | G                   |
| 6/29/07 16:15  | 6/29/07 20:00  | 3.7              | 13.5          | 24.1                | 5.4                  | 21.3                | G                   |
| 7/19/07 21:00  | 7/19/07 23:45  | 2.7              | 15.7          | 7.4                 | 5.2                  | 18.3                | G                   |
| 7/27/07 11:00  | 7/27/07 12:30  | 1.5              | 13.5          | 3.8                 | 9.0                  | 24.4                | G                   |
| 7/27/07 20:45  | 7/28/07 15:15  | 18.5             | 20.1          | 17.3                | 2.1                  | 11.2                | G                   |
| 8/3/07 15:00   | 8/3/07 15:30   | 0.5              | 15.0          | 0.0                 | 20.0                 | 53.8                | G                   |
| 9/11/07 4:00   | 9/11/07 10:15  | 6.2              | 18.3          | 2.5                 | 3.9                  | 11.2                | G                   |
| 10/19/07 4:00  | 10/19/07 5:15  | 1.2              | 13.7          | 3.3                 | 9.1                  | 21.3                | G                   |
| 10/23/07 11:15 | 10/24/07 20:00 | 32.7             | 47.8          | 15.2                | 2.4                  | 14.2                | G                   |
| 11/5/07 20:00  | 11/5/07 23:00  | 3.0              | 14.7          | 0.0                 | 5.4                  | 15.2                | NG                  |
| 11/14/07 17:30 | 11/15/07 2:45  | 9.2              | 33.3          | 5.8                 | 4.8                  | 15.2                | NG                  |

Table 2.4: Continued

| Start Date     | End Date   | Duration (hr) | Depth (mm) | 5 Day Prior (mm) | Avg. Int. (mm/hr) | Max Int. (mm/hr) | Season <sup>1</sup> |
|----------------|--|---------------|------------|------------------|-------------------|------------------|---------------------|
| 11/25/07 23:00 | 11/26/07 10:15   | 11.2          | 14.0       | 5.8              | 1.9               | 7.1              | NG                  |
| 11/26/07 17:45 | 11/26/07 19:30   | 1.7           | 12.2       | 19.8             | 7.0               | 19.3             | NG                  |
| 12/10/07 2:00  | 12/10/07 18:30   | 16.5          | 27.2       | 21.3             | 2.6               | 10.2             | NG                  |
| 12/13/07 8:30  | 12/13/07 14:15   | 5.7           | 14.7       | 42.9             | 2.9               | 16.3             | NG                  |
| 12/20/07 22:00 | 12/21/07 5:45  | 7.7           | 15.0       | 4.1              | 2.3               | 4.1              | NG                  |
| 12/28/07 11:15 | 12/28/07 20:45   | 9.5           | 15.0       | 1.0              | 2.5               | 7.1              | NG                  |
| 1/29/08 14:00  | 1/29/08 23:30  | 9.5           | 26.4       | 0.5              | 3.8               | 8.1              | NG                  |
| 2/6/08 4:00    | 2/6/08 15:15   | 11.3          | 14.7       | 11.7             | 2.8               | 10.2             | NG                  |
| 3/4/08 4:30    | 3/4/08 9:00  | 4.5           | 11.2       | 5.8              | 2.5               | 4.1              | NG                  |
| 3/7/08 7:45    | 3/7/08 17:15   | 9.5           | 23.1       | 22.1             | 2.6               | 5.1              | NG                  |
| 3/9/08 10:00   | 3/9/08 11:45   | 1.8           | 11.7       | 34.5             | 5.8               | 13.2             | NG                  |
| 3/15/08 13:15  | 3/15/08 22:15  | 9.0           | 11.9       | 6.9              | 2.3               | 7.1              | NG                  |
| 4/3/08 9:15    | 4/3/08 19:00   | 9.8           | 18.3       | 5.6              | 2.9               | 9.1              | NG                  |
| 4/11/08 16:00  | 4/11/08 23:45  | 7.8           | 16.5       | 1.0              | 4.1               | 22.4             | NG                  |
| 4/27/08 21:00  | 4/28/08 8:15   | 11.3          | 25.7       | 0.3              | 3.8               | 19.3             | G                   |
| 5/11/08 15:00  | 5/12/08 3:30   | 12.5          | 14.0       | 24.1             | 1.4               | 3.0              | G                   |
| 6/3/08 8:30    | 6/3/08 13:45   | 5.2           | 12.4       | 27.2             | 3.6               | 18.3             | G                   |
| 7/28/08 15:15  | 7/28/08 17:00  | 1.8           | 10.9       | 0.0              | 6.2               | 18.3             | G                   |
| 7/30/08 13:45  | 7/30/08 15:30  | 1.8           | 16.8       | 10.9             | 9.6               | 35.6             | G                   |
| 7/31/08 5:45   | 7/31/08 12:00  | 6.2           | 19.1       | 27.7             | 4.2               | 14.2             | G                   |
| 8/26/08 14:30  | 8/27/08 2:00   | 11.5          | 29.5       | 0.0              | 3.1               | 10.2             | G                   |
| 10/8/08 6:30   | 10/8/08 13:30  | 7.0           | 13.0       | 0.0              | 2.6               | 6.1              | G                   |
| 10/24/08 13:45 | 10/25/08 3:00  | 13.3          | 19.3       | 0.0              | 1.9               | 5.1              | G                   |
| 11/13/08 2:30  | 11/13/08 7:30  | 5.0           | 18.3       | 0.5              | 4.9               | 29.5             | NG                  |
| 11/14/08 23:15 | 11/15/08 9:00  | 9.8           | 23.6       | 18.5             | 2.7               | 8.1              | NG                  |
| 11/24/08 11:00 | 11/24/08 21:15   | 10.2          | 12.2       | 2.0              | 1.5               | 5.1              | NG                  |
| 11/30/08 9:15  | 11/30/08 19:30   | 10.3          | 15.7       | 0.8              | 2.0               | 6.1              | NG                  |
| 12/10/08 0:30  | 12/10/08 8:00  | 7.5           | 33.3       | 0.8              | 4.3               | 9.1              | NG                  |
| 12/10/08 23:45 | 12/11/08 16:30   | 16.7          | 22.9       | 35.8             | 2.1               | 5.1              | NG                  |
| 12/15/08 13:15 | 12/15/08 21:00   | 7.8           | 13.2       | 39.9             | 1.8               | 4.1              | NG                  |
| 12/16/08 14:00 | 12/16/08 18:45   | 4.7           | 13.0       | 40.6             | 2.9               | 6.1              | NG                  |
|                | Hydrograph was plotted for storm event.  |               |            |                  |                   |                  |                     |
|                | Missing precipitation data was filled in from Little Millseat weather station. |               |            |                  |                   |                  |                     |

<sup>1</sup>G-Growing season runs from April 20 - October 26.

NG-Non-Growing season runs from October 27 - April 19.




Figure 2.9: Yearly precipitation sums and 27-year precipitation average. Data were taken from nearby Jackson County Airport for 1999 and 2000.

21.2 mm and in the second time period it was 21.3 mm. Average storm depths pre-harvest and post-harvest were 19.6 mm and 21.4 mm, respectively. Table 2.5 provides a summary of the yearly, pre-harvest and post-harvest average storm depths, maximum storm depths, cumulative storm depths, and storm events per period. For the years when storm intensity was evaluated (2002-2008), the years 2006 and 2007 had the highest average intensities at 6.0 mm hr<sup>-1</sup> and 6.6 mm hr<sup>-1</sup>, respectively, 2008 had the lowest at 3.4 mm hr<sup>-1</sup>. The years 2006 and 2007 also had the highest average maximum intensity at 20.7 mm hr<sup>-1</sup> and 20.5 mm hr<sup>-1</sup>, respectively, and 2008 was again the lowest at 11.3 mm hr<sup>-1</sup>. Table 2.6 provides a summary of the yearly, pre-harvest and post-harvest average storm intensities, maximum storm intensities, and storm events per period. No statistical significance was found between periods for the storm characteristics of depth, 5-day antecedent moisture, average storm intensity, and maximum storm intensity.

Table 2.5: Average storm depth  $\pm$  Std. dev. (mm), average 5-day antecedent depth  $\pm$  std. dev. (mm), maximum storm depth (mm), cumulative storm depth (mm), and sample size 1982-1993 and 2002-2008.

| Year         | Avg. Storm Depth<br>$\pm$ Std. Dev. | Avg. 5-day antecedent<br>rainfall $\pm$ Std. Dev. | Max Storm<br>Depth | Annual Cumulative<br>Storm Depth | Sample<br>size |
|--------------|-------------------------------------|---|--------------------|----------------------------------|----------------|
| 1982         | 19.7 $\pm$ 9.4                      | 14.7 $\pm$ 11.4                                   | 41.4               | 710.9                            | 36             |
| 1983         | 19.5 $\pm$ 8.8                      | 13.2 $\pm$ 15.1                                   | 48.5               | 585.7                            | 30             |
| 1984         | 22.4 $\pm$ 13.6                     | 17.6 $\pm$ 19.8                                   | 66.0               | 850.4                            | 38             |
| 1985         | 21.8 $\pm$ 9.2                      | 16.3 $\pm$ 13.5                                   | 51.3               | 720.6                            | 33             |
| 1986         | 18.4 $\pm$ 10.3                     | 16.4 $\pm$ 25.0                                   | 66.8               | 607.1                            | 33             |
| 1987         | 18.8 $\pm$ 5.4                      | 13.3 $\pm$ 13.5                                   | 35.1               | 639.3                            | 34             |
| 1988         | 21.4 $\pm$ 10.5                     | 10.1 $\pm$ 13.8                                   | 47.5               | 576.6                            | 27             |
| 1989         | 22.9 $\pm$ 11.3                     | 19.7 $\pm$ 17.4                                   | 66.5               | 1029.2                           | 45             |
| 1990         | 22.4 $\pm$ 12.4                     | 16.3 $\pm$ 21.6                                   | 80.0               | 872.2                            | 39             |
| 1991         | 23.3 $\pm$ 11.3                     | 18.3 $\pm$ 22.1                                   | 73.2               | 1001.8                           | 43             |
| 1992         | 20.4 $\pm$ 12.4                     | 10.0 $\pm$ 9.5                                    | 65.3               | 613.2                            | 30             |
| 1993         | 21.5 $\pm$ 11.6                     | 13.1 $\pm$ 15.6                                   | 57.9               | 796.5                            | 37             |
| 2002         | 22.1 $\pm$ 12.8                     | 16.9 $\pm$ 16.3                                   | 61.2               | 816.4                            | 37             |
| 2003         | 22.1 $\pm$ 18.1                     | 14.6 $\pm$ 14.9                                   | 105.7              | 773.9                            | 35             |
| 2004         | 25.9 $\pm$ 17.2                     | 21.6 $\pm$ 30.4                                   | 92.7               | 1141.5                           | 44             |
| 2005         | 18.9 $\pm$ 8.3                      | 10.9 $\pm$ 12.2                                   | 49.8               | 528.8                            | 28             |
| 2006         | 19.2 $\pm$ 10.0                     | 11.6 $\pm$ 10.8                                   | 61.7               | 537.0                            | 28             |
| 2007         | 20.0 $\pm$ 9.2                      | 9.7 $\pm$ 11.8                                    | 47.8               | 580.6                            | 29             |
| 2008         | 17.9 $\pm$ 6.2                      | 12.7 $\pm$ 14.4                                   | 33.3               | 446.5                            | 25             |
| Pre-harvest  | 19.6 $\pm$ 9.1                      | 14.0 $\pm$ 13.1                                   | 48.5               | 1296.67                          | 63             |
| Post-harvest | 21.4 $\pm$ 11.9                     | 15.2 $\pm$ 18.2                                   | 105.7              | 12531.598                        | 588            |

Table 2.6: Average storm intensity  $\pm$  standard deviation (mm/hr), average maximum intensity  $\pm$  standard deviation (mm/hr), and sample size 2002-2008.

| Year | Avg. Storm Intensity<br>$\pm$ Std. Dev. | Avg. Max Storm<br>Intensity $\pm$ Std. Dev. | Sample<br>size |
|------|---|---|----------------|
| 2002 | 4.4 $\pm$ 4.0                           | 15.4 $\pm$ 11.2                             | 37             |
| 2003 | 4.9 $\pm$ 3.5                           | 17.7 $\pm$ 12.7                             | 35             |
| 2004 | 5.3 $\pm$ 3.8                           | 18.7 $\pm$ 13.3                             | 44             |
| 2005 | 5.1 $\pm$ 4.0                           | 16.7 $\pm$ 14.3                             | 28             |
| 2006 | 6.0 $\pm$ 4.3                           | 20.7 $\pm$ 11.1                             | 28             |
| 2007 | 6.6 $\pm$ 6.6                           | 20.5 $\pm$ 18.9                             | 29             |
| 2008 | 3.4 $\pm$ 1.8                           | 11.3 $\pm$ 8.3                              | 25             |



### 2.3.2 Hydrograph Characteristics

From 1982-1993, 345 hydrographs were identified of which 190 (55.1%) were deemed acceptable for use in the analysis of hydrograph characteristics. For the period of 2002-2008, 210 hydrographs were identified of which 104 (49.5%) were acceptable. Hydrographs with similar storm depths (26.4 mm-49.8 mm) from each year of the study period are shown in Figures 2.10-2.28. The purpose of the following hydrographs is to show the discharge relationship between the study subcatchments and how that relationship varies from pre-harvest to post-harvest as well as its variability with regards to seasonality, storm intensity, and antecedent moisture conditions. Tables 2.2 and 2.3 detail all of these variables for all storms that occurred during the study period, including from which the storms the hydrographs plotted below represent. Due to this large amount of variability, statistical analyses of hydrograph characteristics were only performed on those hydrographs that were matched to a storm event across all watersheds.




Figure 2.10: Pre-harvest hydrographs for WSA, WSB, and WSC for a storm event (38.1 mm) on August 5, 1982 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.11: Pre-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.9 mm) on May 22, 1983 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.12: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.4 mm) on February 27, 1984 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.13: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (44.7 mm) on February 11, 1985 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.14: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (48.0 mm) on February 2, 1986 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.15: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (41.1 mm) on February 26, 1987 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.16: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (44.7 mm) on August 23, 1988 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.17: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.8 mm) on September 30, 1989 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.18: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.1 mm) on February 3, 1990 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.1: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (35.6 mm) on June 25, 1991 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.20: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (49.8 mm) on August 27, 1992 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.21: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (36.6 mm) on October 18, 1993 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.22: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (42.2 mm, 7.2 hr) on May 13, 2002 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.23: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (46.2 mm, 18.2 hr) on November 18, 2003 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.24: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (37.3 mm, 3 hr) on June 15, 2004 (G). WSA=control, WSB=BMP, and WSC=no BMPs.






Figure 2.25: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (30.5 mm, 6.7 hr) on May 19, 2005 (G). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.26: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (31.5 mm, 10.8 hr) on January 23, 2006 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.




Figure 2.27: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (33.3 mm, 9.2 hr) on November 14, 2007 (NG). WSA=control, WSB=BMP, and WSC=no BMPs




Figure 2.28: Post-harvest hydrographs for WSA, WSB, and WSC for a storm event (26.4 mm, 9.5 hr) on January 29, 2008 (NG). WSA=control, WSB=BMP, and WSC=no BMPs.

Results from the statistical analyses performed on the hydrograph parameters are shown in Tables 2.7-2.11. Table 2.7 details the results from the baseflow volume and stormflow volume statistical analyses. Results from the baseflow volume statistical analyses found significant differences between WSA and WSC in 1985 and 2007. The total yearly baseflow volume was 1.3 times as much in WSC in 1985 and 1.6 times as much in 2007. Significant statistical differences were also found between WSA and WSB in 1985; total yearly baseflow was 1.4 times greater than WSA. Finally, statistically significant differences were observed between WSB and WSC in 2007. Total yearly baseflow volume was 1.3 times higher in WSC compared to WSB.

More significant variations were seen in stormflow volume between the subcatchments throughout the study. Statistically significant differences in stormflow volume were seen between WSA and WSC in 1984, 1985, 2006 and 2007 with total yearly stormflow volume in WSC being greater by 2.3, 2.5, 3.1, and 1.4 times, respectively. Statistically significant differences were also seen in stormflow between WSA and WSB in 1985 where WSB was 2.2 times higher than WSA. Finally, statistically significant differences in stormflow were observed between WSB and WSC in 2007. Results indicated that WSC was statistically elevated over WSB; however, the total annual stormflow volume in WSB was slightly larger than WSC in 2007 ( $2,019.7 \text{ m}^3 \text{ ha}^{-1}$  vs.  $1,862.7 \text{ m}^3 \text{ ha}^{-1}$ ) (Table 2.11). This may indicate that the suitable hydrographs in 2007 did not accurately represent the stormflow volume relationship between WSB and WSC.

Table 2.8 details the results from the peak flow statistical analysis. No statistical significance differences were found between the subcatchments for peak flow during the study period.

Table 2.9 details the results from the storm volume as a percentage of rainfall statistical analysis. Significant statistical differences between WSA and WSC were observed in 1984, 1985, 1989, 2006, and 2007. Average [S/P]% for WSC during those years were 56%, 25%, 47%, 38%, and 14%, respectively, compared to average [S/P]% in WSA being 35%, 12%, 28%, 19%, and 7%. Statistical differences between WSA and WSB were only present in 1985;

Table 2.7: ANCOVA baseflow volume and stormflow volume results for WSA, WSB, and WSC 1982-1993 and 2002-2008.<sup>1</sup>

| Year | Baseflow Volume (m <sup>3</sup> ha <sup>-1</sup> ) |        |                 | Stormflow Volume (m <sup>3</sup> ha <sup>-1</sup> ) |          |         |
|------|--|--------|-----------------|---|----------|---------|
|      | WSA  | WSB    | WSC             | WSA   | WSB      | WSC     |
| 1982 | 420.5  | 283.6  | 531.1           | 1161.2  | 817.3    | 1155.6  |
| 1983 | 470.3  | 466.1  | 425.2           | 1198.1  | 1214.1   | 1164.9  |
| 1984 | 1212.5   | 1174.2 | 1175.0          | 2713.6b   | 2759.4ab | 3138.5a |
| 1985 | 208.2b   | 526.6a | 579.6a          | 608.8b  | 1212.3a  | 1286.1a |
| 1986 | 339.6  | 273.1  | 603.5           | 878.6   | 788.9    | 1450.2  |
| 1987 | 659.9  | 707.8  | 811.2           | 1395.1  | 1533.2   | 1789.5  |
| 1988 | 499.6  | 704.3  | 625.4           | 1130.5  | 923.1    | 1237.8  |
| 1989 | 1444.5   | 1766.3 | 2265.4          | 3205.7  | 3221.8   | 4234.5  |
| 1990 | 936.5  | 1287.9 | 1064.4          | 1828.1  | 2280.2   | 2342.0  |
| 1991 | 1142.7   | 1550.1 | 1234.1          | 2270.6  | 2867.6   | 2856.9  |
| 1992 | 806.7  | 595.6  | 655.4           | 1160.3  | 1182.9   | 1239.8  |
| 1993 | 254.0  | 379.3  | 345.2           | 624.6   | 827.6    | 822.8   |
| 2002 | 835.5  | 920.2  | -- <sup>2</sup> | 1444.3  | 1820.9   | --      |
| 2003 | 482.4  | 568.9  | --              | 1037.2  | 1176.9   | --      |
| 2004 | 1706.1   | 1715.3 | --              | 2898.0  | 3282.1   | --      |
| 2005 | 578.1  | 1271.6 | 574.2           | 620.6   | 774.3    | 758.4   |
| 2006 | 251.0  | 243.5  | 332.3           | 549.5b  | 835.1ab  | 1116.5a |
| 2007 | 58.5b  | 56.0b  | 113.6a          | 131.6b  | 142.5b   | 305.8a  |
| 2008 | 54.6   | 80.3   | 114.7           | 223.4   | 452.3    | 449.4   |

<sup>1</sup>Within each row, constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds. Constituents with no letters indicate no statistical significance.

<sup>2</sup>No data available.

Table 2.8: ANCOVA peak flow (Qp) ( $\text{m}^3 \text{s}^{-1} \text{ha}^{-1} \times 10^{-2}$ ) results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup>

| Year | Average Peak Flow |      |                 |
|------|-------------------|------|-----------------|
|      | WSA               | WSB  | WSC             |
| 1982 | 0.23              | 0.16 | 0.16            |
| 1983 | 0.14              | 0.12 | 0.11            |
| 1984 | 0.35              | 0.26 | 0.22            |
| 1985 | 0.07              | 0.09 | 0.09            |
| 1986 | 0.16              | 0.15 | 0.20            |
| 1987 | 0.10              | 0.09 | 0.09            |
| 1988 | 0.21              | 0.17 | 0.17            |
| 1989 | 0.34              | 0.26 | 0.29            |
| 1990 | 0.14              | 0.14 | 0.12            |
| 1991 | 0.16              | 0.15 | 0.13            |
| 1992 | 0.17              | 0.15 | 0.15            |
| 1993 | 0.12              | 0.15 | 0.42            |
| 2002 | 0.12              | 0.15 | -- <sup>2</sup> |
| 2003 | 0.13              | 0.15 | --              |
| 2004 | 0.37              | 0.33 | --              |
| 2005 | 0.02              | 0.01 | 0.03            |
| 2006 | 0.34              | 0.24 | 0.20            |
| 2007 | 0.03              | 0.03 | 0.05            |
| 2008 | 0.05              | 0.07 | 0.08            |

<sup>1</sup>Within each row, constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds. Constituents with no letters indicate no statistical significance.

<sup>2</sup>No data available.

Table 2.9: ANCOVA storm volume as a (%) of rainfall results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup>

| Year | Average [(S/P)%] |        |                |
|------|------------------|--------|----------------|
|      | WSA              | WSB    | WSC            |
| 1982 | 25.9             | 19.2   | 27.6           |
| 1983 | 21.5             | 23.5   | 22.5           |
| 1984 | 34.9b            | 48.5ab | 56.2a          |
| 1985 | 11.8b            | 23.0a  | 24.9a          |
| 1986 | 18.9             | 25.9   | 32.7           |
| 1987 | 22.7             | 25.2   | 28.1           |
| 1988 | 28.2             | 25.4   | 32.6           |
| 1989 | 35.4b            | 41.1ab | 47.2a          |
| 1990 | 26.7             | 32.9   | 34.4           |
| 1991 | 34.2             | 38.5   | 37.3           |
| 1992 | 18.0             | 20.4   | 20.7           |
| 1993 | 18.8             | 22.5   | 21.6           |
| 2002 | 35.6             | 45.8   | - <sup>2</sup> |
| 2003 | 29.7             | 33.4   | --             |
| 2004 | 32.9             | 37.2   | --             |
| 2005 | 17.8             | 21.1   | 17.9           |
| 2006 | 18.7b            | 28.9ab | 38.6a          |
| 2007 | 6.6b             | 6.9b   | 14.0a          |
| 2008 | 13.5             | 26.1   | 26.9           |

<sup>1</sup>Within each row, constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds. Constituents with no letters indicate no statistical significance.

<sup>2</sup>No data available.

average [S/P]% for WSB was 23% compared to WSA's 12%. Finally, statistically significant differences between WSB and WSC were observed in 2007 with average [S/P]% for WSC being 14% compared to 7% in WSB.

Table 2.10 details the results from the time to peak statistical analysis. No statistical significance differences were found between the subcatchments for time to peak during the study period.

Table 2.11 details the results from the CN statistical analysis. Statistically significant differences between WSA and WSC were observed in 1984, 1985, 2006, and 2007 with average CNs ( $\lambda=0.05$ ) in WSC of 91, 79, 88, and 81 respectively; average CNs in WSA were 82, 70, 80, and 75, respectively. Statistically significant differences between WSA and WSB only occurred in 1985 with WSB having an average CN of 78 while WSA's was 70. Finally, statistically significant differences between WSB and WSC were observed in 2007; the average CN for WSC in 2007 was 81 while WSB's was 74.

Table 2.10: ANCOVA time to peak (Tp) results for WSA, WSB, and WSC 2002-2008. <sup>1</sup>

| Year | Average Time to Peak (hr) |      |      |
|------|---------------------------|------|------|
|      | WSA                       | WSB  | WSC  |
| 2002 | 14.4                      | 12.1 |      |
| 2003 | 10.1                      | 12.9 |      |
| 2004 | 10.5                      | 10.1 |      |
| 2005 | 6.0                       | 6.1  | 8.9  |
| 2006 | 11.2                      | 9.8  | 10.7 |
| 2007 | 4.9                       | 9.8  | 6.5  |
| 2008 | 10.5                      | 7.3  | 8.9  |

<sup>1</sup>Within each row, constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds.

<sup>2</sup>No data available.



Table 2.11: ANCOVA CN ( $\lambda=0.2$  &  $\lambda=0.05$ ) results for WSA, WSB, and WSC 1982-1993 and 2002-2008. <sup>1</sup>

| Year | Average CN ( $\lambda=0.2$ ) |      |                 | Average CN ( $\lambda=0.05$ ) |      |     |
|------|------------------------------|------|-----------------|-------------------------------|------|-----|
|      | WSA                          | WSB  | WSC             | WSA                           | WSB  | WSC |
| 1982 | 83                           | 81   | 82              | 77                            | 75   | 76  |
| 1983 | 82                           | 83   | 83              | 76                            | 77   | 76  |
| 1984 | 86b                          | 91ab | 93a             | 81b                           | 88ab | 91a |
| 1985 | 78b                          | 84a  | 84a             | 70b                           | 78a  | 79a |
| 1986 | 82                           | 86   | 88              | 76                            | 80   | 84  |
| 1987 | 82                           | 84   | 86              | 76                            | 78   | 81  |
| 1988 | 82                           | 82   | 85              | 76                            | 76   | 80  |
| 1989 | 85                           | 87   | 89              | 80                            | 82   | 85  |
| 1990 | 86                           | 87   | 88              | 81                            | 83   | 84  |
| 1991 | 86                           | 87   | 86              | 80                            | 83   | 81  |
| 1992 | 80                           | 82   | 82              | 73                            | 75   | 76  |
| 1993 | 77                           | 80   | 80              | 68                            | 73   | 73  |
| 2002 | 89                           | 91   | -- <sup>2</sup> | 85                            | 88   | --  |
| 2003 | 87                           | 88   | --              | 82                            | 84   | --  |
| 2004 | 87                           | 88   | --              | 83                            | 84   | --  |
| 2005 | 85                           | 85   | 86              | 80                            | 80   | 81  |
| 2006 | 85b                          | 88ab | 91a             | 80b                           | 84ab | 88a |
| 2007 | 81b                          | 81b  | 86a             | 75b                           | 74b  | 81a |
| 2008 | 83                           | 87   | 86              | 76                            | 82   | 82  |

<sup>1</sup>Within each row, constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds. Constituents with no letters indicate no statistical significance.

<sup>2</sup> No data available.

### 2.3.2.1 Baseflow, Storm Flow and Total Flow Volumes

Figures 2.29-2.34 and Tables 2.12-2.14 show annual, annual growing, and annual non-growing season baseflow, stormflow, and total flow (sum of baseflow and stormflow) volumes for WSA, WSB, and WSC. The year 2004 had the highest cumulative baseflow volume, stormflow volume, and total flow volume for WSA and WSB with WSA having volumes of  $4,312 \text{ m}^3 \text{ ha}^{-1}$ ,  $6,030 \text{ m}^3 \text{ ha}^{-1}$ , and  $10,342 \text{ m}^3 \text{ ha}^{-1}$ , respectively; WSB had volumes of  $4,547 \text{ m}^3 \text{ ha}^{-1}$ ,  $6,989 \text{ m}^3 \text{ ha}^{-1}$ , and  $11,536 \text{ m}^3 \text{ ha}^{-1}$ , respectively. Note that 2004 was the year with the highest cumulative precipitation during the study period; WSC was not analyzed in 2004. The year with the highest baseflow volume for WSC was 1985 with a volume of  $3,516 \text{ m}^3 \text{ ha}^{-1}$ , and the highest stormflow volume and total flow volume in WSC occurred in 1989 with volumes of  $8,652 \text{ m}^3 \text{ ha}^{-1}$  and  $10,789 \text{ m}^3 \text{ ha}^{-1}$ , respectively. During the study period, 1989 was the year with the second highest cumulative precipitation. The lowest baseflow volume, stormflow volume, and total flow volume for WSA occurred in 2008 with volumes of  $490 \text{ m}^3 \text{ ha}^{-1}$ ,  $1,192 \text{ m}^3 \text{ ha}^{-1}$ , and  $1,682 \text{ m}^3 \text{ ha}^{-1}$ , respectively. The lowest baseflow volume and total flow volume for WSB also occurred in 2008 with volumes of  $819 \text{ m}^3 \text{ ha}^{-1}$  and  $2,769 \text{ m}^3 \text{ ha}^{-1}$ , respectively. The lowest stormflow volume in WSB occurred in 1982 with a volume of  $1,717 \text{ m}^3 \text{ ha}^{-1}$ . The lowest baseflow volume, stormflow volume, and total flow volume in WSC occurred in 1983 with volumes of  $1,064 \text{ m}^3 \text{ ha}^{-1}$ ,  $1,776 \text{ m}^3 \text{ ha}^{-1}$ , and  $2,840 \text{ m}^3 \text{ ha}^{-1}$ , respectively. The years 1982, 1983 and 2008 were all periods with below average cumulative precipitation.

Results from the statistical analysis found large significant differences in baseflow volume between WSA and WSC two years post-harvest in 1985 and 24 years post-harvest in 2007. Statistically significant differences in stormflow volume were observed between WSA and WSC in 1984, 1985, 2006, and 2007. Results from the precipitation analysis showed that 2006 and 2007 were years with high average storm intensity and high average maximum storm intensity (Table 2.4-2.6). Although these years were found not to be statistically higher than the period average intensity and average maximum intensity, they were found to be statistically higher than the period minimum occurring in 2008.

Table 2.12: Yearly WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area<sup>1</sup> 1982-1993 and 2002-2008.

| Year | WSA  |   |   | WSB  |   |   | WSC  |   |   |
|------|--|---|---|--|---|---|--|---|---|
|      | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| 1982 | 1928.8   | 1598.5  | 3527.2  | 1510.2   | 1716.7  | 3226.9  | 1116.8   | 2426.3  | 3543.1  |
| 1983 | 1374.1   | 1596.1  | 2970.2  | 950.5  | 2013.9  | 2964.4  | 1064.4   | 1776.3  | 2840.7  |
| 1984 | 2445.1   | 2923.5  | 5368.6  | 1903.0   | 5640.4  | 7543.4  | 2212.2   | 6607.3  | 8819.6  |
| 1985 | 2705.2   | 1596.7  | 4301.9  | 3766.7   | 3473.7  | 7240.4  | 3516.4   | 3969.3  | 7485.7  |
| 1986 | 1953.1   | 1705.1  | 3658.1  | 1791.4   | 3245.3  | 5036.6  | 1919.5   | 3541.8  | 5461.3  |
| 1987 | 1896.1   | 1786.5  | 3682.6  | 1502.7   | 2636.4  | 4139.1  | 1270.1   | 3635.5  | 4905.6  |
| 1988 | 1743.3   | 1610.5  | 3353.9  | 1107.5   | 2381.8  | 3489.3  | 1507.5   | 2800.8  | 4308.4  |
| 1989 | 4052.4   | 4352.7  | 8405.1  | 1533.5   | 7759.4  | 9292.9  | 2136.8   | 8652.4  | 10789.3                                       |
| 1990 | 2469.8   | 2928.2  | 5398.0  | 1371.7   | 5097.2  | 6468.9  | 1798.4   | 5088.9  | 6887.3  |
| 1991 | 4163.4   | 3643.4  | 7806.8  | 1855.6   | 6578.3  | 8433.9  | 1879.3   | 6180.6  | 8059.9  |
| 1992 | 2994.2   | 2117.8  | 5112.0  | 1520.6   | 3745.7  | 5266.3  | 1852.5   | 4103.3  | 5955.8  |
| 1993 | 2880.9   | 1726.8  | 4607.7  | 2062.8   | 3355.6  | 5418.4  | 1758.2   | 3634.2  | 5392.5  |
| 2002 | 2163.3   | 3216.8  | 5380.1  | 2587.0   | 4308.8  | 6895.8  | -- <sup>2</sup>                                | --  | --  |
| 2003 | 2457.3   | 3828.9  | 6286.2  | 3713.0   | 6090.9  | 9803.9  | --   | --  | --  |
| 2004 | 4312.3   | 6030.1  | 10342.4                                       | 4546.8   | 6989.1  | 11535.9                                       | --   | --  | --  |
| 2005 | 2572.9   | 2698.0  | 5270.9  | 2621.6   | 2650.6  | 5272.2  | 3069.2   | 4741.9  | 7811.1  |
| 2006 | 824.4  | 1502.3  | 2326.7  | 1183.3   | 2459.5  | 3642.7  | 2219.4   | 4649.0  | 6868.4  |
| 2007 | 651.6  | 1345.0  | 1996.5  | 823.4  | 2019.7  | 2843.2  | 1032.0   | 1862.7  | 2894.7  |
| 2008 | 490.4  | 1191.8  | 1682.3  | 819.1  | 1950.3  | 2769.3  | 1033.2   | 2076.5  | 3109.7  |

<sup>1</sup>Drainage areas: WSA (16.1 ha), WSB (11.2 ha), and WSC (10.5 ha).

<sup>2</sup>No data available.




Figure 2.29: Yearly baseflow and stormflow WSA, WSB, and WSC (1982-1993).




Figure 2.30: Yearly baseflow and stormflow WSA, WSB, and WSC (2002-2008).

Table 2.13: Growing season WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area<sup>1</sup> 1982-1993 and 2002-2008.

| Year | WSA  |   |   | WSB  |   |   | WSC  |   |   |
|------|--|---|---|--|---|---|--|---|---|
|      | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| 1982 | 267.2  | 115.9   | 383.2   | 276.7  | 163.9   | 440.6   | 423.9  | 132.0   | 555.9   |
| 1983 | 683.8  | 1011.5  | 1695.2  | 345.9  | 1274.1  | 1620.0  | 379.8  | 1196.5  | 1576.4  |
| 1984 | 1067.6   | 1374.4  | 2442.0  | 631.4  | 2758.9  | 3390.3  | 908.4  | 3469.3  | 4377.8  |
| 1985 | 277.3  | 140.4   | 417.6   | 747.2  | 805.0   | 1552.3  | 975.5  | 837.8   | 1813.3  |
| 1986 | 292.4  | 71.6  | 363.9   | 375.0  | 343.9   | 718.9   | 546.0  | 358.1   | 904.1   |
| 1987 | 601.5  | 75.3  | 676.8   | 505.0  | 243.1   | 748.1   | 743.6  | 410.4   | 1153.9  |
| 1988 | 705.1  | 339.9   | 1045.0  | 400.0  | 393.0   | 793.0   | 529.0  | 600.6   | 1129.5  |
| 1989 | 1708.9   | 2047.2  | 3756.1  | 697.0  | 3682.8  | 4379.9  | 901.9  | 4557.3  | 5459.2  |
| 1990 | 835.4  | 454.4   | 1289.8  | 552.6  | 939.1   | 1491.7  | 759.6  | 1265.1  | 2024.7  |
| 1991 | 952.5  | 776.0   | 1728.5  | 692.0  | 1528.9  | 2220.9  | 719.5  | 1255.5  | 1975.0  |
| 1992 | 1073.4   | 734.3   | 1807.7  | 525.0  | 1153.6  | 1678.5  | 759.7  | 1266.0  | 2025.7  |
| 1993 | 795.7  | 471.0   | 1266.7  | 854.5  | 949.1   | 1803.5  | 807.7  | 843.0   | 1650.7  |
| 2002 | 1431.2   | 1660.7  | 3091.9  | 1759.4   | 2102.6  | 3862.0  | -- <sup>2</sup>                                | --  | --  |
| 2003 | 893.3  | 1333.3  | 2226.6  | 843.8  | 1280.0  | 2123.9  | --   | --  | --  |
| 2004 | 2086.2   | 2470.9  | 4557.1  | 2768.8   | 3892.4  | 6661.1  | --   | --  | --  |
| 2005 | 1883.8   | 1384.7  | 3268.6  | 605.3  | 1201.5  | 1806.8  | 1295.1   | 1165.8  | 2460.8  |
| 2006 | 404.2  | 627.2   | 1031.4  | 1002.6   | 1935.9  | 2938.5  | 792.4  | 1591.5  | 2383.9  |
| 2007 | 302.7  | 562.0   | 864.7   | 576.0  | 1143.6  | 1719.7  | 277.4  | 303.7   | 581.0   |
| 2008 | 251.5  | 367.0   | 618.6   | 40.8   | 320.3   | 361.0   | 282.8  | 287.5   | 570.3   |

<sup>1</sup>Drainage areas: WSA (16.1 ha), WSB (11.2 ha), and WSC (10.5 ha).

<sup>2</sup>No data available.




Figure 2.31: Growing season stormflow and baseflow WSA, WSB, and WSC (1982-1993).




Figure 2.32: Growing Season baseflow and stormflow WSA, WSB, and WSC (2002-2008).

Table 2.14: Non-growing season WSA, WSB, and WSC stormflow, baseflow, and total flow normalized by drainage area<sup>1</sup> 1982-1993 and 2002-2008.

| Year | WSA  |   |   | WSB  |   |   | WSC  |   |   |
|------|--|---|---|--|---|---|--|---|---|
|      | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Total Q<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| 1982 | 1661.5   | 1482.5  | 3144.1  | 1233.5   | 1552.8  | 2786.3  | 692.9  | 2294.3  | 2987.2  |
| 1983 | 690.3  | 584.7   | 1275.0  | 604.6  | 739.8   | 1344.4  | 684.5  | 579.8   | 1264.3  |
| 1984 | 1377.5   | 1549.1  | 2926.6  | 1271.6   | 2881.6  | 4153.2  | 1303.8   | 3138.0  | 4441.8  |
| 1985 | 2428.0   | 1456.4  | 3884.3  | 3019.4   | 2668.7  | 5688.1  | 2540.9   | 3131.5  | 5672.4  |
| 1986 | 1660.7   | 1633.5  | 3294.2  | 1416.4   | 2901.3  | 4317.7  | 1373.5   | 3183.7  | 4557.2  |
| 1987 | 1294.6   | 1711.3  | 3005.9  | 997.7  | 2393.3  | 3391.0  | 526.5  | 3225.2  | 3751.7  |
| 1988 | 1038.2   | 1270.7  | 2308.9  | 707.5  | 1988.9  | 2696.3  | 978.5  | 2200.3  | 3178.8  |
| 1989 | 2343.5   | 2305.4  | 4648.9  | 836.5  | 4076.5  | 4913.0  | 1235.0   | 4095.1  | 5330.1  |
| 1990 | 1634.4   | 2473.8  | 4108.3  | 819.1  | 4158.1  | 4977.2  | 1038.7   | 3823.8  | 4862.5  |
| 1991 | 3210.9   | 2867.4  | 6078.3  | 1163.6   | 5049.4  | 6213.0  | 1159.8   | 4925.2  | 6084.9  |
| 1992 | 1920.8   | 1383.6  | 3304.4  | 995.6  | 2592.2  | 3587.8  | 1092.8   | 2837.2  | 3930.0  |
| 1993 | 2085.2   | 1255.8  | 3341.1  | 1208.3   | 2406.6  | 3614.9  | 950.5  | 2791.3  | 3741.8  |
| 2002 | 732.1  | 1556.1  | 2288.2  | 827.6  | 2206.2  | 3033.8  | -- <sup>2</sup>                                | --  | --  |
| 2003 | 1564.0   | 2495.6  | 4059.6  | 2869.2   | 4810.9  | 7680.1  | --   | --  | --  |
| 2004 | 2226.1   | 3559.2  | 5785.3  | 1778.1   | 3096.7  | 4874.8  | --   | --  | --  |
| 2005 | 689.1  | 1313.3  | 2002.3  | 2016.4   | 1449.0  | 3465.4  | 1774.2   | 3576.1  | 5350.3  |
| 2006 | 420.2  | 875.1   | 1295.2  | 180.6  | 523.5   | 704.2   | 1427.0   | 3057.5  | 4484.5  |
| 2007 | 348.9  | 783.0   | 1131.9  | 247.4  | 876.1   | 1123.5  | 754.6  | 1559.1  | 2313.6  |
| 2008 | 212.1  | 740.1   | 952.3   | 778.3  | 1630.0  | 2408.3  | 750.4  | 1789.0  | 2539.4  |

<sup>1</sup>Drainage areas: WSA (16.1 ha), WSB (11.2 ha), and WSC (10.5 ha).

<sup>2</sup>No data available.




Figure 2.33: Non-growing season baseflow and stormflow WSA, WSB, and WSC (1982-1993).




Figure 2.2: Non-growing season baseflow and stormflow WSA, WSB, and WSC (2002-2008).



This could explain the large statistical differences in baseflow volume and stormflow volume between WSA and WSC 23 and 24 years post-harvest following a large time gap of no statistical differences. In contrast, statistically significant differences in baseflow volume and stormflow volume between WSA and WSB only occurred in 1985.

Pre-harvest WSC had a yearly baseflow volume that was on average 0.7 times as much as WSA and a yearly stormflow volume that was 1.3 times higher. Similar values were observed in WSB with a yearly baseflow volume that was on average 0.7 times as much as WSA and a yearly stormflow volume that was 1.2 times higher. The post-harvest (1984-1993 and 2005-2008) yearly baseflow volume in WSC was on average 1.1 times higher than WSA and yearly stormflow volume was 2.0 times higher. For the same post-harvest time period the yearly baseflow volume in WSB was on average 0.9 times as much and yearly stormflow volume was 1.7 times higher.

During the growing season, the differences that were previously explored were even larger (Figure 2.31 and 2.32; Table 2.13). Pre-harvest WSC had a growing season baseflow volume that was on average 0.7 times as much as WSA and a yearly stormflow volume that was 1.8 times higher. Similar values were observed in WSB with a growing season baseflow volume that was on average 0.6 times as much as WSA and a yearly stormflow volume that was 1.6 times higher. The yearly baseflow volume and stormflow volume in WSC for the same post-harvest time period during the growing season was on average 1.2 and 2.5 times respectively, higher than WSA. The baseflow volume and stormflow volume for WSB during the same time period and growing period were 1.0 and 2.4 times respectively, higher than WSA.

Differences in baseflow volume and stormflow volume were also observed during the non-growing season (Figure 2.33 and 2.34; Table 2.14). Pre-harvest WSC had a non-growing season baseflow volume that was on average 0.7 times as much as WSA and a yearly stormflow volume that was 1.3 times higher. Similar values were observed in WSB with a growing season baseflow volume that was on average 0.8 times as much as WSA and a yearly stormflow volume

that was 1.2 times higher. The yearly baseflow volume and stormflow volume in WSC for the same post-harvest time period during the non-growing season was 1.3 and 2.1 times, respectively higher than WSA. The baseflow volume and stormflow volume for WSB during the same time period and growing period were 1.0 and 1.6 times, respectively higher than WSA.

These results show that the stormflow volume in WSB is on average more aligned with the control during the non-growing season than during the growing-season but has not quite reached pre-harvest conditions. A similar study conducted in the Allegheny Mountains observed seasonality effects with statistically significant elevated peak flows and stormflow volumes six years post-harvest during the growing season (Kochenderfer et al., 1997). Seasonality had very little effect on WSC; stormflow volumes were somewhat more aligned with the control during the non-growing season while baseflow volume experienced no change. This would indicate that even though the watersheds have been allowed to regrow for 25 years, the role of vegetation in the hydrologic cycle still does not match the control watershed. Another important result from this analysis was that the stormflow volumes in WSB and WSC remained elevated at the conclusion of the study in 2008. Variations in temporal rainfall distribution have been shown to play a major role in determining storm runoff volumes. Results from (Warner et al., 2010) found that rainfall events with a temporal distribution classified as intense ( $\mu=14.8$  mm/hr) produced larger storm runoff volumes and subsequently elevated CNs in the Appalachian region of Kentucky compared to rainfall events with lower intensities. Similar results were found during this study with the highest amount of variability in runoff volumes between the harvested watersheds and the control occurring during years with the highest storm intensities: 2006 and 2007 average storm intensity was 6 and 6.6 mm hr<sup>-1</sup>, respectively, and average maximum intensity was 20.7 and 20.5 mm hr<sup>-1</sup>, respectively. Comparatively, 2008 had the lowest average storm intensity at 3.4 mm hr<sup>-1</sup> and lowest average maximum storm intensity at 11.3 mm hr<sup>-1</sup> (Table 2.6). Finally, baseflow volumes were less impacted by treatment than stormflow volumes; nonetheless, statistical variation was present two years after harvest for

both WSB and WSC and again in 2007 for WSC, a year with high intensity storms.

#### 2.3.2.2 Peak Discharge

Figures 2.35-2.40 and Table 2.15 show annual, growing season, and non-growing season average peak flow for WSA, WSB, and WSC.

No statistical differences were recorded during the study period. Graphically, there only seems to be marginal differences between the watersheds and no discernible trends in the data. The main difference that can be seen from the graph is in 1993 when the average storm peak flow for WSC was much higher. This difference mainly came from one event (July 13, 1993) when the peak flow in WSC was 15 times higher than WSA and 16 times higher than WSB. In the second time period, these differences become even less noticeable and the watersheds have nearly identical results for most years.

These results are surprising because higher peak flows were expected to accompany higher stormflows but that was hardly the case. For comparison, the pre-harvest average peak flow for WSA, WSB, and WSC was 1.9, 1.4, and 1.3 ( $\text{m}^3 \text{ s}^{-1} \text{ ha}^{-1}$ ), respectively. Due to their respective areas, these values seem reasonable. Average peak flow post-harvest for WSA, WSB, and WSC was 1.8, 1.6, and 1.7 ( $\text{m}^3 \text{ s}^{-1} \text{ ha}^{-1}$ ), respectively. These differences seem pretty insignificant, and it would seem that harvest might have had some small effect on peak flow. With limited pre-harvest data, not much emphasis should be placed on this result. Witness accounts from the first time period (1982-1993) noted logging debris left in the stream in WSC following harvest was trapping large amounts of sediment. This effectively increased frictional effects in the stream channel, which in turn would have decreased stream velocity. This may explain why increases to stream velocity were not observed in WSC as expected. It was also observed that a large storm event, perhaps the event on July 13, 1993, washed out all of this debris and sediment and stripped the channel in WSC to bedrock. This may explain the large elevated peak flow in WSC during the event on July 13, 1993. Differences between the watersheds in  $Q_p$  due to seasonality were not observed during the analyzed time periods.

Table 2.15: WSA, WSB, and WSC average yearly, growing, and non-growing season peak flow ( $Q_p$ )  $\pm$  standard deviation ( $m^3 ha^{-1} s^{-1} \times 10^{-2}$ ) normalized by drainage area<sup>1</sup> 1982-1993 and 2002-2008.

| Year | WSA             |                 |                   | WSB             |                 |                   | WSC             |                 |                   |
|------|-----------------|-----------------|-------------------|-----------------|-----------------|-------------------|-----------------|-----------------|-------------------|
|      | Qp Year         | QP Growing      | Qp Non-Growing    | Qp Year         | QP Growing      | Qp Non-Growing    | Qp Year         | QP Growing      | Qp Non-Growing    |
| 1982 | 0.23 $\pm$ 0.46 | 0.09 $\pm$ 0.14 | 0.36 $\pm$ 0.61   | 0.16 $\pm$ 0.25 | 0.06 $\pm$ 0.09 | 0.24 $\pm$ 0.32   | 0.16 $\pm$ 0.29 | 0.06 $\pm$ 0.06 | 0.25 $\pm$ 0.37   |
| 1983 | 0.14 $\pm$ 0.22 | 0.18 $\pm$ 0.26 | 0.06 $\pm$ 0.04   | 0.12 $\pm$ 0.16 | 0.15 $\pm$ 0.19 | 0.06 $\pm$ 0.04   | 0.11 $\pm$ 0.16 | 0.14 $\pm$ 0.18 | 0.04 $\pm$ 0.02   |
| 1984 | 0.35 $\pm$ 0.79 | 0.55 $\pm$ 1.14 | 0.17 $\pm$ 0.13   | 0.26 $\pm$ 0.30 | 0.28 $\pm$ 0.39 | 0.24 $\pm$ 0.21   | 0.22 $\pm$ 0.26 | 0.24 $\pm$ 0.31 | 0.21 $\pm$ 0.21   |
| 1985 | 0.07 $\pm$ 0.04 | 0.06 $\pm$ 0.05 | 0.09 $\pm$ 0.01   | 0.09 $\pm$ 0.06 | 0.07 $\pm$ 0.05 | 0.15 $\pm$ 0.05   | 0.09 $\pm$ 0.06 | 0.07 $\pm$ 0.05 | 0.14 $\pm$ 0.07   |
| 1986 | 0.16 $\pm$ 0.20 | 0.05 $\pm$ 0.03 | 0.29 $\pm$ 0.24   | 0.15 $\pm$ 0.19 | 0.03 $\pm$ 0.01 | 0.30 $\pm$ 0.21   | 0.20 $\pm$ 0.30 | 0.04 $\pm$ 0.02 | 0.38 $\pm$ 0.39   |
| 1987 | 0.10 $\pm$ 0.12 | 0.06 $\pm$ 0.06 | 0.13 $\pm$ 0.15   | 0.09 $\pm$ 0.12 | 0.03 $\pm$ 0.02 | 0.13 $\pm$ 0.14   | 0.09 $\pm$ 0.10 | 0.05 $\pm$ 0.03 | 0.13 $\pm$ 0.12   |
| 1988 | 0.21 $\pm$ 0.28 | 0.10 $\pm$ 0.15 | 0.31 $\pm$ 0.34   | 0.17 $\pm$ 0.23 | 0.10 $\pm$ 0.19 | 0.23 $\pm$ 0.26   | 0.17 $\pm$ 0.20 | 0.13 $\pm$ 0.21 | 0.20 $\pm$ 0.20   |
| 1989 | 0.34 $\pm$ 0.44 | 0.38 $\pm$ 0.54 | 0.28 $\pm$ 0.22   | 0.26 $\pm$ 0.29 | 0.26 $\pm$ 0.35 | 0.25 $\pm$ 0.17   | 0.29 $\pm$ 0.33 | 0.33 $\pm$ 0.41 | 0.23 $\pm$ 0.16   |
| 1990 | 0.14 $\pm$ 0.11 | 0.07 $\pm$ 0.06 | 0.21 $\pm$ 0.09   | 0.14 $\pm$ 0.13 | 0.04 $\pm$ 0.03 | 0.24 $\pm$ 0.10   | 0.12 $\pm$ 0.09 | 0.05 $\pm$ 0.04 | 0.19 $\pm$ 0.08   |
| 1991 | 0.16 $\pm$ 0.10 | 0.13 $\pm$ 0.09 | 0.18 $\pm$ 0.12   | 0.15 $\pm$ 0.12 | 0.11 $\pm$ 0.11 | 0.20 $\pm$ 0.13   | 0.13 $\pm$ 0.09 | 0.11 $\pm$ 0.09 | 0.14 $\pm$ 0.09   |
| 1992 | 0.17 $\pm$ 0.23 | 0.18 $\pm$ 0.25 | 0.15 $\pm$ 0.06   | 0.15 $\pm$ 0.21 | 0.14 $\pm$ 0.23 | 0.15 $\pm$ 0.07   | 0.15 $\pm$ 0.23 | 0.16 $\pm$ 0.26 | 0.12 $\pm$ 0.05   |
| 1993 | 0.12 $\pm$ 0.09 | 0.11 $\pm$ 0.09 | 0.22 <sup>2</sup> | 0.15 $\pm$ 0.12 | 0.13 $\pm$ 0.11 | 0.36 <sup>2</sup> | 0.42 $\pm$ 0.69 | 0.39 $\pm$ 0.72 | 0.74 <sup>2</sup> |
| 2002 | 0.24 $\pm$ 0.45 | 0.35 $\pm$ 0.60 | 0.12 $\pm$ 0.11   | 0.24 $\pm$ 0.41 | 0.33 $\pm$ 0.55 | 0.15 $\pm$ 0.15   | -- <sup>3</sup> | --              | --                |
| 2003 | 0.12 $\pm$ 0.16 | 0.11 $\pm$ 0.15 | 0.13 $\pm$ 0.19   | 0.13 $\pm$ 0.20 | 0.11 $\pm$ 0.19 | 0.15 $\pm$ 0.23   | --              | --              | --                |
| 2004 | 0.31 $\pm$ 0.31 | 0.27 $\pm$ 0.32 | 0.37 $\pm$ 0.32   | 0.31 $\pm$ 0.32 | 0.30 $\pm$ 0.37 | 0.33 $\pm$ 0.23   | --              | --              | --                |
| 2005 | 0.25 $\pm$ 0.48 | 0.29 $\pm$ 0.51 | 0.02 $\pm$ 0.02   | 0.22 $\pm$ 0.45 | 0.26 $\pm$ 0.47 | 0.01 $\pm$ 0.01   | 0.24 $\pm$ 0.53 | 0.27 $\pm$ 0.56 | 0.03 $\pm$ 0.00   |
| 2006 | 0.18 $\pm$ 0.24 | 0.12 $\pm$ 0.13 | 0.34 $\pm$ 0.41   | 0.13 $\pm$ 0.15 | 0.10 $\pm$ 0.11 | 0.24 $\pm$ 0.22   | 0.15 $\pm$ 0.15 | 0.13 $\pm$ 0.14 | 0.20 $\pm$ 0.17   |
| 2007 | 0.05 $\pm$ 0.06 | 0.08 $\pm$ 0.07 | 0.03 $\pm$ 0.02   | 0.03 $\pm$ 0.02 | 0.03 $\pm$ 0.03 | 0.03 $\pm$ 0.02   | 0.07 $\pm$ 0.06 | 0.10 $\pm$ 0.07 | 0.05 $\pm$ 0.02   |
| 2008 | 0.04 $\pm$ 0.03 | 0.02 $\pm$ 0.02 | 0.05 $\pm$ 0.03   | 0.05 $\pm$ 0.04 | 0.01 $\pm$ 0.01 | 0.07 $\pm$ 0.02   | 0.07 $\pm$ 0.05 | 0.04 $\pm$ 0.04 | 0.08 $\pm$ 0.05   |

<sup>1</sup>Drainage areas: WSA (16.1 ha), WSB (11.2 ha), and WSC (10.5 ha).

<sup>2</sup>One storm suitable for analysis.

<sup>3</sup>No data available.




Figure 2.35: Average Storm Peak Flow WSA, WSB, WSC (1982-1993).




Figure 2.36: Average Storm Peak Flow WSA, WSB, WSC (2002-2008).




Figure 2.37: Growing Season Average Storm Peak Flow WSA, WSB, WSC (1982-1993).




Figure 2.38: Growing Season Average Storm Peak Flow WSA, WSB, WSC (2002-2008).




Figure 2.39: Non-Growing Season Average Storm Peak Flow WSA, WSB, WSC (1982-1993).




Figure 2.40: Non-Growing Season Average Storm Peak Flow WSA, WSB, WSC (2002-2008).

### 2.3.2.3 Stormflow Volume as a Percentage of Rainfall

Results from comparisons of the average storm volume as a percentage of rainfall ([S/P]%) across watersheds are presented in Figures 2.41-2.46 and Tables 2.16-2.18. The yearly maximum average [S/P]% for WSA occurred in 1989 with a value of 35.4% and the yearly maximum average [S/P]% for WSB and WSC occurred directly after harvest in 1984 with values of 48.5% and 65.2%, respectively.

Significant statistical differences between WSA and WSC were observed in 1984, 1985, 1989, 2006, and 2007. Average [S/P]% for WSC was 56%, 25%, 47%, 38%, and 14%, respectively, compared to average [S/P]% in WSA being 35%, 12%, 28%, 19%, and 7%. Statistical differences between WSA and WSB were only present in 1985; average [S/P]% for WSB was 23% compared to WSA's 12%. Finally, statistically significant differences between WSB and WSC were only observed in 2007 with average [S/P]% for WSC being 14% compared to 7% in WSB.

These results, along with the stormflow results, show that WSB and WSC became more responsive to precipitation following harvest, which in turn led to larger stormflow volumes. The largest of these effects were seen in the first few years following harvest. By the end of the first time period, hardly any variation existed. Oddly enough, when monitoring was resumed in WSC (2005), large spikes in storm volumes were observed from 2006-2008. This again was most likely caused by high intensity rainfall events. Seasonal variations followed a similar pattern as the baseflow and stormflow volume results already explored. Larger variations between the harvested watersheds and WSA were present during the growing season when vegetation plays an active role in the hydrologic cycle. Less variation was present during the non-growing season when vegetation was dormant.



Table 2.16: Yearly average storm volume as a (%) of rainfall  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year | WSA             | WSB             | WSC             |
|------|-----------------|-----------------|-----------------|
| 1982 | 25.9 $\pm$ 30.2 | 19.2 $\pm$ 20.6 | 27.6 $\pm$ 33.7 |
| 1983 | 21.5 $\pm$ 22.3 | 23.5 $\pm$ 24.6 | 22.5 $\pm$ 24.4 |
| 1984 | 34.9 $\pm$ 28.9 | 48.5 $\pm$ 26.9 | 56.2 $\pm$ 29.9 |
| 1985 | 11.8 $\pm$ 16.8 | 23.0 $\pm$ 25.9 | 24.9 $\pm$ 26.8 |
| 1986 | 18.9 $\pm$ 25.9 | 25.9 $\pm$ 29.7 | 32.7 $\pm$ 34.2 |
| 1987 | 22.7 $\pm$ 30.8 | 25.2 $\pm$ 30.8 | 28.1 $\pm$ 31.3 |
| 1988 | 28.2 $\pm$ 31.4 | 25.4 $\pm$ 24.3 | 32.6 $\pm$ 30.1 |
| 1989 | 35.4 $\pm$ 25.9 | 41.1 $\pm$ 26.3 | 47.2 $\pm$ 30.0 |
| 1990 | 26.7 $\pm$ 25.5 | 32.9 $\pm$ 31.1 | 34.4 $\pm$ 31.1 |
| 1991 | 34.2 $\pm$ 30.7 | 38.5 $\pm$ 32.0 | 37.3 $\pm$ 34.2 |
| 1992 | 18.0 $\pm$ 22.5 | 20.4 $\pm$ 24.8 | 20.7 $\pm$ 23.0 |
| 1993 | 18.8 $\pm$ 29.5 | 22.5 $\pm$ 27.9 | 21.6 $\pm$ 26.5 |
| 2002 | 35.6 $\pm$ 28.2 | 45.8 $\pm$ 33.5 | -- <sup>1</sup> |
| 2003 | 29.7 $\pm$ 27.7 | 33.4 $\pm$ 27.1 | --              |
| 2004 | 32.9 $\pm$ 25.3 | 37.2 $\pm$ 29.8 | --              |
| 2005 | 17.8 $\pm$ 23.8 | 21.1 $\pm$ 30.9 | 17.9 $\pm$ 24.8 |
| 2006 | 18.7 $\pm$ 17.5 | 28.9 $\pm$ 24.1 | 38.6 $\pm$ 27.8 |
| 2007 | 6.6 $\pm$ 8.2   | 7.0 $\pm$ 8.6   | 14.0 $\pm$ 9.3  |
| 2008 | 13.5 $\pm$ 16.8 | 26.1 $\pm$ 28.0 | 26.9 $\pm$ 30.9 |

<sup>1</sup>No data available.




Figure 2.3: Average storm volume as a percentage of rainfall WSA, WSB, WSC (1982-1993).




Figure 2.42: Average storm volume as a percentage of rainfall WSA, WSB, WSC (2002-2008).

Table 2.17: Growing season average storm volume as a (%) of rainfall  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year | WSA             | WSB             | WSC             |
|------|-----------------|-----------------|-----------------|
| 1982 | 2.0 $\pm$ 1.0   | 2.1 $\pm$ 1.3   | 2.4 $\pm$ 2.2   |
| 1983 | 15.9 $\pm$ 20.8 | 18.2 $\pm$ 24.2 | 17.9 $\pm$ 24.0 |
| 1984 | 23.7 $\pm$ 32.7 | 36.4 $\pm$ 33.0 | 42.9 $\pm$ 35.6 |
| 1985 | 3.0 $\pm$ 4.5   | 10.5 $\pm$ 16.1 | 11.7 $\pm$ 14.3 |
| 1986 | 2.9 $\pm$ 1.9   | 5.0 $\pm$ 3.2   | 8.3 $\pm$ 3.8   |
| 1987 | 3.1 $\pm$ 1.7   | 5.3 $\pm$ 4.3   | 7.7 $\pm$ 6.8   |
| 1988 | 12.6 $\pm$ 25.5 | 6.7 $\pm$ 11.8  | 13.3 $\pm$ 20.3 |
| 1989 | 19.5 $\pm$ 18.1 | 24.5 $\pm$ 19.1 | 29.7 $\pm$ 25.3 |
| 1990 | 4.1 $\pm$ 5.6   | 4.7 $\pm$ 5.8   | 6.0 $\pm$ 6.7   |
| 1991 | 8.2 $\pm$ 11.0  | 11.7 $\pm$ 12.3 | 8.2 $\pm$ 10.9  |
| 1992 | 8.8 $\pm$ 12.0  | 9.8 $\pm$ 9.2   | 11.5 $\pm$ 9.4  |
| 1993 | 14.5 $\pm$ 27.2 | 16.8 $\pm$ 21.5 | 15.9 $\pm$ 19.4 |
| 2002 | 29.4 $\pm$ 31.2 | 36.6 $\pm$ 37.8 | -- <sup>1</sup> |
| 2003 | 8.1 $\pm$ 8.5   | 13.3 $\pm$ 14.3 | --              |
| 2004 | 20.3 $\pm$ 18.1 | 25.6 $\pm$ 26.3 | --              |
| 2005 | 19.4 $\pm$ 25.2 | 23.7 $\pm$ 32.6 | 18.3 $\pm$ 26.8 |
| 2006 | 10.6 $\pm$ 7.6  | 17.8 $\pm$ 12.4 | 28.5 $\pm$ 17.9 |
| 2007 | 2.9 $\pm$ 2.2   | 2.3 $\pm$ 2.2   | 8.8 $\pm$ 4.8   |
| 2008 | 3.2 $\pm$ 1.8   | 5.5 $\pm$ 4.3   | 7.0 $\pm$ 4.8   |

<sup>1</sup>No data available.




Figure 2.43: Growing season average storm volume as a percentage of rainfall WSA, WSB, WSC (1982-1993).




Figure 2.4: Growing season average storm volume as a percentage of rainfall WSA, WSB, WSC (2002-2008).

Table 2.18: Non-growing season average storm volume as a (%) of rainfall  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year              | WSA             | WSB             | WSC              |
|-------------------|-----------------|-----------------|------------------|
| 1982              | 46.7 $\pm$ 27.5 | 34.2 $\pm$ 17.4 | 49.6 $\pm$ 32.8  |
| 1983              | 33.8 $\pm$ 22.5 | 35.2 $\pm$ 23.6 | 32.6 $\pm$ 24.4  |
| 1984              | 45.0 $\pm$ 21.8 | 59.3 $\pm$ 14.0 | 68.2 $\pm$ 18.0  |
| 1985              | 36.2 $\pm$ 13.2 | 57.4 $\pm$ 11.2 | 61.3 $\pm$ 16.5  |
| 1986              | 38.0 $\pm$ 28.8 | 50.9 $\pm$ 27.5 | 62.1 $\pm$ 30.5  |
| 1987              | 40.1 $\pm$ 34.3 | 43.0 $\pm$ 33.6 | 46.2 $\pm$ 33.7  |
| 1988              | 41.2 $\pm$ 31.7 | 41.0 $\pm$ 20.7 | 48.7 $\pm$ 28.2  |
| 1989              | 60.2 $\pm$ 12.9 | 67.2 $\pm$ 7.1  | 74.8 $\pm$ 6.2   |
| 1990              | 49.3 $\pm$ 14.6 | 61.1 $\pm$ 15.8 | 62.9 $\pm$ 14.2  |
| 1991              | 60.3 $\pm$ 19.0 | 65.3 $\pm$ 20.6 | 66.4 $\pm$ 21.8  |
| 1992              | 54.7 $\pm$ 14.7 | 62.8 $\pm$ 22.1 | 57.5 $\pm$ 26.1  |
| 1993 <sup>1</sup> | 62.1            | 79.7            | 78.9             |
| 2002              | 42.7 $\pm$ 24.7 | 56.2 $\pm$ 26.5 | - - <sup>2</sup> |
| 2003              | 51.2 $\pm$ 22.6 | 53.6 $\pm$ 21.1 | - -              |
| 2004              | 53.2 $\pm$ 22.2 | 55.8 $\pm$ 26.1 | - -              |
| 2005              | 7.4 $\pm$ 6.6   | 4.5 $\pm$ 2.7   | 15.5 $\pm$ 3.8   |
| 2006              | 40.3 $\pm$ 19.1 | 58.4 $\pm$ 23.9 | 65.6 $\pm$ 35.0  |
| 2007              | 10.2 $\pm$ 10.5 | 11.6 $\pm$ 10.3 | 19.2 $\pm$ 10.2  |
| 2008              | 18.7 $\pm$ 18.8 | 36.4 $\pm$ 29.3 | 36.8 $\pm$ 34.2  |

<sup>1</sup>One storm suitable for analysis.

<sup>2</sup>No data available.




Figure 2.45: Non-growing season average storm volume as a percentage of rainfall WSA, WSB, WSC (1982-1993).




Figure 2.46: Non-growing season average storm volume as a percentage of rainfall WSA, WSB, WSC (2002-2008).

#### 2.3.2.4 Time to Peak

Differences in the  $T_p$  for the three watersheds were analyzed (Figures 2.47-2.49; Table 2.19). Unfortunately, the  $T_p$  was only analyzed in the second time period due to 15-minute rainfall data only becoming available in 2000. No statistical differences were found during the analyzed period. Interestingly, average annual time to peak across all subcatchments was 6.6 hours. This value may seem large due to the size of the subcatchments and the steep sideslopes but may be explained by the site-specific geologic conditions and forest hydrology. The flow path of rain before it reaches the stream may look something like the following. Rain falls through the canopy where a fraction is intercepted by leaves. The rain that makes it through the canopy then has to travel down past leaf litter and infiltrates down into a layer of organic material and soil (no overland flow occurs). After, the water will reach a tightly woven layer of interbedded sandstone, siltstone, or shale, where a fraction travels towards the stream as subsurface flow and the other fraction infiltrates further. The portion that infiltrates further then may reach a coal seam where again a fraction travels toward the stream as subsurface flow and the other fraction infiltrates. Eventually, the water that is remaining will reach a layer of bedrock where it has to travel toward the stream as subsurface flow. A final portion of the water will get tied up into soil pore spaces where it will be utilized by flora and fauna. Moving through all of these layers, especially the layers of sandstone, siltstone, shale, and coal, dampen the movement of water towards the stream and prolong the time to peak.

An observation on the effect of seasonality indicated an average longer time to peak during the non-growing season compared to the growing season during all analyzed years. Temporal rainfall distribution is expected to have played a role in the longer average time to peak during the non-growing season. Convective storms mainly occur during the warm season months and produce large intense rainfall events. From June 1990 to September 1991, 42 rainfall events were separated by temporal rainfall distribution into classifications of intense, steady, multi-interval intense, and multi-interval steady in the work by

Table 2.19: WSA, WSB, and WSC average time to peak (Tp) ± standard deviation (hr) 2002-2008.

| Year | WSA     |            |                | WSB     |            |                | WSC             |            |                |
|------|---------|------------|----------------|---------|------------|----------------|-----------------|------------|----------------|
|      | Tp Year | Tp Growing | Tp Non-Growing | Tp Year | Tp Growing | Tp Non-Growing | Tp Year         | Tp Growing | Tp Non-Growing |
| 2002 | 9.4±7.5 | 5.0±5.2    | 14.4±6.8       | 9.4±6.5 | 7.0±5.0    | 12.1±7.1       | -- <sup>1</sup> | --         | --             |
| 2003 | 6.5±6.9 | 2.9±2.6    | 10.1±8.1       | 8.6±6.5 | 4.4±3.7    | 12.9±6.0       | --              | --         | --             |
| 2004 | 7.2±8.2 | 5.0±7.4    | 10.5±8.7       | 6.5±6.4 | 4.3±4.7    | 10.1±7.5       | --              | --         | --             |
| 2005 | 4.4±6.1 | 4.1±6.6    | 6.0±0.4        | 4.7±6.6 | 4.5±7.1    | 6.1±1.6        | 5.1±6.6         | 4.6±6.8    | 8.9±4.1        |
| 2006 | 6.0±6.3 | 4.0±5.7    | 11.2±5.3       | 7.7±6.9 | 6.9±7.4    | 9.8±6.4        | 7.4±5.8         | 6.2±5.4    | 10.7±6.7       |
| 2007 | 3.0±4.9 | 1.1±0.9    | 4.9±6.5        | 5.2±6.6 | 0.7±0.4    | 9.8±6.9        | 3.6±4.7         | 0.6±0.3    | 6.5±5.2        |
| 2008 | 7.9±8.2 | 2.6±3.4    | 10.5±8.9       | 7.1±3.3 | 6.9±5.6    | 7.3±2.2        | 8.2±6.1         | 6.8±10.8   | 8.9±3.5        |

<sup>1</sup>No data available.






Figure 2.47: Average Time to Peak WSA, WSB, WSC (2002-2008).




Figure 2.48: Growing season average Time to Peak WSA, WSB, WSC (2002-2008).




Figure 2.49: Non-growing season average Time to Peak WSA, WSB, WSC (2002-2008).

Warner et al., (2010b). Of these 42 events, seven occurred during the non-growing season and only one was classified as intense. However, it should be noted that this one storm had the lowest rainfall intensity of the 14 storms classified as intense and occurred during a month that experienced 3 times as much precipitation as the long-term monthly average. As noted by Bauer (1974), infiltration rates exhibit a period of recovery when rainfall intensities are less than infiltration capacities, or during periods with absent rainfall. This would indicate that multi-interval and less intense rainfall events would allow soil infiltration rates and depressional storage to rebound and subsequently prolong the time it takes for rainfall to reach the stream, inherently increasing the time to peak.

#### 2.3.2.5 Curve Number

Results of the CN are presented in Figures 2.50-2.67 and Tables 2.20-2.25. Statistically significant differences between WSA and WSC were observed in 1984, 1985, 2006, and 2007 with average CNs ( $\lambda=0.05$ ) in WSC of 91, 79, 88, and 81, respectively; average CNs in WSA were 82, 70, 80, and 75, respectively. Statistically significant differences between WSA and WSB only occurred in 1985 with WSB having an average CN of 78 while WSA's was 70. Finally, statistical differences between WSB and WSC were observed in 2007. The average CN for WSC in 2007 was 81 while WSB's was 74. The maximum CN ( $\lambda=0.05$ ) for WSA occurred in 2004 with a value of 87 while the maximum CNs in WSB and WSC occurred directly after harvest in 1984 with values of 88 and 91, respectively. The average pre-harvest CN of WSA, WSB, and WSC were all 76 while post-harvest WSA had an average CN of 78, WSB's was 81, and WSC's was 82.

It was observed that the yearly CN in WSB became elevated directly following harvest and aligned back with WSA around 1987. WSC behaved similarly and was the most elevated from 1984-1987 but remained elevated or somewhat elevated throughout the study. As expected, CNs were the highest during the non-growing season when vegetation was dormant. With the exception of the first few years after harvest, minimal differences were observed between CNs during the non-growing season. Conversely, throughout much of

Table 2.20: Yearly curve numbers (CNs)  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year | CN ( $\lambda=0.2$ ) |             |                | CN ( $\lambda=0.05$ ) |             |             |
|------|----------------------|-------------|----------------|-----------------------|-------------|-------------|
|      | WSA                  | WSB         | WSC            | WSA                   | WSB         | WSC         |
| 1982 | 83 $\pm$ 14          | 81 $\pm$ 13 | 82 $\pm$ 15    | 77 $\pm$ 20           | 75 $\pm$ 18 | 76 $\pm$ 20 |
| 1983 | 82 $\pm$ 10          | 83 $\pm$ 11 | 83 $\pm$ 10    | 76 $\pm$ 15           | 77 $\pm$ 15 | 76 $\pm$ 15 |
| 1984 | 86 $\pm$ 10          | 91 $\pm$ 7  | 93 $\pm$ 7     | 82 $\pm$ 14           | 88 $\pm$ 10 | 91 $\pm$ 10 |
| 1985 | 78 $\pm$ 9           | 84 $\pm$ 9  | 84 $\pm$ 9     | 70 $\pm$ 13           | 78 $\pm$ 13 | 79 $\pm$ 13 |
| 1986 | 82 $\pm$ 8           | 86 $\pm$ 7  | 88 $\pm$ 7     | 76 $\pm$ 12           | 80 $\pm$ 10 | 84 $\pm$ 10 |
| 1987 | 82 $\pm$ 11          | 84 $\pm$ 11 | 86 $\pm$ 9     | 76 $\pm$ 15           | 78 $\pm$ 15 | 81 $\pm$ 13 |
| 1988 | 82 $\pm$ 14          | 82 $\pm$ 13 | 85 $\pm$ 13    | 76 $\pm$ 19           | 76 $\pm$ 18 | 80 $\pm$ 17 |
| 1989 | 85 $\pm$ 9           | 87 $\pm$ 8  | 89 $\pm$ 8     | 80 $\pm$ 12           | 82 $\pm$ 12 | 85 $\pm$ 12 |
| 1990 | 86 $\pm$ 7           | 87 $\pm$ 8  | 88 $\pm$ 7     | 81 $\pm$ 10           | 83 $\pm$ 11 | 84 $\pm$ 10 |
| 1991 | 86 $\pm$ 10          | 87 $\pm$ 10 | 86 $\pm$ 11    | 80 $\pm$ 14           | 83 $\pm$ 14 | 81 $\pm$ 15 |
| 1992 | 80 $\pm$ 9           | 82 $\pm$ 9  | 82 $\pm$ 9     | 73 $\pm$ 13           | 75 $\pm$ 13 | 76 $\pm$ 12 |
| 1993 | 77 $\pm$ 14          | 80 $\pm$ 13 | 80 $\pm$ 12    | 68 $\pm$ 19           | 73 $\pm$ 18 | 73 $\pm$ 17 |
| 2002 | 89 $\pm$ 9           | 91 $\pm$ 10 | - <sup>1</sup> | 85 $\pm$ 12           | 88 $\pm$ 14 | - -         |
| 2003 | 87 $\pm$ 10          | 88 $\pm$ 10 | - -            | 82 $\pm$ 14           | 84 $\pm$ 14 | - -         |
| 2004 | 87 $\pm$ 9           | 88 $\pm$ 10 | - -            | 83 $\pm$ 13           | 84 $\pm$ 13 | - -         |
| 2005 | 85 $\pm$ 8           | 85 $\pm$ 8  | 86 $\pm$ 8     | 80 $\pm$ 11           | 80 $\pm$ 11 | 81 $\pm$ 12 |
| 2006 | 85 $\pm$ 7           | 88 $\pm$ 7  | 91 $\pm$ 7     | 80 $\pm$ 10           | 84 $\pm$ 10 | 88 $\pm$ 9  |
| 2007 | 81 $\pm$ 8           | 81 $\pm$ 8  | 86 $\pm$ 6     | 75 $\pm$ 11           | 74 $\pm$ 12 | 81 $\pm$ 9  |
| 2008 | 83 $\pm$ 10          | 87 $\pm$ 10 | 86 $\pm$ 10    | 76 $\pm$ 14           | 82 $\pm$ 14 | 82 $\pm$ 15 |

<sup>1</sup>No data available.




Figure 2.50: Average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993).




Figure 2.51: Average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008).




Figure 2.52: Average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993).




Figure 2.53: Average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008).

Table 2.21: Growing season curve numbers (CNs)  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year | CN ( $\lambda=0.2$ ) |             |                 | CN ( $\lambda=0.05$ ) |             |             |
|------|----------------------|-------------|-----------------|-----------------------|-------------|-------------|
|      | WSA                  | WSB         | WSC             | WSA                   | WSB         | WSC         |
| 1982 | 71 $\pm$ 12          | 71 $\pm$ 12 | 71 $\pm$ 13     | 60 $\pm$ 15           | 60 $\pm$ 16 | 60 $\pm$ 17 |
| 1983 | 79 $\pm$ 10          | 80 $\pm$ 10 | 80 $\pm$ 10     | 72 $\pm$ 14           | 73 $\pm$ 15 | 72 $\pm$ 15 |
| 1984 | 82 $\pm$ 10          | 88 $\pm$ 9  | 91 $\pm$ 9      | 76 $\pm$ 14           | 84 $\pm$ 13 | 88 $\pm$ 12 |
| 1985 | 76 $\pm$ 7           | 81 $\pm$ 8  | 82 $\pm$ 8      | 66 $\pm$ 10           | 73 $\pm$ 11 | 75 $\pm$ 11 |
| 1986 | 80 $\pm$ 4           | 82 $\pm$ 3  | 84 $\pm$ 4      | 73 $\pm$ 6            | 76 $\pm$ 4  | 79 $\pm$ 6  |
| 1987 | 79 $\pm$ 6           | 81 $\pm$ 5  | 83 $\pm$ 6      | 71 $\pm$ 8            | 74 $\pm$ 7  | 76 $\pm$ 8  |
| 1988 | 75 $\pm$ 16          | 73 $\pm$ 13 | 77 $\pm$ 14     | 66 $\pm$ 21           | 63 $\pm$ 18 | 69 $\pm$ 19 |
| 1989 | 80 $\pm$ 7           | 82 $\pm$ 7  | 84 $\pm$ 8      | 72 $\pm$ 10           | 76 $\pm$ 10 | 79 $\pm$ 11 |
| 1990 | 80 $\pm$ 4           | 81 $\pm$ 3  | 82 $\pm$ 3      | 72 $\pm$ 6            | 74 $\pm$ 5  | 75 $\pm$ 5  |
| 1991 | 79 $\pm$ 6           | 81 $\pm$ 6  | 78 $\pm$ 6      | 70 $\pm$ 9            | 74 $\pm$ 9  | 70 $\pm$ 9  |
| 1992 | 77 $\pm$ 7           | 78 $\pm$ 7  | 80 $\pm$ 7      | 69 $\pm$ 10           | 70 $\pm$ 11 | 72 $\pm$ 10 |
| 1993 | 75 $\pm$ 14          | 78 $\pm$ 13 | 78 $\pm$ 11     | 66 $\pm$ 18           | 70 $\pm$ 18 | 70 $\pm$ 16 |
| 2002 | 85 $\pm$ 10          | 86 $\pm$ 11 | -- <sup>1</sup> | 79 $\pm$ 14           | 81 $\pm$ 16 | --          |
| 2003 | 79 $\pm$ 7           | 81 $\pm$ 9  | --              | 71 $\pm$ 10           | 74 $\pm$ 13 | --          |
| 2004 | 84 $\pm$ 10          | 85 $\pm$ 10 | --              | 78 $\pm$ 14           | 80 $\pm$ 14 | --          |
| 2005 | 85 $\pm$ 8           | 85 $\pm$ 9  | 85 $\pm$ 9      | 80 $\pm$ 12           | 80 $\pm$ 12 | 80 $\pm$ 12 |
| 2006 | 83 $\pm$ 6           | 86 $\pm$ 6  | 89 $\pm$ 6      | 77 $\pm$ 9            | 81 $\pm$ 9  | 85 $\pm$ 9  |
| 2007 | 79 $\pm$ 6           | 78 $\pm$ 6  | 84 $\pm$ 4      | 71 $\pm$ 9            | 70 $\pm$ 8  | 78 $\pm$ 6  |
| 2008 | 80 $\pm$ 10          | 81 $\pm$ 12 | 83 $\pm$ 11     | 73 $\pm$ 14           | 75 $\pm$ 17 | 76 $\pm$ 16 |

<sup>1</sup>No data available.




Figure 2.54: Growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993).




Figure 2.55: Growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008).






Figure 2.56: Growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993).




Figure 2.57: Growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008).

Table 2.22: Non-growing season curve numbers (CNs)  $\pm$  standard deviation for WSA, WSB, and WSC 1982-1993 and 2002-2008.

| Year              | CN ( $\lambda=0.2$ ) |             |                 | CN ( $\lambda=0.05$ ) |             |             |
|-------------------|----------------------|-------------|-----------------|-----------------------|-------------|-------------|
|                   | WSA                  | WSB         | WSC             | WSA                   | WSB         | WSC         |
| 1982              | 93 $\pm$ 5           | 90 $\pm$ 5  | 93 $\pm$ 7      | 91 $\pm$ 7            | 88 $\pm$ 7  | 90 $\pm$ 9  |
| 1983              | 89 $\pm$ 9           | 90 $\pm$ 8  | 89 $\pm$ 9      | 85 $\pm$ 13           | 86 $\pm$ 11 | 85 $\pm$ 12 |
| 1984              | 90 $\pm$ 8           | 94 $\pm$ 4  | 95 $\pm$ 4      | 87 $\pm$ 12           | 92 $\pm$ 6  | 94 $\pm$ 5  |
| 1985              | 85 $\pm$ 10          | 92 $\pm$ 5  | 93 $\pm$ 5      | 80 $\pm$ 15           | 90 $\pm$ 7  | 90 $\pm$ 7  |
| 1986              | 85 $\pm$ 11          | 90 $\pm$ 8  | 92 $\pm$ 8      | 79 $\pm$ 16           | 86 $\pm$ 11 | 90 $\pm$ 11 |
| 1987              | 85 $\pm$ 14          | 87 $\pm$ 14 | 88 $\pm$ 12     | 80 $\pm$ 19           | 82 $\pm$ 19 | 84 $\pm$ 16 |
| 1988              | 89 $\pm$ 9           | 89 $\pm$ 8  | 91 $\pm$ 8      | 85 $\pm$ 13           | 86 $\pm$ 11 | 88 $\pm$ 11 |
| 1989              | 93 $\pm$ 3           | 94 $\pm$ 2  | 96 $\pm$ 2      | 91 $\pm$ 4            | 93 $\pm$ 3  | 95 $\pm$ 2  |
| 1990              | 91 $\pm$ 4           | 94 $\pm$ 3  | 94 $\pm$ 3      | 89 $\pm$ 6            | 93 $\pm$ 5  | 93 $\pm$ 4  |
| 1991              | 93 $\pm$ 8           | 94 $\pm$ 8  | 94 $\pm$ 9      | 91 $\pm$ 11           | 92 $\pm$ 11 | 92 $\pm$ 12 |
| 1992              | 93 $\pm$ 2           | 94 $\pm$ 4  | 93 $\pm$ 5      | 91 $\pm$ 3            | 93 $\pm$ 5  | 91 $\pm$ 7  |
| 1993 <sup>1</sup> | 94                   | 97          | 97              | 92                    | 97          | 96          |
| 2002              | 94 $\pm$ 4           | 96 $\pm$ 3  | -- <sup>2</sup> | 92 $\pm$ 5            | 95 $\pm$ 4  | --          |
| 2003              | 94 $\pm$ 4           | 95 $\pm$ 4  | --              | 92 $\pm$ 6            | 93 $\pm$ 6  | --          |
| 2004              | 94 $\pm$ 4           | 94 $\pm$ 5  | --              | 92 $\pm$ 5            | 92 $\pm$ 7  | --          |
| 2005              | 86 $\pm$ 1           | 84 $\pm$ 3  | 89 $\pm$ 5      | 81 $\pm$ 1            | 78 $\pm$ 4  | 86 $\pm$ 7  |
| 2006              | 91 $\pm$ 5           | 95 $\pm$ 4  | 95 $\pm$ 6      | 89 $\pm$ 7            | 93 $\pm$ 5  | 94 $\pm$ 8  |
| 2007              | 84 $\pm$ 9           | 85 $\pm$ 9  | 88 $\pm$ 8      | 78 $\pm$ 12           | 79 $\pm$ 13 | 84 $\pm$ 12 |
| 2008              | 84 $\pm$ 11          | 89 $\pm$ 9  | 88 $\pm$ 10     | 78 $\pm$ 15           | 86 $\pm$ 12 | 84 $\pm$ 14 |

<sup>1</sup>One suitable storm for analysis.

<sup>2</sup>No data available.




Figure 2.58: Non-growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (1982-1993).

:




Figure 2.59: Non-growing season average curve number ( $\lambda=0.2$ ) WSA, WSB, WSC (2002-2008)




Figure 2.60: Non-growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (1982-1993).




Figure 2.61: Non-growing season average curve number ( $\lambda=0.05$ ) WSA, WSB, WSC (2002-2008).

Table 2.23: Average curve numbers (CNs)  $\pm$  standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.

| Period    | CN ( $\lambda=0.2$ ) |             |                 | CN ( $\lambda=0.05$ ) |             |             |
|-----------|----------------------|-------------|-----------------|-----------------------|-------------|-------------|
|           | WSA                  | WSB         | WSC             | WSA                   | WSB         | WSC         |
| 1982-1983 | 77 $\pm$ 14          | 77 $\pm$ 13 | 77 $\pm$ 14     | 69 $\pm$ 19           | 68 $\pm$ 18 | 69 $\pm$ 20 |
| 1984-1988 | 81 $\pm$ 13          | 84 $\pm$ 12 | 86 $\pm$ 11     | 74 $\pm$ 18           | 79 $\pm$ 16 | 82 $\pm$ 16 |
| 1989-1993 | 83 $\pm$ 11          | 86 $\pm$ 11 | 87 $\pm$ 11     | 78 $\pm$ 16           | 81 $\pm$ 16 | 82 $\pm$ 16 |
| 2002-2004 | 85 $\pm$ 11          | 87 $\pm$ 12 | -- <sup>1</sup> | 80 $\pm$ 15           | 82 $\pm$ 16 | --          |
| 2005-2008 | 77 $\pm$ 9           | 81 $\pm$ 10 | 83 $\pm$ 11     | 69 $\pm$ 13           | 74 $\pm$ 15 | 77 $\pm$ 15 |

<sup>1</sup>No data available.




Figure 2.62: Average curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.




Figure 2.63: Average curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.

Table 2.24: Growing season curve numbers (CNs)  $\pm$  standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.

| Period    | CN ( $\lambda=0.2$ ) |             |                 | CN ( $\lambda=0.05$ ) |             |             |
|-----------|----------------------|-------------|-----------------|-----------------------|-------------|-------------|
|           | WSA                  | WSB         | WSC             | WSA                   | WSB         | WSC         |
| 1982-1983 | 71 $\pm$ 11          | 71 $\pm$ 12 | 71 $\pm$ 12     | 60 $\pm$ 15           | 61 $\pm$ 16 | 61 $\pm$ 16 |
| 1984-1988 | 72 $\pm$ 11          | 76 $\pm$ 11 | 79 $\pm$ 12     | 62 $\pm$ 16           | 67 $\pm$ 16 | 71 $\pm$ 17 |
| 1989-1993 | 74 $\pm$ 9           | 77 $\pm$ 9  | 78 $\pm$ 9      | 64 $\pm$ 11           | 68 $\pm$ 13 | 70 $\pm$ 13 |
| 2002-2004 | 80 $\pm$ 11          | 82 $\pm$ 13 | -- <sup>1</sup> | 72 $\pm$ 15           | 75 $\pm$ 18 | --          |
| 2005-2008 | 76 $\pm$ 9           | 80 $\pm$ 11 | 82 $\pm$ 10     | 67 $\pm$ 13           | 72 $\pm$ 15 | 76 $\pm$ 15 |

<sup>1</sup>No data available.




Figure 2.64: Growing season curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.




Figure 2.65: Growing season curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.



Table 2.25: Non-growing season curve numbers (CNs)  $\pm$  standard deviation for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.

| Period    | CN ( $\lambda=0.2$ ) |             |                 | CN ( $\lambda=0.05$ ) |             |             |
|-----------|----------------------|-------------|-----------------|-----------------------|-------------|-------------|
|           | WSA                  | WSB         | WSC             | WSA                   | WSB         | WSC         |
| 1982-1983 | 91 $\pm$ 10          | 88 $\pm$ 7  | 90 $\pm$ 10     | 88 $\pm$ 14           | 84 $\pm$ 10 | 87 $\pm$ 14 |
| 1984-1988 | 86 $\pm$ 12          | 89 $\pm$ 9  | 91 $\pm$ 9      | 81 $\pm$ 16           | 85 $\pm$ 13 | 88 $\pm$ 12 |
| 1989-1993 | 92 $\pm$ 5           | 94 $\pm$ 5  | 94 $\pm$ 6      | 90 $\pm$ 8            | 92 $\pm$ 7  | 93 $\pm$ 8  |
| 2002-2004 | 93 $\pm$ 4           | 93 $\pm$ 5  | -- <sup>1</sup> | 91 $\pm$ 5            | 92 $\pm$ 7  | --          |
| 2005-2008 | 79 $\pm$ 10          | 82 $\pm$ 11 | 85 $\pm$ 12     | 71 $\pm$ 14           | 76 $\pm$ 16 | 79 $\pm$ 16 |

<sup>1</sup>No data available.




Figure 2.66: Non-growing season curve number ( $\lambda=0.2$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.





Figure 2.67: Non-growing season curve number ( $\lambda=0.05$ ) for storms with cumulative depth equal to or greater than 25.4 mm in WSA, WSB, and WSC for the periods of 1982-1983, 1984-1988, 1989-1993 and 2002-2008.

the study, large differences in CN were observed between the watersheds during the growing season.


### **2.3.3 Forest Stand Characteristics**

In 2010, species composition in WSA (Figure 2.68) was dominated by 19% each of *Liriodendron tulipifera* L. (Figure 2.69) and *Quercus rubra* (Figure 2.70) followed by *Acer rubrum* (Figure 2.71) at 14% and *Magnolia acuminata* (Figure 2.72) at 12%. Species composition in WSB (Figure 2.68) was dominated by 19% *A. rubrum*, 17% *Tilia Americana* (Figure 2.73), 14% *Q. Montana* (Figure 2.74), and 9% *L. tulipifera* L. Species composition in WSC (Figure 2.68) was dominated by 26% *L. tulipifera* L., 19% *A. rubrum*, 11%, *T. americana*, and 9% *Magnolia macrophylla* (Figure 2.75). Arthur et al. (1998) noted that before the harvest began, *Quercus* spp. accounted for 39% of canopy tree density with *Carya* spp. and *L. tulipifera* L. comprising 17% and 15%, respectively. Over a 27-year period, density of *Quercus* spp. decreased throughout all three watersheds with the largest reductions in WSB and WSC, which now contain 18% and 17% *Quercus* spp., respectively. Density of *Carya* spp. also decreased and is now at 4% in WSA and slightly over 2% each in WSB and WSC. Interestingly, *L. tulipifera* L. density increased to 19% in WSA, decreased to 9% in WSB, and increased in WSC to 26%.

WSA Species Composition



WSB Species Composition



WSC Species Composition




Figure 2.68 Percent stand composition in WSA, WSB, and WSC in 2010.



Figure 2.69: Leaf of *Liriodendron tulipifera* L.-Yellow Poplar (Mohlenbrock, 1995).



Figure 2.70: Leaf of *Quercus rubra*-Red Oak (Seiberling et al., 2006).




Figure 2.71: Leaf of *Acer rubrum*-Red Maple (Seiberling et al., 2006).



Figure 2.72: Leaf of *Magnolia accuminata*-Cucumber Tree (Chamuris, 2016).






Figure 2.73: Leaf of *Tilia americana*-Basswood (Herman, D.E., et al., 1996).



Figure 2.74: Leaf of *Q. montana*-Chestnut Oak (Mohlenbrock, 1995).



Figure 2.75: Leaves of *Magnolia macrophylla*-Bigleaf Magnolia (Profant, 2010).



The differences in tree species composition among the three watersheds could have an impact on watershed hydrology due to differences in transpiration demand and leaf area. A shift to species with lower leaf area could result in decreased canopy interception and ET and increased water yield (or vice versa). Shifts in species composition from ring-porous species (*Quercus*) to diffuse porous species (*Acer*, *Liriodendron*, and *Betula*) have been shown to effectively increase total stand transpiration (Wulschleger et al., 2001), thereby decreasing water yield. As demonstrated by Ford et al. (2011), for a given tree diameter, yellow-poplar (*Liriodendron tulipifera*) transpiration rates were nearly two-fold greater than hickory (*Carya* spp.) and four-fold greater than oaks (*Quercus* spp.). However, the mechanism by which this hydrologic alteration occurs is related to the proportional increase in diffuse porous sapwood area.

Ground cover percentages for WSA, WSB and WSC were found to average 17%, 13% and 11%, respectively. Of the total groundcover, average herbaceous cover comprised 11%, 7% and 9% for WSA, WSB and WSC, respectively. In both WSB and WSC, the exotic grass *Microstegium* sp. (Figure 2.76) was encountered. In WSB, *Microstegium* sp. was found only as large patches on logging roads whereas in WSC it was found in two separate plots away from any roads. The invasive *Lonicera maackii* (Figure 2.77) was encountered at one plot in WSB and was roughly at shoulder height and had produced numerous stems. The only other invasive plant species encountered throughout the surveying process was *Ailanthus altissima* (Figure 2.78), which was found in the upper rear reaches of WSC near an active forest road, and had produced two separate canopy sized stems.




Figure 2.76: Invasive grass *Microstegium* sp.-Nepalese browntop (Mohlenbrock, 1991).



Figure 2.77: Invasive *Lonicera maackii*-Honeysuckle (Herman, D.E., et al., 1996).



Figure 2.78: *Ailanthus altissima*-Tree of heaven (Goldman, 2012).

Stand data for WSA were found to have an average basal area of  $22.5 \text{ m}^2 \text{ ha}^{-1}$  and an average diameter of 11.2 cm in the 2010 survey. WSB was found to have an average basal area of  $22.9 \text{ m}^2 \text{ ha}^{-1}$  and average diameter of 12.6 cm. WSC was found to have an average basal area of  $21.3 \text{ m}^2 \text{ ha}^{-1}$  and average diameter of 12.1 cm. Overall, these three parameters are fairly similar among the three catchments. However, stand density does vary substantially from one catchment to the other. Stand densities in WSA, WSB and WSC were 1,144 trees  $\text{ha}^{-1}$ , 1,392 trees  $\text{ha}^{-1}$ , and 1,479 trees  $\text{ha}^{-1}$ , respectively. That is a 17.8% and 22.7% lower density in WSA than WSB and WSC, respectively.

The lower stand density in WSA may be an attribute of its age (approximately 90 years; see Cotton et al., 2013), which would be reflected by taller trees and more canopy cover. Use of LiDAR data in 2013 confirmed this relationship (Figure 2.79). Average tree height was 22.8, 18.5 and 16.5 m for WSA, WSB and WSC, respectively (Table 2.26). Moreover, the distribution of trees in the 22 to 40 m height class was much greater for WSA than WSB or WSC (Figure 2.80). The retention of trees within the SMZ also contributed to an overall increase in stand height in WSB over WSC.






Figure 2.79: Vegetation height collected by LiDAR in 2013 for WSA, WSB, and WSC.




Figure 2.80: Vegetation height distributions in 2013 in WSA, WSB, and WSC.

Table 2.26: Vegetation height distribution in WSA, WSB, and WSC.

| Vegetation attributes (m) | Watershed |       |       |
|---------------------------|-----------|-------|-------|
|                           | A         | B     | C     |
| Minimum                   | 0.00      | 0.00  | 0.00  |
| 25th percentile           | 17.37     | 15.83 | 13.95 |
| Median value              | 22.91     | 18.58 | 17.00 |
| 75th percentile           | 28.97     | 21.12 | 19.59 |
| Maximum                   | 45.74     | 44.26 | 31.80 |
| Average                   | 22.88     | 18.56 | 16.55 |

Structurally speaking, all three watersheds exhibited similar mean basal area and stem diameters in 2010, but mean stand density was lowest in WSA and highest in WSC. Figures 2.81-2.83 depict the stand density in 2015 along the stream and into the riparian areas. The greater stand density in WSB and especially in WSC over WSA may be large enough to affect competition among individual stems, which could result in greater than expected streamflow over the long-term (Ford et al., 2011). Experience has shown that changes in stand density and composition can influence the water budget. As a result of reduced ET, annual water yield has been shown to increase after forest harvesting (Hibbert 1987; Bosch 1982; Arthur et al. 1998). The magnitude and duration of the increase is related to the percentage of vegetation cover removed, climate, and forest type (Bosch 1982; Stednick 1996). Regardless, ET has been repeatedly identified as a major regulator of streamflow, soil moisture, and groundwater recharge in forested ecosystems and is mainly controlled by solar radiation, air temperature, precipitation, and vegetation characteristics including amount of leaf biomass (Sun et al. 2011). Even after nearly 30 years, mean tree heights in WSA remain higher than that observed in WSB or WSC. Given the relationship of height to biomass, it seems likely that WSA would have higher water demand and subsequently lower water yield (Pflugmacher et al., 2008).



Figure 2.81: Stand density in WSA (2015).





Figure 2.82: Stand density and regrowth in WSB (2015).



Figure 2.83: Stand density and regrowth in WSC (2015).

## 2.4 CONCLUSIONS

Harvesting has resulted in long-term impacts to the hydrologic regime of the treated areas. The implementation of BMPs in WSB has led to less severe long-term harvesting effects on streamflow than those observed in WSC. A similar observation was made by Arthur et al. (1998) who attributed these differences among treatments mainly to the retirement of haul roads and the use of a riparian buffer within WSB. Elevated stormflow and altered storm responses are the main mechanisms by which observed streamflow increases have occurred. It has been shown at Coweeta hydrologic laboratory in North Carolina that storm hydrograph response and baseflow are sensitive to logging intensity and road disturbance (Swank et al., 2001). This observation is further supported by another study performed at Coweeta, where 65% removal of stand basal area by commercial logging and tractor skidding with high road density led to roughly double the stormflow volume and peak flow rate produced by cable logging (Douglas and Swank, 1976; Swank et al., 2001).

The magnitude of streamflow increase in WSC when compared to WSA and WSB is further evidence that greater harvest intensity and greater road disturbance in logging operations leads to greater hydrologic response. In addition, changes in stand density and structure also appear to have had a significant effect on watershed hydrology, likely through increased ET that is related to the size of trees and their respective biomass. The hydrologic recovery in the treatment watersheds may depend on the long-term regrowth in WSB and WSC and the transformation of logging roads into the forest habitat. Presently, elevated stormflows at the treatment sites, especially following intense rainfall events, suggest that there may still be direct flow paths from the old logging operations to the stream. However, changes in species composition and subsequently sapwood area may alter the hydrologic function of these subcatchments and streamflow response may not reach pre-harvest conditions.

Nonetheless, there are some indications that hydrologic recovery in WSB has occurred or nearly occurred while WSC continues to show some impacts

from the harvest. The use of BMPs in WSB, most notably the retirement of haul roads and use of a riparian buffer, has effectively decreased the severity of harvesting impacts to overall hydrology. Although water yield within WSB has attenuated to a level that is no longer significantly different than the control, complete hydrologic recovery under all rainfall intensities in either watershed has yet to be observed. The true mechanisms by which such long-term hydrologic response has occurred remain unclear, although several hypotheses exist. 1) The intensity of cutting coupled with the design and placement of haul roads has significantly increased stormflow as well as hydrologic sensitivity to high precipitation events. These effects are greatest in WSC, which experienced a greater intensity of treatment. 2) Shifts in belowground flow and storage processes have resulted in significantly greater hydrologic input to the stream channels. These treatment effects have been greatest in WCS. 3) Increased stand density in treatment watersheds has been sufficient to affect an increase in competition, thereby reducing leaf area index and stand transpiration. This reduction in stand transpiration has caused a significant switch in streamflow from baseflow to stormflow throughout the growing season in the treatment watersheds. These three hypotheses are not mutually exclusive, and it is certainly possible that each are contributing to some extent to the observations made within the treatment watersheds B and C.



## **CHAPTER 3: LONG-TERM EFFECTS OF BEST MANAGEMENT PRACTICES ON WATER CHEMISTRY FOR THREE APPALACHIAN HEADWATER CATCHMENTS**

### **3.1 INTRODUCTION**

Headwater streams are generally 1<sup>st</sup>-3<sup>rd</sup> order ephemeral, intermittent, and perennial reaches (Vannote et al., 1980) and relatively small in stature; however, cumulatively they contribute 60 to 85% of the total stream length in a river network, and drain 70 to 90% of the total drainage basin area (Leopold et al., 1964; Benda et al., 2005; MacDonald and Coe, 2007). Various definitions exist for what constitutes ephemeral, intermittent, and perennial reaches but the following has been chosen for this study. Ephemeral reaches are defined as streams that only flow in direct response to precipitation. Intermittent reaches are defined as streams that only flow for a portion of the year, usually the wet season, and when the water table is above the streambed. Perennial reaches are the streams that flow continuously under normal precipitation conditions (USEPA, 2013). These headwater stream systems and their riparian zones provide valuable habitat to sensitive flora and fauna as well as macroinvertebrate populations (Meyer et al., 2007; Richardson and Danehy, 2007) and serve as significant contributors to down-gradient stream and river health by filtering upland derived sediment, processing and cycling nutrients, and providing energy inputs for larger downstream organisms (Pritchett and Fisher, 1987; Kaplan et al., 2008).

Average annual concentrations for common constituents in forested systems are; NO<sub>3</sub> (0.01-1.7 mg L<sup>-1</sup>), total suspended solids are generally less than 10 mg L<sup>-1</sup> but can range from 100-1000 mg L<sup>-1</sup> during stormflows, and PO<sub>4</sub> (μ=.08 mg L<sup>-1</sup>) (Binkley and Brown, 1993; Ice and Binkley, 2003). Water quality of forested systems is usually considered superior compared to other land uses but factors such as: season, geology and soil chemistry, past land use, severity of erosion, air pollution inputs, streamflow levels and sources, and types and age of

vegetation present can significantly alter stream quality (Stuart and Edwards, 2006). Forest harvesting is another factor that has been shown to significantly impact stream water quality as well. Studies done in the northeastern United States on stream water quality following harvest have found that harvesting a forested watershed leads to an increase in  $\text{NO}_3$  concentration as well as an increase in base cations (Mg, Ca, K, Na), aluminum, and acidity (Martin et al., 2000; Wang et al., 2005).

Nitrogen is a key nutrient for sustaining plant health and is often the limiting nutrient in forested systems. However, removal of vegetation from a site reduces the amount of nitrogen taken up by plants and can lead to excess nitrogen leaching through the soil and into the stream. Multiple studies have confirmed this and shown that forest harvesting resulted in a change in soil processes such as N-mineralization and nitrification which then lead to an excess of nutrients that leached through the soil and into the streams (Hornbeck and Leak, 1992; Pierce et al., 1993; Burns and Murdoch, 2005). The primary inputs of nitrogen into forested systems are via long-term inputs of small amounts of nitrogen in precipitation, particulates, dry deposition, and nitrogen fixation (Binkley et al., 2000). The nitrogen cycle consists of mineralization of soluble or insoluble organic nitrogen to  $\text{NH}_4$ , followed by either immobilization via microbial uptake or nitrification to  $\text{NO}_3$ . Nitrate may be leached from the soil, immobilized by microbial uptake, taken up by plants, or lost via denitrification (Binkley and Fisher, 2012).

Phosphorous is another key nutrient vital for plant growth. The primary pathway that phosphorous enters into forested systems is through weathering of minerals. Phosphate ions bind tightly to soil molecules and are highly insoluble, this means very little leaches out with water runoff. From there, phosphate can either be taken up by vegetation or erode into the stream via sediment transport. The phosphorus cycle is unique in that it does not have a gas phase. Therefore, the amount of phosphate at a site is dictated by the rate of mineral weathering, the amount of phosphate in soil suspension, and the amount of phosphate in biological pools (Binkley and Fisher, 2012). Forest harvesting removes

phosphate from the available phosphate in biological pools but, surprisingly, does not usually result in phosphate increases in the stream (Aubertin and Patric, 1974; Stuart and Dunshie, 1976; Hornbeck et al., 1987; Richardson, 1988; Jewett et al., 1995; Clark et al., 2000). One paired watershed study, similar in design to this research (control watershed, BMP watershed, no BMP watershed), done in the coastal plain physiographic province in Virginia did record significant increases in phosphate concentration in the stream for the no BMP watershed following clear-cutting (Wynn et al., 2000).

Another factor that can impact stream quality in forested systems is the effects from atmospheric deposition. This is especially important in Kentucky where a large fraction of energy production is produced from coal-fired power plants. Kentucky is home to 21 coal-fired power plants with 45 more plants in the surrounding states of Indiana, Ohio, Tennessee, Virginia, and West Virginia. As of 2012 Kentucky ranked 6<sup>th</sup> for the top states that rely on coal-fired power production (USEIA, 2012). Coal-fired power plants produce large amounts of SO<sub>2</sub> and NO<sub>x</sub> that enter into the atmosphere and subsequently fall back as acidic precipitation otherwise known as acid rain. Several experiments that were conducted on small forested headwater catchments found that sites that suffer from soil acidification due to atmospheric deposition experience large changes in stream water alkalinity or stream buffering capacity from small changes in ionic inputs that leach through the soil column (Lange et al., 1996).

The Clean Air Act was enacted in 1970 by the United States Environmental Protection Agency (USEPA) to combat air pollution. Later in 1990 an amendment was made to the Clean Air Act to specifically address the issue of acid rain by limiting the amount of SO<sub>2</sub> and NO<sub>x</sub> that entered into the atmosphere. From 1984 to 2013, decreases of 62% and 34% in the wet deposition of SO<sub>4</sub> and NO<sub>3</sub>, respectively, were recorded at the Lilley Cornett Woods National Atmospheric Deposition Program (NAPD) monitoring site in Letcher County, Kentucky. Studies have shown that a reduction in acid deposition leads to an increase in alkalinity and subsequently increases a stream's buffering capacity (Chen and Lin 2009; Neal et al., 2010;). Although reductions in SO<sub>2</sub> and NO<sub>x</sub>

emissions have been recorded, recent studies conducted at the Daniel Boone National Forest have shown that the effects from atmospheric deposition on forest soils are still prevalent and noted that inputs from  $\text{SO}_2$  and  $\text{NO}_x$  can lead to a loss of base cations that leach into the stream, increase soil acidity, mobilize Al on exchange complexes, increase the weathering of clay minerals, and leach heavy metals and pollutants into solution (Reuss, 1983; Barton et al., 2002; Sanderson, 2014). While coal-fired power plant emissions have been reduced through emission controlling technology, such as scrubbers, production increases in other sectors, such as aerosols and livestock, have increased the amount of  $\text{NH}_4$  in precipitation. A study that was conducted close to the study site in Rowan county Kentucky from 1990-1998 found that the concentration of  $\text{NH}_4$  in precipitation had significantly increased from an initial concentration of  $0.2 \text{ mg L}^{-1}$  in 1990 to a value of  $0.35 \text{ mg L}^{-1}$  by 1998 (Aneja, 2003).

Best management practices (BMPs) are employed all over the United States to reduce soil erosion and subsequent suspended sediment from land use changes. The problems that arise from soil erosion are numerous and are not centralized to the location they originate. Once vegetation is removed from a site, the soil becomes more vulnerable to transport (Aust et al., 2011). Storm events collect this soil and transport it into nearby streams and rivers; this can decrease water quality, negatively impact aquatic biota, reshape streams and rivers, and lower soil fertility (Stuart and Edwards, 2006; Frady et al., 2007; Boggs et al., 2016). Headwater catchments such as the ones being studied here are especially vulnerable to erosion because of their steep gradient and the fact that they serve as sediment traps until large precipitation events periodically flush this sediment downstream (Benda et al., 2005). From visual observations and from Arthur et al., (1998), it would seem that harvesting exacerbates this phenomenon and sediment transport events occur more readily and are in general larger. The focus of this research is to assess the nutrient transport that takes place post-harvest due to erosional processes as well as inputs from atmospheric deposition by comparing yearly, growing, and non-growing season concentrations of  $\text{SO}_4$ , Mg, Ca, K, Na, total alkalinity,  $\text{NO}_3$ ,  $\text{PO}_4$ , Cl,  $\text{NH}_4$ , and total organic carbon to



determine significant pairwise differences and long-term trends for the monitored time period. Due to the long-term nature of this study, it is hypothesized that some of the monitored constituents that enter through atmospheric deposition may gradually decline due to more stringent air quality regulations imposed by the 1990 amendments made to the Clean Air Act.

## 3.2 METHODS

### 3.2.1 Water Quality Data

Grab samples for the three watersheds were collected on a weekly basis starting in February 1982 and have continued until present. Constituents and their analyzation method from 1982-1993 were:  $\text{SO}_4$  (Bausch-Lomb mass spectrophotometer); Mg, Ca, K, Na (Perkin Elmer Atomic Absorption Spectrophotometer); total alkalinity (titration to methyl orange endpoint);  $\text{NO}_3$  (nitrate reductase method; 1982-1990); and  $\text{PO}_4$  (Bausch-Lomb mass spectrophotometer). For 2002-2008 samples,  $\text{SO}_4$  and Cl concentrations were determined using a quantitative ion chromatography procedure on a Dionex Ion Chromatograph (IC) 2000. Measurement of Mg, Ca, K, and Na concentrations were made with a GBC SDS 270 Atomic Adsorption Spectrophotometer. Total alkalinity was determined using an Orion pH meter and auto titrator with a titrant endpoint pH of 4.6. Analysis of  $\text{NO}_3$  and  $\text{NH}_4$  were performed using a Bran-Luebbe Autoanalyzer (continuous-flow multi-test methods; MT7/MT8 (EPA 353.2) and MT15/16 (EPA 350.1), respectively). Total organic carbon was measured on samples of  $\leq 2\text{mL}$  with a Shimadzu TOC 5000A Analyzer (Shimadzu Corporation).

The constituents were summarized into yearly, growing, and non-growing season flow-weighted mean concentrations (FWMC) (Equation 3.1).

$$\bar{c} \cong \frac{\sum_{i=1}^n \left( \frac{c_i + c_{i+1}}{2} \right) \left( \frac{Q_i + Q_{i+1}}{2} \right)}{\sum_{i=1}^n \left( \frac{Q_i + Q_{i+1}}{2} \right)} \quad \text{Equation 3.1}$$

In the equation above,  $\bar{c}$  represents the FWMC,  $c_i$  is the concentration at time  $i$ , and  $Q_i$  is the discharge at time  $i$ . This method is advantageous because it summarizes all of the weekly concentrations for a specified time period into a single concentration, thus making comparison easier. It also takes into account the effects from variable flow. According to Rickert (1985), constituent concentrations from metals, nutrients, and suspended sediments increase with increasing flow. The FWMC achieves this by placing more weight on constituent concentrations that occur with higher flows and less weight on concentrations that occur under lower flows. The one negative to using the FWMC is that it is flow dependent, meaning flow data are required for analysis. Suspended sediment samples were also collected and analyzed using a filtration technique (1982-1986 and 1988-1990). Due to extended equipment failure (ISCO Model 1680), suspended sediment data were limited. The data available were analyzed by a previous study (Arthur et al., 1998), and because of this, no further analysis was conducted as part of this study.

### **3.2.2 Statistical Analysis**

Significant temporal differences in water quality constituents due to treatment were analyzed by a before-after-control-impact (BACI) statistical design. Data were categorized into before (pre-harvest) and after (post-harvest) and then a one-way analysis of variance (ANOVA) was conducted to test for significant ( $p \leq 0.5$ ) time and treatment interactions. If significant differences were found in treatment only, then that would suggest that the watersheds were weakly paired for that constituent. If significant differences were found for time only, then that would suggest there is an independent background time effect not related to treatment. Finally, if significant differences were found for treatment\*time, then that would suggest that treatment has an effect compared to the control and that the before and after trends vary among treatments. In addition to the above analyses, a linear regression was conducted in order to determine the long-term trends in the data and to assess whether the trends

were significantly different from zero. Statistical analyses were performed in SAS version 9.4 (SAS Institute, Inc., 2015).

### **3.3 RESULTS AND DISCUSSION**

#### **3.3.1 Time and Treatment Effects**

The results regarding sediment loading found in the Arthur et al. (1998) study concluded that WSB and WSC were significantly higher than WSA from 1984-1986, WSC was significantly higher than WSA in 1988, and in 1990 all three watersheds were significantly different from each other with WSC being the highest and WSA being the lowest. Another interesting result from the study was that:

*Ninety-five percent of the sediment flux from Watershed B in that year (1987) occurred during a single early November 7.5-cm rain event that produced 733 kg/ha of suspended sediment. It is likely that Watershed C had a similarly large sediment discharge during that single storm because the two clear-cut watersheds generally paralleled each other in sediment production during this period. Watershed A produced only 9.7 kg/ha suspended sediment during the same rainfall event. -Arthur et al. (1998).*

Steep gradient headwater catchments, such as the ones being studied here, are susceptible to mass wasting and serve as sediment traps, large precipitation events periodically flush this sediment downstream (Benda et al., 2005). From visual observations and from Arthur et al. (1998), it would seem that harvesting exacerbates this phenomenon and sediment transport events occur more readily and are in general larger (Arthur et al., 1998). A more recent study (2001) on coarse woody debris (CWD) was conducted on these watersheds and found that the watersheds differed in the type of CWD and in the recruitment of CWD. Based on their findings, the authors suggested that WSB may be more prone to windthrow or slumping, which in turn would affect its capacity to buffer sediment

from reaching the stream on the long-term (McClure et al., 2004). Although sediment data were no longer collected after 1990, it seems that sediment buildup is still presently an issue. Figures 3.1-3.3 depict the sediment buildup that was present behind the flumes during a site visit in July 2016. During the visit, it was noted that considerable amounts of sediment were trapped behind the flumes in WSB and WSC even though sediment is removed every 2-3 months and as needed after large storm events.



Figure 3.1: Sediment and debris buildup behind WSA flume (2016).



Figure 3.2: Sediment and debris buildup behind WSB flume (2016).





Figure 3.3: Sediment and debris buildup behind WSC flume (2016).

Results from the statistical analysis on the significant time interactions on constituent concentrations across the watersheds are presented in Table 3.1. From the analysis, it was found that  $\text{SO}_4$ , Ca (WSB), K (WSC), Na (WSB), ALK,  $\text{NO}_3$ , and  $\text{PO}_4$  were statistically different from pre-harvest to post-harvest. Results also showed statistically similar results from pre-harvest to post-harvest for Ca (WSA, WSC), K (WSA, WSB), and Na (WSA, WSC) while Mg was deemed not statistically significant for any of the watersheds. Constituents whose monitoring period began in 2002 (Cl,  $\text{NH}_4$ , TOC) were not analyzed in this section because no pre-harvest data were available.



Table 3.1: ANOVA time interaction results, pre- and post-harvest concentration ( $\text{mg L}^{-1}$ ) plus sample size (pre)x(post).<sup>1</sup>

| Constituent     | WSA         |              |             | WSB         |              |             | WSC         |              |             |
|-----------------|-------------|--------------|-------------|-------------|--------------|-------------|-------------|--------------|-------------|
|                 | Pre-harvest | Post-harvest | Sample Size | Pre-harvest | Post-harvest | Sample Size | Pre-harvest | Post-harvest | Sample Size |
| SO <sub>4</sub> | 12.7a       | 9.3b         | (75)x(791)  | 13.4a       | 9.9b         | (72)x(738)  | 11.4a       | 8.6b         | (68)x(769)  |
| Mg              | 1.8         | 1.7          | (75)x(791)  | 2.0         | 1.9          | (72)x(774)  | 1.7         | 2.1          | (68)x(770)  |
| Ca              | 2.2a        | 2.3a         | (75)x(791)  | 2.6a        | 2.3b         | (72)x(774)  | 2.5a        | 2.5a         | (68)x(770)  |
| K               | 1.3a        | 1.4a         | (75)x(791)  | 1.6a        | 1.6a         | (72)x(774)  | 1.5b        | 1.8a         | (68)x(770)  |
| Na              | 1.0a        | 1.0a         | (75)x(791)  | 1.2a        | 1.1b         | (72)x(774)  | 0.9a        | 1.0a         | (68)x(770)  |
| ALK             | 12.2b       | 23.5a        | (75)x(791)  | 12.9b       | 23a          | (69)x(772)  | 13.4b       | 27.9a        | (66)x(770)  |
| NO <sub>3</sub> | 0.13b       | 0.38a        | (75)x(778)  | 0.17b       | 0.49a        | (72)x(760)  | 0.1b        | 0.5a         | (68)x(770)  |
| PO <sub>4</sub> | 0.07b       | 0.1a         | (75)x(484)  | 0.06b       | 0.1a         | (72)x(472)  | 0.04b       | 0.1a         | (68)x(465)  |

Constituents with different letters (a,b) from pre to post-harvest indicate statistical differences within watershed between time periods, while constituents with the same letter denote statistical similarity between time periods

Constituents with no letters indicate no statistical significance

Significant statistical differences between treatments were observed for  $\text{SO}_4$ , Ca, K, Na, and TOC. Slightly less significant differences ( $p \leq 0.10$ ) were also observed for  $\text{NO}_3$  and Cl. Magnesium, ALK,  $\text{PO}_4$ , and  $\text{NH}_4$  exhibited no statistical significance during the study period. Results from the analysis showed all watersheds were statistically different for  $\text{SO}_4$  with WSB having the highest concentration and WSC having the lowest. WSB and WSC had nearly identical Ca concentrations and were statistically higher than WSA for the study period. Next, all three watersheds had statistically different concentrations for K with WSC being the highest and WSA being the lowest. Sodium was the next constituent to be analyzed with WSB having a statistically higher concentration than both WSA and WSC. Furthermore, results from  $\text{NO}_3$  showed WSB having the highest concentration and WSA having the lowest with WSC having a statistically similar concentration to both WSA and WSB. Next, Cl was analyzed and the results showed WSA and WSB had a statistically higher concentration than WSC. Finally, results from TOC displayed statistical similarity for WSA and WSC and they had a statistically higher concentration than WSB.

Table 3.2: ANOVA treatment interaction results, concentration (mg L<sup>-1</sup>) plus sample.<sup>1</sup>

| Constituent                  | WSA           |             | WSB           |             | WSC           |             |
|------------------------------|---------------|-------------|---------------|-------------|---------------|-------------|
|                              | Concentration | Sample Size | Concentration | Sample Size | Concentration | Sample Size |
| SO <sub>4</sub>              | 11.0b         | 877         | 11.6a         | 821         | 10.0c         | 844         |
| Mg                           | 1.8           | 877         | 1.9           | 857         | 1.9           | 846         |
| Ca                           | 2.2b          | 877         | 2.5a          | 857         | 2.5a          | 846         |
| K                            | 1.38c         | 877         | 1.57b         | 857         | 1.63a         | 846         |
| Na                           | 0.97b         | 877         | 1.17a         | 857         | 0.97b         | 846         |
| ALK                          | 17.8          | 877         | 18.0          | 852         | 20.6          | 844         |
| NO <sub>3</sub> <sup>*</sup> | 0.26b         | 864         | 0.33a         | 843         | 0.28ab        | 836         |
| PO <sub>4</sub>              | 0.08          | 570         | 0.08          | 555         | 0.08          | 541         |
| Cl <sup>*</sup>              | 0.82a         | 307         | 0.83a         | 301         | 0.78b         | 305         |
| NH <sub>4</sub>              | 0.04          | 184         | 0.0           | 173         | 0.1           | 184         |
| TOC                          | 4.5a          | 275         | 4.0b          | 268         | 4.6a          | 269         |

\* Significance level of  $\alpha \leq 0.10$

Constituents with different letters (a,b,c) indicate statistical differences between watersheds, while constituents with the same letter denote statistical similarity between watersheds

Constituents with no letters indicate no statistical significance

Significant statistical differences were observed for  $\text{SO}_4$ , Ca, K, and Na while no statistical significance differences were found for Mg, ALK,  $\text{NO}_3$ , and  $\text{PO}_4$ ; this was true for both pre- and post-harvest. During pre-harvest, all three watersheds were statistically different for  $\text{SO}_4$  (WSB, WSA, WSC), Ca (WSB, WSC, WSA), and K (WSB, WSC, WSA); watersheds are listed in order of highest to lowest concentration. Watershed B had a statistically higher concentration for Na pre-harvest while the concentrations in WSA and WSC were lower and statistically similar. Post-harvest statistical differences were exhibited across all three watersheds for  $\text{SO}_4$  (WSB, WSA, WSC) and Ca (WSC, WSB, WSA). Watershed B and WSC had a statistically similar and higher concentration for K post-harvest compared to WSA. Similar to the pre-harvest result, WSB had a statistically higher concentration for Na while the concentration in WSA and WSC was lower and statistically similar.

Table 3.3: ANOVA time\*treatment interaction results, pre and post-harvest concentration (mg L<sup>-1</sup>) plus sample size (pre)x(post).<sup>1</sup>

| Constituent     | Pre-harvest |       |       | Post-harvest |      |      | Sample Size |            |            |
|-----------------|-------------|-------|-------|--------------|------|------|-------------|------------|------------|
|                 | WSA         | WSB   | WSC   | WSA          | WSB  | WSC  | WSA         | WSB        | WSC        |
| SO <sub>4</sub> | 12.8b       | 13.4a | 11.4c | 9.3b         | 9.9a | 8.6c | (75)x(791)  | (72)x(738) | (68)x(769) |
| Mg              | 1.8         | 2.0   | 1.7   | 1.7          | 1.9  | 2.1  | (75)x(791)  | (72)x(774) | (68)x(770) |
| Ca              | 2.2c        | 2.6a  | 2.5b  | 2.3c         | 2.6a | 2.5b | (75)x(791)  | (72)x(774) | (68)x(770) |
| K               | 1.3c        | 1.6a  | 1.5b  | 1.4b         | 1.6a | 1.8a | (75)x(791)  | (72)x(774) | (68)x(770) |
| Na              | 1.0b        | 1.2a  | 1.0b  | 1.0b         | 1.1a | 1.0b | (75)x(791)  | (72)x(774) | (68)x(770) |
| ALK             | 12.2        | 12.9  | 13.4  | 23.5         | 23.0 | 27.9 | (75)x(791)  | (69)x(772) | (66)x(770) |
| NO <sub>3</sub> | 0.13        | 0.17  | 0.1   | 0.39         | 0.49 | 0.47 | (75)x(778)  | (72)x(760) | (68)x(770) |
| PO <sub>4</sub> | 0.07        | 0.06  | 0.04  | 0.10         | 0.10 | 0.11 | (75)x(484)  | (72)x(472) | (68)x(465) |

Constituents with different letters (a,b,c) indicate statistical differences within time period and between the watersheds, while constituents with the same letter denote statistical similarity between watersheds within the time period

Constituents with no letters indicate no statistical significance

The final set of results to be analyzed was the long-term regression analysis that was run on the constituent concentrations (Table 3.4). The goal of the regression analysis was to determine if there were any long-term trends in the data and to assess whether constituent concentrations had increased or decreased throughout the study period and whether that change was significantly different from zero. Constituents that showed a statistically significant long-term decrease in concentration were  $\text{SO}_4$ , Mg (WSA, WSB), Ca, K (WSB, WSC),  $\text{NO}_3$ ,  $\text{PO}_4$ , and TOC. Constituents that showed a statistically significant long-term increase in concentration were Na, ALK (WSA), Cl (WSA, WSC), and  $\text{NH}_4$ .

Table 3.4: Water quality regression analysis, slope ( $\text{mg L}^{-1} \text{ d}^{-1} \times 10^{-4}$ ),  $p$ -value, and sample size.<sup>1</sup>

| Constituent     | Slope |       |       | p-value |      |       | Sample Size |     |     |
|-----------------|-------|-------|-------|---------|------|-------|-------------|-----|-----|
|                 | WSA   | WSB   | WSC   | WSA     | WSB  | WSC   | WSA         | WSB | WSC |
| SO <sub>4</sub> | -2.0  | -2.2  | -1.7  | *       | *    | *     | 877         | 821 | 844 |
| Mg              | -0.28 | -0.48 | -0.06 | *       | *    | 0.9   | 877         | 857 | 846 |
| Ca              | -1.1  | -1.4  | -1.8  | *       | *    | *     | 877         | 857 | 846 |
| K               | 0.03  | -0.20 | -0.24 | 0.6     | *    | 0.006 | 877         | 857 | 846 |
| Na              | 0.27  | 0.09  | 0.10  | *       | 0.02 | *     | 877         | 857 | 846 |
| ALK             | 3.4   | 0.8   | 7.5   | 0.02    | 0.6  | 0.2   | 877         | 852 | 844 |
| NO <sub>3</sub> | -0.10 | -0.30 | -0.40 | 0.05    | *    | *     | 864         | 843 | 836 |
| PO <sub>4</sub> | -0.07 | -0.07 | -0.10 | 0.02    | 0.06 | 0.005 | 570         | 555 | 541 |
| Cl              | 0.66  | 0.18  | 1.1   | 0.05    | 0.7  | 0.002 | 307         | 301 | 305 |
| NH <sub>4</sub> | 0.37  | 0.18  | 0.25  | *       | *    | *     | 184         | 173 | 184 |
| TOC             | -4.6  | -4.8  | -7.3  | 0.06    | 0.02 | 0.003 | 275         | 268 | 269 |

\*  $p < 0.001$

Results from the statistical analyses showed some very interesting time and treatment effects. One of the main aspects of this analysis was to assess the combined contribution of time and treatments interactions (Table 3.3) on constituent concentrations during the study period. If the statistical relationship varied from pre-harvest to post-harvest, then we could confidently say that treatment had an effect on constituent concentration. This was the case for the base cations Ca and K which are shown in Figures 3.4-3.6 and 3.7-3.9, respectively. For Ca, all three watersheds were significantly different from each other pre-harvest with WSB having the highest concentration and WSA having the lowest. Post-harvest, all three watersheds remained significantly different but WSC had the highest Ca concentration while WSA remained the lowest. When looking at the time only interaction for Ca, it can be seen that the concentration for Ca remained statistically similar in WSA and WSC throughout the study while following treatment, Ca concentrations in WSB declined. For K, all three watersheds were significantly different pre-harvest with WSB having the highest concentration and WSA having the lowest. Post-harvest, WSB and WSC were statistically similar and also had a statistically higher concentration compared to WSA. When looking at the time interaction as well, it can be seen that the K concentration in WSA and WSB remained statistically similar from pre-harvest to post-harvest while the K concentration in WSC was statistically higher following harvest. These results indicate that the BMPs in WSB helped retain the base cations of Ca and K compared with WSC.






Figure 3.4: Yearly calcium FVMC (1982-1993) and (2002-2008).




Figure 3.5: Growing season calcium FVMC (1982-1993) and (2002-2008).




Figure 3.6: Non-growing season calcium FVMC (1982-1993) and (2002-2008).




Figure 3.7: Yearly potassium FPMC (1982-1993) and (2002-2008).




Figure 3.8: Growing season potassium FPMC (1982-1993) and (2002-2008).




Figure 3.9: Non-growing season potassium FPMC (1982-1993) and (2002-2008).

The base cations that did not have significant time\*treatment interactions (Mg, Na) exhibited some mixed results in this study. Magnesium concentration was deemed not statistically significant for all three interactions (time, treatment, and time\*treatment), and from the regression analysis, WSA and WSB showed a statistically significant decline in Mg concentration over the study period (Figures 3.10-3.12). Although it was deemed not statistically significant, spikes in Mg concentration can be seen graphically in WSB and WSC post-harvest. As for Na, a significant time interaction was observed in WSB where Na concentration was statistically higher pre-harvest compared to post-harvest. However, results from the regression analysis showed a statistically significant increase in Na concentration for all three watersheds throughout the study period (Figures 3.13-3.15). These results may be explained by some recent studies that were conducted at the Daniel Boone National Forest. The studies examined the effects of atmospheric deposition on forest soils and noted that inputs from  $\text{SO}_4$  and  $\text{NO}_x$  can lead to a loss of base cations that leach into the stream (Barton et al., 2002; Reuss, 1983; Sanderson, 2014). Interestingly, large spikes in Mg, Ca, and Na concentrations were observed around 1988, five years post-harvest. These spikes may have been caused by deposition from dust produced by surface mining operations that took place next to Clemons Fork around that time (Figure 2.3).




Figure 3.10: Yearly magnesium FVMC (1982-1993) and (2002-2008).




Figure 3.11: Growing season magnesium FVMC (1982-1993) and (2002-2008).




Figure 3.12: Non-growing season magnesium FVMC (1982-1993) and (2002-2008).




Figure 3.13: Yearly sodium FWMC (1982-1993) and (2002-2008).




Figure 3.14: Growing season sodium FWMC (1982-1993) and (2002-2008).




Figure 3.15: Non-growing season sodium FWMC (1982-1993) and (2002-2008).

Some other interesting results were seen for those constituents that enter through atmospheric deposition. Looking at the time interaction, the  $\text{SO}_4$  concentration in all three watersheds was significantly reduced following harvest (Figures 3.16-3.18). This was also seen in the results from the regression analysis where the concentration of  $\text{SO}_4$  had significantly declined throughout the study period across all watersheds. This outcome is most likely the result of more stringent air quality regulations imposed by a 1990 amendment to the Clean Air Act, which affected many nearby coal-fired power plants (Sanderson, 2014). Nitrate, another constituent affected by the Clean Air Act's regulations also saw an overall significant decrease in concentration over the study period across all watersheds (Figures 3.19-3.21). However, it should be noted that  $\text{NO}_3$  concentration spiked for both harvested watersheds directly after treatment and remained elevated for roughly five years. This effect is largely a result of a cumulative reduction in plant  $\text{NO}_3$  uptake due to removed vegetation. This resulted in a change in soil processes such as N-mineralization and nitrification which then lead to an excess of nutrients that leached through the soil and into the streams (Hornbeck and Leak, 1992; Pierce et al., 1993; Burns and Murdoch, 2005). A statistically significant increase in  $\text{NH}_4$  concentration was also exhibited from 2002-2008 across all watersheds (Figures 3.22-3.24). This result is interesting because  $\text{NH}_4$  is usually pretty immobile in soil because it binds to cation exchange sites. Also,  $\text{NH}_4$  that is in soil solution usually gets converted to  $\text{NO}_3$  through nitrification;  $\text{NO}_3$  does not bind to cation exchange sites and is therefore easily leachable. This would indicate that the increase in  $\text{NH}_4$  concentration is likely an input from precipitation. A study conducted in nearby Rowan county Kentucky from 1990-1998 confirmed that the concentration of  $\text{NH}_4$  had significantly increased in precipitation and nearly doubled during the study period (Aneja, 2003).




Figure 3.16: Yearly sulfate FVMC (1982-1993) and (2002-2008).




Figure 3.17: Growing season sulfate FVMC (1982-1993) and (2002-2008).




Figure 3.18: Non-growing season sulfate FVMC (1982-1993) and (2002-2008).




Figure 3.19: Yearly nitrate FPMC (1982-1990) and (2002-2008).




Figure 3.20: Growing season nitrate FPMC (1982-1990) and (2002-2008).




Figure 3.21: Non-growing season nitrate FPMC (1982-1990) and (2002-2008).






Figure 3.22: Yearly ammonium FPMC (2002-2008).




Figure 3.23: Growing season ammonium FPMC (2002-2008).




Figure 3.24: Non-growing season ammonium FPMC (2002-2008).

Total alkalinity was another constituent that was affected by soil acidity and subsequently affected by changes to the Clean Air Act in 1990. Several experiments that were conducted on small forested headwater catchments found that sites that suffer from soil acidification due to atmospheric deposition experience large changes in stream water alkalinity from small changes in ionic inputs that leach through the soil column (Lange et al., 1996). Changes such as this were exhibited in this study and can be seen from the figures for  $\text{SO}_4$  (Figures 3.16-3.18) and (Figures 3.25-3.27). ALK where changes in  $\text{SO}_4$  concentration had large inverse effects on ALK concentration. During the second time period, following the 1990 amendment to the Clean Air Act,  $\text{SO}_4$  concentration significantly declined and was much less variable year to year. This correlated into less variability for ALK as well and results from the time interaction showed a statistical increase in ALK concentration throughout the study period.




Figure 3.25: Yearly alkalinity FWMC (1982-1993) and (2002-2008).




Figure 3.26: Growing season alkalinity FWMC (1982-1993) and (2002-2008).




Figure 3.27: Non-growing season alkalinity FWMC (1982-1993) and (2002-2008).

Results from the time interaction exhibited a statistical increase in  $\text{PO}_4$  concentrations across all watersheds from pre-harvest to post-harvest (Figures 3.28-3.30). Increases in  $\text{PO}_4$  concentrations in WSB and WSC were observed for two years post-harvest and then realign with WSA in 1986. Afterwards, all three watersheds experience a large increase in phosphate concentration around 1988. This increase in  $\text{PO}_4$  may be linked to the surface mining operations that took place next to Clemons Fork, as increases were seen around the same time as Mg, Ca, and Na.




Figure 3.28: Yearly phosphate FPMC (1982-1993).




Figure 3.29: Growing season phosphate FPMC (1982-1993).




Figure 3.30: Non-growing season phosphate FPMC (1982-1993).

Figures 3.31-3.33 and 3.34-3.36 depict the CI and TOC concentrations from 2002-2008. Since no pre-harvest data were available for these constituents, a regression analysis was only used to assess long-term trends. The CI concentration in both WSA and WSC exhibited a significant statistical increase in concentration from 2002-2008. The reason behind this increase in concentration is unclear. Results from the regression analysis for TOC showed a significant decrease in TOC concentration from 2002-2008 across all watersheds. A large decrease can be seen in TOC in 2008; this year TOC data were scarce and mainly came from the non-growing season. Looking at the TOC data, it is apparent that TOC concentration is season dependent with high concentrations occurring during the growing season and low concentrations occurring in the non-growing season. The data from 2008 may have skewed the results, as the trend from the previous years is seemingly linear.




Figure 3.31: Yearly chloride FPMC (2002-2008).




Figure 3.32: Growing season chloride FPMC (2002-2008).




Figure 3.33: Non-growing season chloride FPMC (2002-2008).




Figure 3.34: Yearly total organic carbon FPMC (2002-2008).




Figure 3.35: Growing season total organic carbon FPMC (2002-2008).




Figure 3.36: Non-growing season total organic carbon FPMC (2002-2008).



### 3.4 CONCLUSIONS

Harvesting the two treatment watersheds resulted in increased sediment loads, and leaching of base cations,  $\text{NO}_3$ , and  $\text{PO}_4$  into the stream. From the results it would seem that the most notable contribution of BMP implementation was that it limited the depletion of base cations and reduced sediment transport compared to the clear-cut watershed. When looking at  $\text{NO}_3$ , BMPs seemingly added no benefit and WSB remained elevated alongside WSC for five years after harvest.

Interestingly, some of the largest changes to stream water chemistry in these watersheds came from atmospheric deposition and not forest harvesting. During the study period  $\text{NO}_3$  and  $\text{SO}_4$  significantly declined across all watersheds. This significant reduction across all watersheds is attributed to regulations imposed by the 1990 amendment to the Clean Air Act that affected many nearby coal-fired power plants. Unlike  $\text{NO}_3$  and  $\text{SO}_4$ , however,  $\text{NH}_4$  significantly increased during its' monitoring period and was likely the result of increasing  $\text{NH}_4$  precipitation inputs due to increases to livestock production and aerosol emissions. Other significant changes to stream water chemistry came in the form of deposition from nearby surface mining activities which is thought to have caused the observed spikes in Mg, Ca, Na, and  $\text{PO}_4$  concentrations in 1988. Although significant increases in constituent concentrations were observed in the treatment watersheds, WSB and WSC are still considered to have excellent stream water quality compared to eastern Kentucky standards.

## **CHAPTER 4: CONCLUSIONS**

This study evaluated the long-term impacts of harvest and the effectiveness of BMPs at mitigating those impacts. The research was conducted at the University of Kentucky's Robinson Forest in eastern Kentucky and consisted of three adjacent headwater catchments. A paired watershed approach was used and yearly hydrograph characteristics plus water quality constituents were analyzed and compared across the watersheds. Results from this study affirm that implementing BMPs are an effective way to limit the damaging impacts of harvest and simply clear-cutting a watershed has significant long-term consequences. However, BMP effectiveness on promoting water quality was less certain and seemed to only add some benefit for some of the monitored constituents. Regardless, with this knowledge and knowing that headwater catchments make up a large majority of stream length and drainage area in a river network, it doesn't make sense that these first and second order streams should continue to go unregulated or under regulated.

Chapter One detailed the background of BMPs and their significance as well as the research objectives.

Chapter Two evaluated the storm hydrograph characteristics of baseflow, stormflow, peak discharge, storm volume as a percentage of rainfall, time to peak (2002-2008), and curve number of the paired watersheds. No differences in the analyzed characteristics were observed between the watersheds pre-harvest indicating the watersheds were strongly paired from a hydrologic standpoint. However, it should be noted that pre-harvest data were quite limited. Considering this was a long-term study, it cannot be definitively said that these watersheds would have behaved in the future as similarly as they did in the 19 pre-harvest months. Following treatment, large hydrologic differences were observed in both harvested watersheds.

Watershed B was statistically different from WSA in 1985 for all calculated hydrograph parameters besides peak discharge. By 1988, all of the evaluated hydrograph parameters besides stormflow in WSB had returned or nearly returned to control conditions. The stormflow in WSB remained elevated

throughout the first time period, and by the beginning of the second time period (2002), was realigned with the control. Watershed C was statistically different from WSA for all hydrograph parameters besides peak discharge and time to peak in 1984, 1985, 2006, and 2007 and was statistically different from WSA in 1989 for storm volume as a percentage of rainfall. Large differences in hydrograph parameters were observed throughout the study period even though massive regrowth is present at the site. The findings here indicate that BMP implementation was very effective at limiting the negative hydrologic effects due to harvest and clear-cutting a watershed can have significant long-term consequences.

Chapter Three compared water quality constituents of  $\text{SO}_4$ , Mg, Ca, K, Na, ALK (1982-2008),  $\text{NO}_3$  (1982-1990 & 2002-2008),  $\text{PO}_4$  (1982-1993), Cl,  $\text{NH}_4\text{-N}$ , and TOC (2002-2008) to assess significant differences between the watersheds due to harvest. Another goal of this study was to determine if the monitored constituents showed any long-term increasing or decreasing trends. The most notable contribution of BMP implementation was that it seemed to limit the depletion of base cations compared with the clear-cut watershed and reduced erosional effects. When looking at  $\text{NO}_3$ , the BMPs seemingly added no benefit and WSB remained elevated alongside WSC for roughly five years after harvest. Nitrate concentration significantly declined after this across all watersheds. This reduction in  $\text{NO}_3$  is attributed to regulations imposed by the Clean Air Act and is also responsible for the significant decline in  $\text{SO}_4$  that was observed across the watersheds. Unlike  $\text{NO}_3$  and  $\text{SO}_4$ , however,  $\text{NH}_4$  significantly increased during its' monitoring period and this was likely the result of increasing  $\text{NH}_4$  precipitation inputs.

## **CHAPTER 5: FUTURE WORK**

Future research should continue at the sites to assess correlations between storm characteristics (max intensity, average intensity, duration, and antecedent moisture conditions) and hydrograph responses across the watersheds. It would be interesting to see what kinds of storms illicit different responses in the watersheds and how they vary temporally. Further research should also be conducted at similar sites to assess the performance of a wider riparian buffer but with a percentage of the buffer being harvested. This will allow the same amount of timber to be harvested but may help limit sediment transport and stormflow directly after harvest. Finally, research on time-based hydrograph characteristics such as lag time and time of concentration should be conducted at similar sites. The headwater catchments studied here were extremely responsive to precipitation, and with the advancement of monitoring equipment, a smaller time-step could be used which would allow an accurate comparison to be made.

Further research should also be done to measure the performance of BMPs on limiting sediment transport in headwater catchments. A previous study at the site found significant differences in sediment loading across the watersheds, but due to extended equipment failure, monitoring was discontinued in 1990 (Arthur et al., 1998). From recent observation, it would seem that sediment transport may still be an issue in both harvested watersheds. Erosion control is one of the main goals of implementing BMPs, and as such, monitoring of suspended sediment should resume so an analysis can be done.

## APPENDIX A: INDIVIDUAL HYDROGRAPH CHARACTERISTICS

Table A.1: WSA, WSB, and WSC Stormflow and Baseflow (1982-1993).

| Date                         | WSA   |  | WSB   |  | WSC   |  |
|------------------------------|---|--|---|--|---|--|
|                              | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| February 8, 1982             | 376.2   | 139.8  | 257.3   | 130.8  | 270.4   | 191.7  |
| February 16, 1982            | 213.5   | 79.6   | 155.0   | 39.9   | 266.3   | 110.2  |
| March 15, 1982               | 281.1   | 81.3   | 145.9   | 24.4   | 272.7   | 83.1   |
| March 31, 1982               | 87.8  | 56.5   | 83.1  | 35.6   | 133.7   | 74.1   |
| May 21, 1982                 | 7.3   | 4.7  | 7.1   | 6.1  | 9.1   | 10.3   |
| May 28, 1982                 | 3.3   | 2.9  | 4.6   | 2.7  | 6.9   | 5.6  |
| June 4, 1982                 | 6.3   | 4.7  | 7.5   | 4.3  | 10.5  | 7.4  |
| July 28, 1982                | 1.9   | 0.3  | 1.9   | 1.4  | 1.3   | 0.4  |
| August 5, 1982               | 6.1   | 2.0  | 6.6   | 2.4  | 4.8   | 2.2  |
| September 13, 1982           | 13.2  | 3.9  | 9.7   | 2.6  | 5.3   | 1.1  |
| September 25, 1982           | 2.4   | 1.1  | 2.1   | 1.1  | 1.7   | 0.9  |
| November 20, 1982            | 15.9  | 3.6  | 12.0  | 2.1  | 7.0   | 0.9  |
| November 30, 1982            | 35.5  | 10.5   | 28.9  | 6.8  | 24.7  | 4.3  |
| December 5, 1982             | 31.5  | 13.0   | 27.0  | 11.9   | 25.4  | 8.9  |
| December 15, 1982            | 79.1  | 16.8   | 68.6  | 11.5   | 115.9   | 30.1   |
| January 21, 1983             | 114.4   | 24.8   | 118.8   | 34.2   | 100.5   | 23.4   |
| February 10, 1983            | 125.6   | 33.7   | 123.9   | 29.6   | 132.0   | 37.4   |
| April 14, 1983               | 116.3   | 23.6   | 131.9   | 34.7   | 120.9   | 32.5   |
| May 3, 1983                  | 176.6   | 54.6   | 173.7   | 54.4   | 173.7   | 54.8   |
| May 13, 1983                 | 412.4   | 200.2  | 375.7   | 173.2  | 371.4   | 169.7  |
| May 22, 1983                 | 138.2   | 96.4   | 163.4   | 100.7  | 149.1   | 72.3   |
| June 3, 1983                 | 10.8  | 1.9  | 7.7   | 1.7  | 7.7   | 1.2  |
| June 4, 1983                 | 55.7  | 18.4   | 66.6  | 21.3   | 69.3  | 18.9   |
| July 3, 1983                 | 0.6   | 0.2  | 1.7   | 0.1  | 1.8   | 0.2  |
| July 5, 1983                 | 2.1   | 2.0  | 3.1   | 2.0  | 1.9   | 1.6  |
| July 18, 1983                | 1.0   | 0.5  | 0.8   | 0.3  | 0.8   | 0.5  |
| August 2, 1983               | 1.2   | 0.5  | 0.7   | 0.2  | 0.8   | 0.2  |
| August 11, 1983              | 3.3   | 0.7  | 3.4   | 1.2  | 3.1   | 1.0  |
| August 27, 1983 <sup>1</sup> | 0.2   | 0.02   | 0.3   | 0.04   | 0.1   | 0.02   |
| November 14, 1983            | 13.7  | 4.3  | 22.6  | 5.2  | 16.1  | 5.2  |
| December 27, 1983            | 26.0  | 8.5  | 20.0  | 7.3  | 15.6  | 6.3  |
| February 27, 1984            | 181.5   | 51.7   | 155.5   | 46.5   | 162.3   | 37.7   |
| March 20, 1984               | 211.8   | 63.9   | 199.2   | 71.0   | 235.4   | 68.7   |
| March 28, 1984               | 141.5   | 55.7   | 127.8   | 60.4   | 154.1   | 54.8   |
| April 4, 1984                | 159.8   | 51.9   | 156.7   | 47.4   | 167.9   | 46.8   |
| April 9, 1984                | 238.5   | 110.9  | 233.3   | 103.8  | 307.0   | 104.3  |
| April 21, 1984               | 288.4   | 179.4  | 309.6   | 188.1  | 335.6   | 201.0  |
| May 4, 1984                  | 174.9   | 28.1   | 187.3   | 43.2   | 179.8   | 31.5   |
| May 6, 1984                  | 851.5   | 515.5  | 225.3   | 30.1   | 235.2   | 26.6   |
| May 23, 1984                 | 2.6   | 7.0  | 4.2   | 16.9   | 3.5   | 11.9   |
| May 28, 1984                 | 12.1  | 5.2  | 117.5   | 52.5   | 139.3   | 78.1   |

Table A.1: Continued

| Date               | WSA   |  | WSB   |  | WSC   |  |
|--------------------|---|--|---|--|---|--|
|                    | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| July 26, 1984      | 1.5   | 0.21   | 2.6   | 0.37   | 5.2   | 1.67   |
| July 27, 1984      | 2.6   | 2.00   | 49.0  | 9.44   | 64.6  | 16.34  |
| July 30, 1984      | 0.6   | 0.76   | 9.9   | 6.55   | 19.4  | 8.25   |
| August 22, 1984    | 4.1   | 0.25   | 26.4  | 16.50  | 52.4  | 38.27  |
| November 4, 1984   | 8.5   | 2.87   | 99.7  | 19.85  | 113.1   | 24.06  |
| November 18, 1984  | 128.3   | 42.01  | 357.2   | 266.85   | 403.6   | 233.03   |
| November 28, 1984  | 69.4  | 25.23  | 144.6   | 97.83  | 148.1   | 88.68  |
| December 20, 1984  | 121.3   | 30.74  | 187.5   | 42.08  | 199.9   | 35.04  |
| December 24, 1984  | 114.5   | 39.04  | 166.3   | 54.76  | 212.0   | 68.36  |
| January 3, 1985    | 117.9   | 32.87  | 189.3   | 56.16  | 225.2   | 58.87  |
| February 11, 1985  | 154.8   | 66.42  | 192.3   | 87.70  | 188.0   | 86.58  |
| May 15, 1985       | 2.2   | 1.14   | 5.0   | 3.42   | 10.2  | 5.90   |
| June 5, 1985       | 5.6   | 3.88   | 20.5  | 6.86   | 19.7  | 7.93   |
| June 10, 1985      | 3.1   | 1.24   | 9.3   | 3.16   | 8.5   | 3.93   |
| June 11, 1985      | 47.1  | 10.07  | 163.9   | 45.65  | 149.6   | 70.13  |
| July 10, 1985      | 3.4   | 1.04   | 8.3   | 14.89  | 12.1  | 18.76  |
| July 30, 1985      | 1.4   | 0.12   | 1.9   | 1.23   | 2.0   | 1.58   |
| August 1, 1985     | 6.9   | 2.73   | 21.9  | 14.48  | 21.3  | 15.71  |
| August 17, 1985    | 3.3   | 0.78   | 14.7  | 6.09   | 12.8  | 6.24   |
| August 25, 1985    | 2.4   | 1.35   | 3.6   | 1.27   | 16.9  | 9.89   |
| August 30, 1985    | 9.1   | 4.52   | 58.1  | 46.61  | 74.4  | 42.13  |
| September 26, 1985 | 0.8   | 0.07   | 4.9   | 9.78   | 8.0   | 9.80   |
| November 2, 1985   | 115.0   | 35.72  | 343.7   | 142.47   | 360.8   | 131.10   |
| December 12, 1985  | 135.9   | 46.26  | 174.9   | 86.88  | 176.5   | 111.00   |
| February 2, 1986   | 347.5   | 74.94  | 223.8   | 34.20  | 454.6   | 96.87  |
| April 28, 1986     | 8.4   | 4.71   | 10.5  | 11.03  | 21.4  | 21.15  |
| May 11, 1986       | 8.2   | 4.27   | 25.6  | 18.10  | 18.6  | 11.36  |
| July 2, 1986       | 3.2   | 1.26   | 9.5   | 2.01   | 9.9   | 9.74   |
| July 20, 1986      | 2.4   | 1.33   | 3.0   | 0.81   | 13.2  | 11.26  |
| July 26, 1986      | 1.9   | 0.61   | 3.3   | 1.05   | 11.0  | 12.44  |
| October 1, 1986    | 6.3   | 2.64   | 6.9   | 2.92   | 12.5  | 7.47   |
| November 5, 1986   | 17.7  | 4.18   | 45.1  | 5.26   | 64.7  | 7.10   |
| November 8, 1986   | 215.8   | 78.71  | 135.6   | 8.89   | 414.5   | 140.20   |
| November 10, 1986  | 49.6  | 50.07  | 89.5  | 82.43  | 124.3   | 131.59   |
| December 8, 1986   | 217.4   | 116.84   | 236.1   | 106.37   | 305.4   | 154.28   |
| January 18, 1987   | 185.9   | 186.25   | 207.8   | 141.95   | 267.3   | 140.65   |
| February 22, 1987  | 146.3   | 36.88  | 163.4   | 38.99  | 163.6   | 38.65  |
| February 26, 1987  | 308.2   | 127.11   | 302.2   | 177.45   | 334.3   | 154.83   |
| March 18, 1987     | 108.3   | 78.54  | 105.6   | 81.91  | 88.1  | 96.71  |
| March 30, 1987     | 473.3   | 147.49   | 501.0   | 158.93   | 544.6   | 193.83   |
| May 12, 1987       | 2.2   | 5.16   | 4.2   | 11.79  | 8.1   | 19.57  |

Table A.1: Continued

| Date               | WSA   |  | WSB   |  | WSC   |  |
|--------------------|---|--|---|--|---|--|
|                    | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| May 21, 1987       | 4.7   | 5.4  | 8.6   | 10.8   | 9.0   | 16.6   |
| May 25, 1987       | 5.5   | 7.3  | 8.9   | 7.2  | 12.3  | 11.4   |
| June 16, 1987      | 4.6   | 4.8  | 4.9   | 5.4  | 7.1   | 7.4  |
| June 25, 1987      | 12.3  | 2.3  | 16.9  | 12.4   | 19.8  | 20.1   |
| July 11, 1987      | 18.9  | 12.6   | 45.0  | 18.3   | 69.0  | 38.2   |
| August 22, 1987    | 2.1   | 0.4  | 4.4   | 1.6  | 6.4   | 5.6  |
| September 12, 1987 | 4.4   | 1.1  | 5.2   | 0.5  | 8.0   | 2.4  |
| November 9, 1987   | 11.2  | 7.1  | 10.0  | 2.5  | 19.5  | 8.0  |
| November 16, 1987  | 3.6   | 4.8  | 9.8   | 3.9  | 7.7   | 4.8  |
| December 14, 1987  | 5.6   | 2.8  | 13.0  | 4.6  | 12.9  | 4.9  |
| December 24, 1987  | 98.3  | 29.8   | 122.4   | 29.7   | 211.8   | 47.5   |
| January 18, 1988   | 208.4   | 121.0  | 163.4   | 194.9  | 233.3   | 112.9  |
| April 6, 1988      | 292.3   | 113.9  | 229.3   | 183.3  | 315.9   | 142.5  |
| May 4, 1988        | 293.3   | 135.3  | 134.3   | 155.9  | 248.9   | 175.5  |
| June 2, 1988       | 1.5   | 2.3  | 1.5   | 14.4   | 8.7   | 22.3   |
| June 9, 1988       | 1.8   | 2.9  | 2.1   | 4.1  | 2.8   | 4.9  |
| August 23, 1988    | 3.7   | 2.1  | 6.0   | 2.3  | 20.1  | 6.1  |
| September 16, 1988 | 7.4   | 6.9  | 8.1   | 4.9  | 16.3  | 15.3   |
| November 4, 1988   | 17.4  | 3.7  | 25.9  | 5.4  | 32.6  | 9.6  |
| November 27, 1988  | 27.1  | 10.9   | 55.3  | 33.1   | 54.4  | 52.3   |
| December 21, 1988  | 53.1  | 6.7  | 99.3  | 16.0   | 93.0  | 15.6   |
| December 24, 1988  | 224.6   | 94.1   | 198.0   | 90.1   | 212.0   | 68.5   |
| January 11, 1989   | 332.1   | 139.7  | 346.1   | 181.1  | 418.6   | 167.7  |
| February 3, 1989   | 151.1   | 90.0   | 182.4   | 87.6   | 227.3   | 97.9   |
| February 13, 1989  | 373.2   | 114.5  | 400.6   | 105.9  | 444.8   | 109.0  |
| February 20, 1989  | 436.5   | 277.8  | 349.8   | 341.5  | 378.3   | 290.4  |
| March 20, 1989     | 213.6   | 85.4   | 254.2   | 158.6  | 272.5   | 143.9  |
| April 3, 1989      | 139.3   | 75.2   | 191.7   | 140.3  | 194.3   | 87.1   |
| May 19, 1989       | 8.5   | 9.0  | 8.6   | 17.2   | 7.9   | 17.1   |
| June 12, 1989      | 533.7   | 239.0  | 495.9   | 242.4  | 547.1   | 315.4  |
| June 22, 1989      | 24.5  | 36.5   | 33.4  | 65.7   | 31.8  | 49.6   |
| July 23, 1989      | 18.2  | 14.6   | 16.3  | 10.5   | 15.6  | 17.9   |
| July 27, 1989      | 3.0   | 2.2  | 4.4   | 2.7  | 3.8   | 4.3  |
| July 31, 1989      | 21.1  | 13.7   | 56.7  | 23.5   | 79.1  | 36.3   |
| August 5, 1989     | 3.4   | 1.3  | 70.6  | 55.1   | 65.9  | 39.5   |
| August 18, 1989    | 5.8   | 5.1  | 8.9   | 8.8  | 13.6  | 16.9   |
| September 22, 1989 | 136.7   | 57.2   | 224.9   | 100.0  | 325.8   | 161.9  |
| September 30, 1989 | 149.3   | 90.6   | 182.0   | 102.1  | 201.4   | 144.0  |
| October 16, 1989   | 463.2   | 117.3  | 139.9   | 14.9   | 729.0   | 424.2  |
| November 14, 1989  | 192.5   | 75.5   | 255.6   | 108.5  | 277.8   | 142.4  |
| January 29, 1990   | 147.7   | 36.1   | 184.3   | 42.3   | 173.7   | 34.1   |



Table A.1: Continued

| Date               | WSA   |  | WSB   |  | WSC   |  |
|--------------------|---|--|---|--|---|--|
|                    | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| February 3, 1990   | 265.4   | 105.5  | 291.7   | 98.5   | 317.1   | 84.9   |
| February 9, 1990   | 197.2   | 119.9  | 259.5   | 128.1  | 274.0   | 115.5  |
| February 15, 1990  | 173.5   | 109.5  | 183.5   | 204.3  | 197.9   | 153.8  |
| March 16, 1990     | 212.1   | 76.8   | 275.6   | 122.5  | 213.5   | 78.3   |
| April 6, 1990      | 129.5   | 48.9   | 127.1   | 51.8   | 142.6   | 55.8   |
| May 26, 1990       | 2.3   | 2.0  | 2.5   | 2.2  | 2.5   | 7.2  |
| May 28, 1990       | 36.4  | 16.6   | 41.5  | 13.8   | 46.9  | 24.2   |
| June 2, 1990       | 55.1  | 26.6   | 56.6  | 19.0   | 70.0  | 26.0   |
| June 21, 1990      | 6.5   | 9.7  | 4.0   | 6.1  | 7.3   | 7.5  |
| July 30, 1990      | 0.6   | 1.2  | 1.5   | 3.5  | 3.0   | 5.3  |
| August 8, 1990     | 1.8   | 0.4  | 4.5   | 8.9  | 6.6   | 5.5  |
| August 29, 1990    | 1.9   | 2.6  | 2.3   | 4.4  | 4.0   | 5.7  |
| September 9, 1990  | 4.9   | 1.6  | 2.3   | 3.8  | 3.8   | 3.7  |
| September 12, 1990 | 1.0   | 1.8  | 1.9   | 3.6  | 2.5   | 3.8  |
| October 4, 1990    | 3.4   | 3.0  | 12.1  | 8.4  | 12.6  | 10.6   |
| December 2, 1990   | 117.5   | 70.3   | 186.7   | 107.0  | 230.1   | 105.7  |
| December 20, 1990  | 278.8   | 103.2  | 384.4   | 202.1  | 330.0   | 111.9  |
| December 27, 1990  | 94.1  | 65.5   | 128.3   | 76.9   | 136.1   | 71.2   |
| December 30, 1990  | 98.5  | 135.5  | 130.1   | 180.7  | 168.0   | 153.6  |
| January 6, 1991    | 276.9   | 170.7  | 311.2   | 173.2  | 293.7   | 166.5  |
| February 6, 1991   | 173.4   | 50.0   | 189.5   | 48.6   | 161.4   | 47.7   |
| February 13, 1991  | 138.2   | 40.9   | 152.6   | 38.6   | 157.1   | 36.4   |
| February 17, 1991  | 108.5   | 13.0   | 477.9   | 219.6  | 607.2   | 230.2  |
| March 22, 1991     | 189.4   | 63.6   | 204.8   | 76.8   | 200.6   | 68.4   |
| March 29, 1991     | 277.7   | 105.7  | 262.0   | 95.6   | 257.2   | 88.2   |
| April 15, 1991     | 178.5   | 72.5   | 209.9   | 72.0   | 204.5   | 59.0   |
| April 19, 1991     | 185.8   | 194.7  | 183.9   | 205.7  | 175.5   | 113.8  |
| May 9, 1991        | 12.5  | 8.8  | 9.6   | 16.4   | 3.3   | 2.9  |
| May 18, 1991       | 4.1   | 4.1  | 14.5  | 20.7   | 4.2   | 5.2  |
| May 27, 1991       | 4.5   | 5.0  | 6.6   | 7.3  | 4.9   | 3.9  |
| May 29, 1991       | 20.8  | 13.4   | 25.7  | 11.9   | 16.5  | 10.1   |
| June 22, 1991      | 32.5  | 3.9  | 77.3  | 38.5   | 58.2  | 26.1   |
| June 25, 1991      | 143.0   | 61.9   | 152.3   | 82.9   | 132.7   | 68.3   |
| July 10, 1991      | 3.2   | 1.8  | 2.9   | 3.2  | 2.4   | 3.1  |
| July 12, 1991      | 16.8  | 19.3   | 28.2  | 23.4   | 27.4  | 13.5   |
| August 7, 1991     | 14.2  | 10.9   | 16.9  | 9.2  | 12.5  | 11.0   |
| October 5, 1991    | 8.8   | 1.7  | 12.6  | 6.8  | 7.3   | 1.8  |
| October 15, 1991   | 11.9  | 14.2   | 31.0  | 24.3   | 10.1  | 11.3   |
| November 21, 1991  | 94.2  | 48.1   | 108.2   | 69.1   | 88.4  | 63.7   |
| December 13, 1991  | 121.3   | 109.1  | 128.4   | 149.5  | 172.3   | 80.5   |
| December 28, 1991  | 254.4   | 129.4  | 261.7   | 156.7  | 259.6   | 122.5  |

Table A.1: Continued

| Date                    | WSA   |  | WSB   |  | WSC   |  |
|-------------------------|---|--|---|--|---|--|
|                         | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| February 24, 1992       | 261.2   | 121.0  | 306.4   | 110.1  | 298.4   | 99.1   |
| March 30, 1992          | 212.7   | 96.1   | 259.9   | 118.5  | 246.2   | 104.8  |
| May 28, 1992            | 2.4   | 0.6  | 4.7   | 2.7  | 25.8  | 15.0   |
| June 17, 1992           | 12.4  | 8.3  | 20.2  | 10.1   | 14.3  | 6.2  |
| June 24, 1992           | 5.3   | 0.7  | 14.7  | 5.9  | 10.7  | 4.3  |
| July 1, 1992            | 375.9   | 344.6  | 253.2   | 166.8  | 261.1   | 188.9  |
| July 14, 1992           | 2.0   | 2.3  | 2.0   | 5.3  | 6.7   | 15.2   |
| July 17, 1992           | 2.9   | 5.4  | 5.4   | 21.2   | 3.7   | 6.9  |
| July 22, 1992           | 4.7   | 4.0  | 6.0   | 5.1  | 4.8   | 3.9  |
| July 24, 1992           | 87.9  | 143.2  | 119.6   | 68.3   | 141.0   | 83.5   |
| August 8, 1992          | 23.7  | 24.0   | 25.1  | 22.2   | 33.0  | 26.0   |
| August 27, 1992         | 47.3  | 26.2   | 43.1  | 17.9   | 57.9  | 35.1   |
| September 4, 1992       | 2.7   | 4.7  | 5.9   | 12.1   | 9.6   | 22.5   |
| September 18, 1992      | 8.5   | 8.1  | 7.1   | 3.7  | 46.4  | 18.5   |
| December 20, 1992       | 110.7   | 17.6   | 109.6   | 25.6   | 80.3  | 25.5   |
| March 29, 1993          | 252.2   | 89.3   | 323.9   | 160.7  | 320.6   | 132.2  |
| April 25, 1993          | 193.3   | 58.1   | 115.3   | 40.3   | 34.2  | 6.9  |
| May 9, 1993             | 22.5  | 15.8   | 24.0  | 17.4   | 9.4   | 7.4  |
| June 9, 1993            | 3.9   | 5.2  | 7.2   | 5.4  | 9.4   | 1.0  |
| June 21, 1993           | 4.8   | 5.2  | 8.2   | 5.4  | 7.2   | 3.9  |
| July 13, 1993           | 9.0   | 3.0  | 15.5  | 3.6  | 51.5  | 37.6   |
| July 15, 1993           | 14.1  | 11.4   | 19.7  | 16.9   | 58.7  | 40.3   |
| July 26, 1993           | 7.2   | 10.2   | 10.2  | 13.5   | 9.6   | 5.2  |
| September 2, 1993       | 9.6   | 9.1  | 37.3  | 6.1  | 39.8  | 6.5  |
| September 15, 1993      | 21.2  | 12.2   | 45.1  | 29.3   | 47.8  | 23.6   |
| October 18, 1993        | 86.8  | 34.5   | 221.2   | 80.6   | 234.6   | 80.5   |
| Mean ± Std. Dev. (Pre)  | 80.0±113.9                                      | 30.3±48.1                                      | 68.6±92.2                                       | 25.4±41.9                                      | 78.9±104.9                                      | 32.6±50.7                                      |
| Mean ± Std. Dev. (Post) | 98.5±130.5                                      | 46.7±69.1                                      | 109.6±118.6                                     | 55.8±66.2                                      | 126.9±142.8                                     | 58.2±68.4                                      |

<sup>1</sup> Final storm before treatment.

Table A.2: WSA, WSB, and WSC Stormflow and Baseflow (2002-2008).

| Date               | WSA   |  | WSB   |  | WSC   |  |
|--------------------|---|--|---|--|---|--|
|                    | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| March 17, 2002     | 161.0   | 18.5   | 201.2   | 30.3   | -- <sup>1</sup>                                 | -- <sup>1</sup>                                |
| April 28, 2002     | 117.2   | 53.4   | 140.7   | 48.4   | --  | --   |
| May 2, 2002        | 451.2   | 359.3  | 513.0   | 353.5  | --  | --   |
| May 13, 2002       | 169.5   | 127.8  | 231.7   | 172.3  | --  | --   |
| May 17, 2002       | 67.6  | 98.3   | 130.5   | 163.4  | --  | --   |
| June 6, 2002       | 11.7  | 13.5   | 19.3  | 18.0   | --  | --   |
| July 19, 2002      | 13.4  | 9.1  | 9.5   | 6.3  | --  | --   |
| September 25, 2002 | 14.9  | 8.9  | 8.6   | 1.5  | --  | --   |
| October 15, 2002   | 8.1   | 5.6  | 3.5   | 1.7  | --  | --   |
| October 29, 2002   | 32.9  | 13.4   | 38.8  | 7.0  | --  | --   |
| November 5, 2002   | 31.3  | 16.2   | 40.9  | 9.6  | --  | --   |
| November 10, 2002  | 27.8  | 18.2   | 105.3   | 23.3   | --  | --   |
| December 10, 2002  | 105.4   | 10.7   | 119.5   | 10.6   | --  | --   |
| December 13, 2002  | 178.8   | 46.8   | 204.7   | 35.1   | --  | --   |
| December 19, 2002  | 53.3  | 35.9   | 53.7  | 39.0   | --  | --   |
| January 1, 2003    | 85.7  | 53.0   | 77.4  | 55.4   | --  | --   |
| February 3, 2003   | 121.0   | 46.1   | 121.3   | 42.3   | --  | --   |
| April 6, 2003      | 131.1   | 11.9   | 147.3   | 13.1   | --  | --   |
| April 17, 2003     | 106.9   | 56.9   | 112.8   | 112.0  | --  | --   |
| May 8, 2003        | 40.7  | 12.4   | 81.1  | 53.4   | --  | --   |
| June 6, 2003       | 54.2  | 24.3   | 84.1  | 29.9   | --  | --   |
| June 11, 2003      | 2.2   | 1.9  | 8.4   | 6.6  | --  | --   |
| July 10, 2003      | 4.3   | 7.3  | 5.3   | 9.0  | --  | --   |
| August 3, 2003     | 2.2   | 0.6  | 4.0   | 1.3  | --  | --   |
| August 17, 2003    | 4.2   | 6.5  | 13.1  | 9.5  | --  | --   |
| September 3, 2003  | 107.3   | 88.1   | 113.9   | 76.0   | --  | --   |
| September 22, 2003 | 7.4   | 8.6  | 2.8   | 2.7  | --  | --   |
| November 12, 2003  | 19.1  | 6.1  | 23.5  | 7.8  | --  | --   |
| November 18, 2003  | 201.6   | 91.8   | 215.8   | 85.2   | --  | --   |
| November 28, 2003  | 75.3  | 43.5   | 74.3  | 41.2   | --  | --   |
| December 10, 2003  | 74.0  | 23.2   | 91.7  | 23.6   | --  | --   |
| January 17, 2004   | 158.2   | 68.3   | 177.3   | 96.3   | --  | --   |
| February 2, 2004   | 80.3  | 17.2   | 101.5   | 25.7   | --  | --   |
| February 5, 2004   | 120.8   | 6.1  | 71.2  | 4.8  | --  | --   |
| March 5, 2004      | 212.1   | 173.7  | 256.0   | 157.5  | --  | --   |
| May 16, 2004       | 4.6   | 6.6  | 3.1   | 5.6  | --  | --   |
| May 24, 2004       | 4.4   | 4.8  | 6.5   | 3.5  | --  | --   |
| May 26, 2004       | 90.1  | 14.9   | 88.2  | 12.9   | --  | --   |
| May 27, 2004       | 180.1   | 90.0   | 221.8   | 60.2   | --  | --   |

Table A.2: Continued

| Date               | WSA   |  | WSB   |  | WSC   |  |
|--------------------|---|--|---|--|---|--|
|                    | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| June 15, 2004      | 148.8   | 118.0  | 290.3   | 226.8  | --  | --   |
| June 22, 2004      | 27.7  | 25.7   | 35.9  | 35.1   | --  | --   |
| June 25, 2004      | 93.5  | 47.1   | 128.9   | 73.2   | --  | --   |
| July 6, 2004       | 7.6   | 3.6  | 9.0   | 6.1  | --  | --   |
| July 26, 2004      | 1.6   | 0.2  | 3.8   | 0.8  | --  | --   |
| July 27, 2004      | 20.8  | 11.8   | 13.1  | 5.3  | --  | --   |
| July 31, 2004      | 7.6   | 7.0  | 2.9   | 3.8  | --  | --   |
| August 5, 2004     | 9.2   | 16.2   | 11.6  | 15.0   | --  | --   |
| September 7, 2004  | 128.1   | 57.6   | 128.3   | 53.2   | --  | --   |
| September 16, 2004 | 346.5   | 295.1  | 409.7   | 225.7  | --  | --   |
| October 13, 2004   | 9.7   | 6.9  | 12.6  | 4.8  | --  | --   |
| October 18, 2004   | 123.1   | 110.3  | 152.2   | 81.4   | --  | --   |
| October 27, 2004   | 191.3   | 107.0  | 188.8   | 111.1  | --  | --   |
| November 4, 2004   | 231.8   | 172.5  | 208.3   | 120.6  | --  | --   |
| November 11, 2004  | 52.6  | 52.2   | 38.1  | 50.9   | --  | --   |
| November 30, 2004  | 328.2   | 91.1   | 363.8   | 116.6  | --  | --   |
| December 9, 2004   | 299.0   | 185.0  | 324.1   | 149.2  | --  | --   |
| December 23, 2004  | 20.2  | 16.8   | 34.9  | 69.3   | --  | --   |
| April 29, 2005     | 307.7   | 206.3  | 399.4   | 825.5  | 495.9   | 364.5  |
| May 19, 2005       | 31.6  | 73.3   | 51.4  | 105.9  | 28.5  | 31.7   |
| May 22, 2005       | 49.6  | 60.6   | 87.2  | 115.0  | 26.2  | 29.9   |
| June 3, 2005       | 104.4   | 133.5  | 121.2   | 141.5  | 42.2  | 47.4   |
| June 20, 2005      | 6.4   | 3.3  | 29.6  | 30.9   | 17.4  | 12.2   |
| July 1, 2005       | 7.3   | 10.5   | 4.6   | 4.4  | 4.1   | 4.3  |
| July 7, 2005       | 8.8   | 13.6   | 1.0   | 2.8  | 30.6  | 4.1  |
| July 17, 2005      | 26.9  | 6.5  | 29.4  | 7.2  | 17.4  | 8.5  |
| July 18, 2005      | 33.4  | 28.9   | 28.4  | 29.2   | 32.9  | 32.2   |
| July 27, 2005      | 4.6   | 9.9  | 1.1   | 2.3  | 8.9   | 14.6   |
| August 5, 2005     | 4.4   | 10.1   | 2.4   | 2.2  | 2.4   | 4.4  |
| August 16, 2005    | 2.5   | 4.6  | 0.7   | 1.3  | 0.9   | 2.0  |
| August 19, 2005    | 5.3   | 10.6   | 1.7   | 1.2  | 3.1   | 4.4  |
| December 3, 2005   | 3.3   | 0.6  | 3.1   | 0.3  | 21.7  | 4.2  |
| December 15, 2005  | 24.6  | 5.8  | 13.0  | 1.8  | 26.0  | 9.9  |
| January 17, 2006   | 53.8  | 11.8   | 86.0  | 15.4   | 63.3  | 6.8  |
| January 23, 2006   | 126.0   | 99.3   | 185.8   | 72.1   | 271.3   | 70.9   |
| March 13, 2006     | 104.3   | 24.5   | 143.6   | 17.0   | 149.8   | 37.8   |
| May 2, 2006        | 7.5   | 19.7   | 13.0  | 29.0   | 53.6  | 29.3   |
| May 25, 2006       | 18.1  | 12.8   | 14.5  | 8.1  | 45.6  | 13.4   |

Table A.2: Continued

| Date                     | WSA   |  | WSB   |  | WSC   |  |
|--------------------------|---|--|---|--|---|--|
|                          | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Stormflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) | Baseflow<br>(m <sup>3</sup> ha <sup>-1</sup> ) |
| May 31, 2006             | 29.5  | 15.7   | 57.5  | 10.4   | 82.9  | 11.4   |
| July 5, 2006             | 63.0  | 21.6   | 98.1  | 35.3   | 139.2   | 43.5   |
| September 23, 2006       | 5.1   | 0.4  | 8.8   | 0.6  | 9.4   | 0.6  |
| September 23, 2006       | 10.0  | 1.8  | 22.9  | 6.1  | 23.3  | 4.9  |
| September 30, 2006       | 4.3   | 2.3  | 10.4  | 4.6  | 9.8   | 3.4  |
| October 26, 2006         | 128.0   | 40.9   | 194.4   | 44.7   | 268.5   | 110.6  |
| March 28, 2007           | 37.0  | 16.4   | 8.1   | 9.6  | 18.4  | 6.9  |
| May 16, 2007             | 4.1   | 2.1  | 8.7   | 8.6  | 26.7  | 11.1   |
| June 1, 2007             | 6.8   | 2.7  | 6.0   | 2.6  | 42.1  | 8.6  |
| June 5, 2007             | 12.1  | 5.2  | 11.1  | 5.3  | 22.8  | 15.3   |
| June 18, 2007            | 1.0   | 0.3  | 0.1   | 0.3  | 2.4   | 1.3  |
| June 29, 2007            | 7.0   | 13.1   | 1.9   | 2.4  | 9.2   | 9.0  |
| August 3, 2007           | 2.3   | 1.5  | 0.7   | 0.5  | 9.8   | 3.8  |
| November 14, 2007        | 15.8  | 4.8  | 7.7   | 1.4  | 21.7  | 4.4  |
| November 26, 2007        | 4.2   | 1.7  | 7.3   | 1.0  | 22.1  | 9.5  |
| December 10, 2007        | 12.4  | 2.7  | 20.0  | 2.2  | 38.0  | 6.1  |
| December 13, 2007        | 24.0  | 4.9  | 43.9  | 11.6   | 50.9  | 15.7   |
| December 20, 2007        | 4.8   | 3.1  | 26.8  | 10.8   | 41.5  | 21.9   |
| January 29, 2008         | 79.6  | 14.6   | 133.5   | 16.4   | 157.1   | 31.5   |
| February 6, 2008         | 73.5  | 21.1   | 128.7   | 21.9   | 135.3   | 32.7   |
| May 11, 2008             | 5.4   | 3.5  | 12.0  | 11.0   | 13.9  | 5.9  |
| June 3, 2008             | 5.6   | 4.1  | 8.9   | 5.3  | 11.9  | 20.9   |
| August 26, 2008          | 3.5   | 0.7  | 1.8   | 0.1  | 4.4   | 0.9  |
| November 14, 2008        | 5.8   | 1.5  | 25.0  | 2.7  | 13.7  | 2.4  |
| December 10, 2008        | 6.9   | 1.0  | 29.7  | 1.4  | 23.1  | 1.4  |
| December 10, 2008        | 17.6  | 4.6  | 77.0  | 9.8  | 37.1  | 7.9  |
| December 16, 2008        | 25.5  | 3.5  | 35.8  | 11.7   | 52.7  | 11.1   |
| Mean ± Std. Dev. (02-04) | 94.4±98.5                                       | 53.1±70.8                                      | 110.2±113.2                                     | 56.2±69.8                                      |   |  |
| Mean ± Std. Dev. (05-08) | 32.4±53.0                                       | 20.0±38.1                                      | 46.9±73.0                                       | 35.1±121.7                                     | 56.0±89.5                                       | 24.1±54.6                                      |

<sup>1</sup>No data available.

TableA.3: WSA, WSB, and WSC peak flow ( $\text{m}^3 \text{ s}^{-1} \text{ ha}^{-1} \times 10^{-2}$ ) (1982-1993).

| Date                         | WSA  | WSB  | WSC  |
|------------------------------|------|------|------|
| February 8, 1982             | 1.81 | 0.94 | 1.10 |
| February 16, 1982            | 0.30 | 0.27 | 0.25 |
| March 15, 1982               | 0.46 | 0.45 | 0.45 |
| March 31, 1982               | 0.12 | 0.13 | 0.08 |
| May 21, 1982                 | 0.09 | 0.05 | 0.07 |
| May 28, 1982                 | 0.02 | 0.03 | 0.02 |
| June 4, 1982                 | 0.03 | 0.01 | 0.02 |
| July 28, 1982                | 0.04 | 0.02 | 0.04 |
| August 5, 1982               | 0.07 | 0.05 | 0.05 |
| September 13, 1982           | 0.40 | 0.27 | 0.18 |
| September 25, 1982           | 0.01 | 0.00 | 0.01 |
| November 20, 1982            | 0.01 | 0.01 | 0.01 |
| November 30, 1982            | 0.03 | 0.03 | 0.02 |
| December 5, 1982             | 0.03 | 0.02 | 0.02 |
| December 15, 1982            | 0.09 | 0.07 | 0.07 |
| January 21, 1983             | 0.06 | 0.08 | 0.05 |
| February 10, 1983            | 0.09 | 0.09 | 0.06 |
| April 14, 1983               | 0.10 | 0.09 | 0.06 |
| May 3, 1983                  | 0.18 | 0.21 | 0.18 |
| May 13, 1983                 | 0.82 | 0.55 | 0.61 |
| May 22, 1983                 | 0.49 | 0.47 | 0.30 |
| June 3, 1983                 | 0.14 | 0.09 | 0.14 |
| June 4, 1983                 | 0.16 | 0.13 | 0.15 |
| July 3, 1983                 | 0.00 | 0.04 | 0.05 |
| July 5, 1983                 | 0.02 | 0.01 | 0.01 |
| July 18, 1983                | 0.01 | 0.01 | 0.02 |
| August 2, 1983               | 0.03 | 0.02 | 0.02 |
| August 11, 1983              | 0.10 | 0.10 | 0.07 |
| August 27, 1983 <sup>1</sup> | 0.01 | 0.00 | 0.01 |
| November 14, 1983            | 0.02 | 0.02 | 0.02 |
| December 27, 1983            | 0.01 | 0.01 | 0.01 |
| February 27, 1984            | 0.19 | 0.19 | 0.12 |
| March 20, 1984               | 0.20 | 0.22 | 0.13 |
| March 28, 1984               | 0.07 | 0.06 | 0.06 |
| April 4, 1984                | 0.08 | 0.09 | 0.08 |
| April 9, 1984                | 0.49 | 0.57 | 0.49 |
| April 21, 1984               | 0.82 | 0.92 | 0.69 |
| May 4, 1984                  | 0.37 | 0.38 | 0.29 |
| May 6, 1984                  | 3.50 | 0.98 | 0.86 |
| May 23, 1984                 | 0.02 | 0.02 | 0.02 |
| May 28, 1984                 | 0.02 | 0.05 | 0.04 |

Table A.3: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| July 26, 1984      | 0.04 | 0.04 | 0.06 |
| July 27, 1984      | 0.03 | 0.05 | 0.06 |
| July 30, 1984      | 0.00 | 0.02 | 0.04 |
| August 22, 1984    | 0.11 | 0.08 | 0.15 |
| November 4, 1984   | 0.03 | 0.07 | 0.07 |
| November 18, 1984  | 0.27 | 0.68 | 0.70 |
| November 28, 1984  | 0.07 | 0.14 | 0.10 |
| December 20, 1984  | 0.14 | 0.19 | 0.13 |
| December 24, 1984  | 0.16 | 0.22 | 0.17 |
| January 3, 1985    | 0.09 | 0.11 | 0.10 |
| February 11, 1985  | 0.10 | 0.16 | 0.11 |
| May 15, 1985       | 0.04 | 0.04 | 0.05 |
| June 5, 1985       | 0.08 | 0.08 | 0.12 |
| June 10, 1985      | 0.03 | 0.04 | 0.05 |
| June 11, 1985      | 0.10 | 0.19 | 0.18 |
| July 10, 1985      | 0.09 | 0.08 | 0.07 |
| July 30, 1985      | 0.01 | 0.01 | 0.01 |
| August 1, 1985     | 0.08 | 0.08 | 0.07 |
| August 17, 1985    | 0.03 | 0.03 | 0.02 |
| August 25, 1985    | 0.02 | 0.03 | 0.05 |
| August 30, 1985    | 0.17 | 0.12 | 0.08 |
| September 26, 1985 | 0.01 | 0.02 | 0.02 |
| November 2, 1985   | 0.07 | 0.21 | 0.25 |
| December 12, 1985  | 0.09 | 0.11 | 0.08 |
| February 2, 1986   | 0.28 | 0.30 | 0.25 |
| April 28, 1986     | 0.02 | 0.02 | 0.02 |
| May 11, 1986       | 0.09 | 0.05 | 0.05 |
| July 2, 1986       | 0.04 | 0.03 | 0.04 |
| July 20, 1986      | 0.07 | 0.04 | 0.06 |
| July 26, 1986      | 0.07 | 0.03 | 0.05 |
| October 1, 1986    | 0.03 | 0.01 | 0.03 |
| November 5, 1986   | 0.05 | 0.06 | 0.06 |
| November 8, 1986   | 0.67 | 0.61 | 1.04 |
| November 10, 1986  | 0.13 | 0.17 | 0.18 |
| December 8, 1986   | 0.34 | 0.35 | 0.37 |
| January 18, 1987   | 0.46 | 0.41 | 0.35 |
| February 22, 1987  | 0.08 | 0.07 | 0.08 |
| February 26, 1987  | 0.27 | 0.27 | 0.22 |
| March 18, 1987     | 0.03 | 0.03 | 0.03 |
| March 30, 1987     | 0.20 | 0.24 | 0.26 |
| May 12, 1987       | 0.03 | 0.02 | 0.02 |
| May 21, 1987       | 0.03 | 0.03 | 0.04 |

Table A.3: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| May 25, 1987       | 0.04 | 0.03 | 0.05 |
| June 16, 1987      | 0.05 | 0.03 | 0.07 |
| June 25, 1987      | 0.21 | 0.07 | 0.09 |
| July 11, 1987      | 0.08 | 0.05 | 0.10 |
| August 22, 1987    | 0.02 | 0.02 | 0.05 |
| September 12, 1987 | 0.02 | 0.01 | 0.02 |
| November 9, 1987   | 0.03 | 0.02 | 0.03 |
| November 16, 1987  | 0.01 | 0.01 | 0.01 |
| December 14, 1987  | 0.01 | 0.01 | 0.02 |
| December 24, 1987  | 0.08 | 0.15 | 0.14 |
| January 18, 1988   | 0.54 | 0.33 | 0.33 |
| April 6, 1988      | 0.28 | 0.20 | 0.19 |
| May 4, 1988        | 0.36 | 0.43 | 0.51 |
| June 2, 1988       | 0.01 | 0.01 | 0.02 |
| June 9, 1988       | 0.01 | 0.01 | 0.01 |
| August 23, 1988    | 0.05 | 0.01 | 0.02 |
| September 16, 1988 | 0.08 | 0.03 | 0.09 |
| November 4, 1988   | 0.02 | 0.01 | 0.02 |
| November 27, 1988  | 0.01 | 0.02 | 0.02 |
| December 21, 1988  | 0.10 | 0.13 | 0.08 |
| December 24, 1988  | 0.88 | 0.70 | 0.54 |
| January 11, 1989   | 0.35 | 0.32 | 0.32 |
| February 3, 1989   | 0.15 | 0.14 | 0.13 |
| February 13, 1989  | 0.20 | 0.21 | 0.22 |
| February 20, 1989  | 0.73 | 0.60 | 0.54 |
| March 20, 1989     | 0.27 | 0.22 | 0.19 |
| April 3, 1989      | 0.07 | 0.07 | 0.06 |
| May 19, 1989       | 0.02 | 0.02 | 0.03 |
| June 12, 1989      | 1.53 | 1.07 | 1.36 |
| June 22, 1989      | 0.12 | 0.07 | 0.09 |
| July 23, 1989      | 0.33 | 0.10 | 0.21 |
| July 27, 1989      | 0.03 | 0.03 | 0.04 |
| July 31, 1989      | 0.09 | 0.14 | 0.30 |
| August 5, 1989     | 0.03 | 0.07 | 0.06 |
| August 18, 1989    | 0.17 | 0.09 | 0.15 |
| September 22, 1989 | 0.16 | 0.20 | 0.21 |
| September 30, 1989 | 0.29 | 0.32 | 0.30 |
| October 16, 1989   | 1.39 | 0.80 | 0.83 |
| November 14, 1989  | 0.21 | 0.21 | 0.18 |
| January 29, 1990   | 0.15 | 0.18 | 0.11 |
| February 3, 1990   | 0.27 | 0.28 | 0.25 |
| February 9, 1990   | 0.34 | 0.33 | 0.30 |



Table A.3: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| February 15, 1990  | 0.30 | 0.29 | 0.21 |
| March 16, 1990     | 0.21 | 0.20 | 0.14 |
| April 6, 1990      | 0.04 | 0.04 | 0.04 |
| May 26, 1990       | 0.02 | 0.01 | 0.02 |
| May 28, 1990       | 0.07 | 0.05 | 0.11 |
| June 2, 1990       | 0.20 | 0.09 | 0.13 |
| June 21, 1990      | 0.08 | 0.05 | 0.06 |
| July 30, 1990      | 0.01 | 0.01 | 0.02 |
| August 8, 1990     | 0.02 | 0.01 | 0.02 |
| August 29, 1990    | 0.02 | 0.01 | 0.01 |
| September 9, 1990  | 0.11 | 0.05 | 0.08 |
| September 12, 1990 | 0.01 | 0.01 | 0.01 |
| October 4, 1990    | 0.11 | 0.07 | 0.08 |
| December 2, 1990   | 0.10 | 0.16 | 0.13 |
| December 20, 1990  | 0.28 | 0.39 | 0.26 |
| December 27, 1990  | 0.21 | 0.29 | 0.24 |
| December 30, 1990  | 0.18 | 0.23 | 0.22 |
| January 6, 1991    | 0.13 | 0.13 | 0.10 |
| February 6, 1991   | 0.08 | 0.08 | 0.07 |
| February 13, 1991  | 0.09 | 0.11 | 0.10 |
| February 17, 1991  | 0.23 | 0.45 | 0.38 |
| March 22, 1991     | 0.12 | 0.12 | 0.10 |
| March 29, 1991     | 0.13 | 0.13 | 0.11 |
| April 15, 1991     | 0.27 | 0.26 | 0.21 |
| April 19, 1991     | 0.49 | 0.41 | 0.16 |
| May 9, 1991        | 0.04 | 0.03 | 0.03 |
| May 18, 1991       | 0.03 | 0.02 | 0.03 |
| May 27, 1991       | 0.04 | 0.03 | 0.04 |
| May 29, 1991       | 0.16 | 0.09 | 0.08 |
| June 22, 1991      | 0.22 | 0.29 | 0.26 |
| June 25, 1991      | 0.22 | 0.34 | 0.22 |
| July 10, 1991      | 0.10 | 0.06 | 0.06 |
| July 12, 1991      | 0.22 | 0.12 | 0.11 |
| August 7, 1991     | 0.22 | 0.13 | 0.21 |
| October 5, 1991    | 0.19 | 0.09 | 0.14 |
| October 15, 1991   | 0.02 | 0.02 | 0.03 |
| November 21, 1991  | 0.17 | 0.21 | 0.15 |
| December 13, 1991  | 0.10 | 0.11 | 0.10 |
| December 28, 1991  | 0.15 | 0.16 | 0.12 |
| February 24, 1992  | 0.21 | 0.22 | 0.17 |
| March 30, 1992     | 0.10 | 0.10 | 0.08 |
| May 28, 1992       | 0.02 | 0.01 | 0.03 |

Table A.3: Continued

| Date                        | WSA             | WSB             | WSC             |
|-----------------------------|-----------------|-----------------|-----------------|
| June 17, 1992               | 0.04            | 0.03            | 0.03            |
| June 24, 1992               | 0.11            | 0.05            | 0.06            |
| July 1, 1992                | 0.88            | 0.38            | 0.62            |
| July 14, 1992               | 0.05            | 0.05            | 0.06            |
| July 17, 1992               | 0.02            | 0.02            | 0.02            |
| July 22, 1992               | 0.06            | 0.07            | 0.04            |
| July 24, 1992               | 0.45            | 0.80            | 0.81            |
| August 8, 1992              | 0.25            | 0.18            | 0.09            |
| August 27, 1992             | 0.21            | 0.11            | 0.13            |
| September 4, 1992           | 0.04            | 0.02            | 0.03            |
| September 18, 1992          | 0.05            | 0.02            | 0.03            |
| December 20, 1992           | 0.13            | 0.13            | 0.12            |
| March 29, 1993              | 0.22            | 0.36            | 0.74            |
| April 25, 1993              | 0.07            | 0.05            | 0.02            |
| May 9, 1993                 | 0.21            | 0.14            | 0.18            |
| June 9, 1993                | 0.11            | 0.05            | 0.15            |
| June 21, 1993               | 0.14            | 0.05            | 0.15            |
| July 13, 1993               | 0.10            | 0.14            | 2.42            |
| July 15, 1993               | 0.02            | 0.03            | 0.11            |
| July 26, 1993               | 0.30            | 0.40            | 0.38            |
| September 2, 1993           | 0.05            | 0.14            | 0.15            |
| September 15, 1993          | 0.05            | 0.14            | 0.15            |
| October 18, 1993            | 0.05            | 0.15            | 0.16            |
| Mean $\pm$ Std. Dev. (Pre)  | 0.20 $\pm$ 0.36 | 0.15 $\pm$ 0.21 | 0.14 $\pm$ 0.23 |
| Mean $\pm$ Std. Dev. (Post) | 0.18 $\pm$ 0.34 | 0.16 $\pm$ 0.20 | 0.18 $\pm$ 0.27 |

<sup>1</sup> Final storm before treatment.

Table A.4: WSA, WSB, and WSC peak flow ( $\text{m}^3 \text{s}^{-1} \text{ha}^{-1} \times 10^{-2}$ ) (2002-2008).

| Date               | WSA  | WSB  | WSC             |
|--------------------|------|------|-----------------|
| March 17, 2002     | 0.34 | 0.46 | -- <sup>1</sup> |
| April 28, 2002     | 0.15 | 0.15 | --              |
| May 2, 2002        | 1.78 | 1.59 | --              |
| May 13, 2002       | 0.50 | 0.64 | --              |
| May 17, 2002       | 0.11 | 0.11 | --              |
| June 6, 2002       | 0.03 | 0.03 | --              |
| July 19, 2002      | 0.20 | 0.07 | --              |
| September 25, 2002 | 0.03 | 0.01 | --              |
| October 15, 2002   | 0.01 | 0.00 | --              |
| October 29, 2002   | 0.10 | 0.09 | --              |
| November 5, 2002   | 0.03 | 0.04 | --              |
| November 10, 2002  | 0.05 | 0.09 | --              |
| December 10, 2002  | 0.13 | 0.13 | --              |
| December 13, 2002  | 0.17 | 0.19 | --              |
| December 19, 2002  | 0.03 | 0.04 | --              |
| January 1, 2003    | 0.03 | 0.03 | --              |
| February 3, 2003   | 0.05 | 0.06 | --              |
| April 6, 2003      | 0.21 | 0.25 | --              |
| April 17, 2003     | 0.06 | 0.08 | --              |
| May 8, 2003        | 0.11 | 0.09 | --              |
| June 6, 2003       | 0.12 | 0.14 | --              |
| June 11, 2003      | 0.05 | 0.03 | --              |
| July 10, 2003      | 0.02 | 0.02 | --              |
| August 3, 2003     | 0.02 | 0.01 | --              |
| August 17, 2003    | 0.07 | 0.03 | --              |
| September 3, 2003  | 0.46 | 0.56 | --              |
| September 22, 2003 | 0.03 | 0.01 | --              |
| November 12, 2003  | 0.04 | 0.02 | --              |
| November 18, 2003  | 0.57 | 0.69 | --              |
| November 28, 2003  | 0.04 | 0.04 | --              |
| December 10, 2003  | 0.05 | 0.05 | --              |
| January 17, 2004   | 0.17 | 0.15 | --              |
| February 2, 2004   | 0.06 | 0.08 | --              |
| February 5, 2004   | 0.59 | 0.53 | --              |
| March 5, 2004      | 0.72 | 0.58 | --              |
| May 16, 2004       | 0.02 | 0.01 | --              |
| May 24, 2004       | 0.06 | 0.02 | --              |
| May 26, 2004       | 0.40 | 0.30 | --              |
| May 27, 2004       | 0.99 | 0.83 | --              |
| June 15, 2004      | 0.40 | 0.69 | --              |
| June 22, 2004      | 0.21 | 0.30 | --              |

Table A.4: Continued

| Date               | WSA  | WSB   | WSC   |
|--------------------|------|-------|-------|
| June 25, 2004      | 0.06 | 0.08  | --    |
| July 6, 2004       | 0.05 | 0.05  | --    |
| July 26, 2004      | 0.04 | 0.05  | --    |
| July 27, 2004      | 0.07 | 0.09  | --    |
| July 31, 2004      | 0.03 | 0.02  | --    |
| August 5, 2004     | 0.11 | 0.08  | --    |
| September 7, 2004  | 0.26 | 0.29  | --    |
| September 16, 2004 | 0.97 | 1.28  | --    |
| October 13, 2004   | 0.17 | 0.09  | --    |
| October 18, 2004   | 0.54 | 0.61  | --    |
| October 27, 2004   | 0.33 | 0.40  | --    |
| November 4, 2004   | 0.98 | 0.42  | --    |
| November 11, 2004  | 0.03 | 0.03  | --    |
| November 30, 2004  | 0.36 | 0.51  | --    |
| December 9, 2004   | 0.48 | 0.52  | --    |
| December 23, 2004  | 0.03 | 0.03  | --    |
| April 29, 2005     | 1.87 | 1.67  | 1.94  |
| May 19, 2005       | 0.09 | 0.12  | 0.08  |
| May 22, 2005       | 0.07 | 0.11  | 0.07  |
| June 3, 2005       | 0.34 | 0.22  | 0.08  |
| June 20, 2005      | 0.10 | 0.11  | 0.08  |
| July 1, 2005       | 0.07 | 0.05  | 0.04  |
| July 7, 2005       | 0.01 | 0.00  | 0.02  |
| July 17, 2005      | 0.72 | 0.81  | 0.96  |
| July 18, 2005      | 0.20 | 0.14  | 0.19  |
| July 27, 2005      | 0.05 | 0.01  | 0.03  |
| August 5, 2005     | 0.11 | 0.06  | 0.07  |
| August 16, 2005    | 0.02 | 0.003 | 0.005 |
| August 19, 2005    | 0.05 | 0.01  | 0.01  |
| December 3, 2005   | 0.01 | 0.004 | 0.02  |
| December 15, 2005  | 0.03 | 0.01  | 0.03  |
| January 17, 2006   | 0.09 | 0.10  | 0.10  |
| January 23, 2006   | 0.81 | 0.49  | 0.40  |
| March 13, 2006     | 0.11 | 0.12  | 0.10  |
| May 2, 2006        | 0.04 | 0.02  | 0.07  |
| May 25, 2006       | 0.03 | 0.02  | 0.06  |
| May 31, 2006       | 0.33 | 0.15  | 0.17  |
| July 5, 2006       | 0.16 | 0.17  | 0.21  |
| September 23, 2006 | 0.04 | 0.03  | 0.04  |
| September 23, 2006 | 0.03 | 0.03  | 0.04  |
| September 30, 2006 | 0.02 | 0.01  | 0.01  |
| October 26, 2006   | 0.31 | 0.33  | 0.44  |

Table A.4: Continued

| Date                         | WSA             | WSB             | WSC             |
|------------------------------|-----------------|-----------------|-----------------|
| March 28, 2007               | 0.02            | 0.01            | 0.03            |
| May 16, 2007                 | 0.04            | 0.02            | 0.08            |
| June 1, 2007                 | 0.20            | 0.06            | 0.10            |
| June 5, 2007                 | 0.14            | 0.06            | 0.24            |
| June 18, 2007                | 0.03            | 0.004           | 0.05            |
| June 29, 2007                | 0.03            | 0.01            | 0.06            |
| August 3, 2007               | 0.03            | 0.00            | 0.05            |
| November 14, 2007            | 0.04            | 0.01            | 0.04            |
| November 26, 2007            | 0.03            | 0.01            | 0.06            |
| December 10, 2007            | 0.03            | 0.04            | 0.05            |
| December 13, 2007            | 0.05            | 0.07            | 0.09            |
| December 20, 2007            | 0.01            | 0.01            | 0.02            |
| January 29, 2008             | 0.07            | 0.09            | 0.13            |
| February 6, 2008             | 0.08            | 0.10            | 0.13            |
| May 11, 2008                 | 0.01            | 0.01            | 0.03            |
| June 3, 2008                 | 0.05            | 0.02            | 0.08            |
| August 26, 2008              | 0.01            | 0.00            | 0.01            |
| November 14, 2008            | 0.01            | 0.03            | 0.02            |
| December 10, 2008            | 0.03            | 0.06            | 0.06            |
| December 10, 2008            | 0.04            | 0.08            | 0.07            |
| December 16, 2008            | 0.05            | 0.07            | 0.06            |
| Mean $\pm$ Std. Dev. (02-04) | 0.24 $\pm$ 0.33 | 0.24 $\pm$ 0.32 | - -             |
| Mean $\pm$ Std. Dev. (05-08) | 0.14 $\pm$ 0.30 | 0.12 $\pm$ 0.27 | 0.14 $\pm$ 0.31 |

<sup>1</sup>No data available.

Table A.5: WSA, WSB, and WSC discharge volume as % of rainfall (1982-1993).

| Date                         | WSA  | WSB  | WSC  |
|------------------------------|------|------|------|
| February 8, 1982             | 76.4 | 52.2 | 54.9 |
| February 16, 1982            | 72.5 | 52.6 | 90.4 |
| March 15, 1982               | 77.9 | 40.4 | 75.6 |
| March 31, 1982               | 37.6 | 35.6 | 57.2 |
| May 21, 1982                 | 2.4  | 2.3  | 2.9  |
| May 28, 1982                 | 2.0  | 2.8  | 4.2  |
| June 4, 1982                 | 3.9  | 4.6  | 6.4  |
| July 28, 1982                | 0.5  | 0.5  | 0.3  |
| August 5, 1982               | 1.6  | 1.7  | 1.3  |
| September 13, 1982           | 2.1  | 1.6  | 0.8  |
| September 25, 1982           | 1.5  | 1.3  | 1.1  |
| November 20, 1982            | 7.5  | 5.6  | 3.3  |
| November 30, 1982            | 26.9 | 21.9 | 18.7 |
| December 5, 1982             | 20.6 | 17.7 | 16.7 |
| December 15, 1982            | 54.6 | 47.4 | 80.1 |
| January 21, 1983             | 42.5 | 44.1 | 37.3 |
| February 10, 1983            | 56.8 | 56.1 | 59.7 |
| April 14, 1983               | 49.2 | 55.8 | 51.2 |
| May 3, 1983                  | 51.5 | 50.6 | 50.6 |
| May 13, 1983                 | 44.1 | 58.7 | 57.6 |
| May 22, 1983                 | 33.0 | 39.0 | 35.6 |
| June 3, 1983                 | 3.9  | 2.7  | 2.8  |
| June 4, 1983                 | 37.2 | 44.4 | 46.2 |
| July 3, 1983                 | 0.8  | 0.5  | 0.6  |
| July 5, 1983                 | 1.6  | 2.3  | 1.5  |
| July 18, 1983                | 0.8  | 0.7  | 0.6  |
| August 2, 1983               | 0.5  | 0.3  | 0.3  |
| August 11, 1983              | 1.0  | 1.1  | 1.0  |
| August 27, 1983 <sup>1</sup> | 0.1  | 0.2  | 0.1  |
| November 14, 1983            | 4.5  | 7.5  | 5.3  |
| December 27, 1983            | 16.0 | 12.3 | 9.6  |
| February 27, 1984            | 43.8 | 37.6 | 39.2 |
| March 20, 1984               | 75.8 | 71.3 | 84.3 |
| March 28, 1984               | 58.6 | 53.0 | 63.9 |
| April 4, 1984                | 57.7 | 56.6 | 60.6 |
| April 9, 1984                | 57.6 | 56.4 | 74.2 |
| April 21, 1984               | 64.5 | 69.2 | 75.1 |
| May 4, 1984                  | 74.1 | 79.3 | 76.1 |
| May 6, 1984                  | 62.5 | 75.6 | 96.0 |
| May 23, 1984                 | 2.3  | 3.6  | 3.0  |
| May 28, 1984                 | 5.5  | 53.2 | 63.1 |

Table A.5: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| July 26, 1984      | 0.7  | 1.2  | 2.4  |
| July 27, 1984      | 1.6  | 30.1 | 39.8 |
| July 30, 1984      | 0.5  | 8.6  | 17.0 |
| August 22, 1984    | 1.7  | 6.8  | 13.5 |
| November 4, 1984   | 4.2  | 49.7 | 56.4 |
| November 18, 1984  | 19.8 | 55.1 | 62.3 |
| November 28, 1984  | 25.3 | 52.7 | 54.0 |
| December 20, 1984  | 54.9 | 84.8 | 90.4 |
| December 24, 1984  | 52.4 | 76.1 | 97.0 |
| January 3, 1985    | 43.0 | 69.0 | 82.1 |
| February 11, 1985  | 34.6 | 43.0 | 42.1 |
| May 15, 1985       | 1.2  | 2.7  | 5.5  |
| June 5, 1985       | 3.1  | 11.2 | 10.8 |
| June 10, 1985      | 1.4  | 4.1  | 3.7  |
| June 11, 1985      | 16.4 | 57.1 | 52.1 |
| July 10, 1985      | 1.2  | 3.0  | 4.3  |
| July 30, 1985      | 0.5  | 0.7  | 0.7  |
| August 1, 1985     | 1.8  | 5.7  | 5.6  |
| August 17, 1985    | 2.3  | 10.2 | 8.9  |
| August 25, 1985    | 1.8  | 2.7  | 12.8 |
| August 30, 1985    | 2.3  | 14.8 | 18.9 |
| September 26, 1985 | 0.5  | 3.3  | 5.4  |
| November 2, 1985   | 18.4 | 55.0 | 57.7 |
| December 12, 1985  | 48.6 | 62.6 | 63.2 |
| February 2, 1986   | 72.4 | 85.4 | 94.7 |
| April 28, 1986     | 6.1  | 7.6  | 15.6 |
| May 11, 1986       | 3.2  | 9.8  | 7.1  |
| July 2, 1986       | 1.7  | 5.0  | 5.2  |
| July 20, 1986      | 1.5  | 1.8  | 8.0  |
| July 26, 1986      | 1.1  | 1.8  | 6.0  |
| October 1, 1986    | 3.8  | 4.2  | 7.6  |
| November 5, 1986   | 5.9  | 15.0 | 21.6 |
| November 8, 1986   | 26.3 | 45.7 | 50.5 |
| November 10, 1986  | 21.2 | 38.3 | 53.2 |
| December 8, 1986   | 64.4 | 69.9 | 90.4 |
| January 18, 1987   | 51.9 | 58.0 | 74.6 |
| February 22, 1987  | 61.9 | 69.2 | 69.3 |
| February 26, 1987  | 74.9 | 73.4 | 81.3 |
| March 18, 1987     | 85.3 | 83.1 | 69.4 |
| March 30, 1987     | 65.4 | 69.2 | 75.2 |
| May 12, 1987       | 1.6  | 3.0  | 5.8  |
| May 21, 1987       | 2.5  | 4.6  | 4.8  |

Table A.5: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| May 25, 1987       | 3.6  | 5.9  | 8.1  |
| June 16, 1987      | 3.3  | 3.5  | 5.1  |
| June 25, 1987      | 3.3  | 4.6  | 5.3  |
| July 11, 1987      | 6.6  | 15.7 | 24.1 |
| August 22, 1987    | 1.1  | 2.3  | 3.4  |
| September 12, 1987 | 2.6  | 3.0  | 4.7  |
| November 9, 1987   | 2.8  | 2.5  | 4.9  |
| November 16, 1987  | 2.3  | 6.2  | 4.9  |
| December 14, 1987  | 4.1  | 9.5  | 9.4  |
| December 24, 1987  | 12.4 | 15.5 | 26.8 |
| January 18, 1988   | 55.1 | 43.2 | 61.6 |
| April 6, 1988      | 81.1 | 63.6 | 87.6 |
| May 4, 1988        | 58.3 | 27.8 | 49.5 |
| June 2, 1988       | 1.2  | 1.3  | 7.1  |
| June 9, 1988       | 1.4  | 1.6  | 2.1  |
| August 23, 1988    | 0.8  | 1.3  | 4.5  |
| September 16, 1988 | 1.5  | 1.6  | 3.2  |
| November 4, 1988   | 5.3  | 7.8  | 9.9  |
| November 27, 1988  | 17.8 | 36.3 | 35.7 |
| December 21, 1988  | 17.7 | 33.1 | 31.0 |
| December 24, 1988  | 70.2 | 61.9 | 66.2 |
| January 11, 1989   | 64.7 | 67.5 | 81.6 |
| February 3, 1989   | 48.7 | 58.8 | 73.3 |
| February 13, 1989  | 61.5 | 66.0 | 73.3 |
| February 20, 1989  | 86.4 | 69.2 | 74.8 |
| March 20, 1989     | 49.2 | 58.5 | 62.7 |
| April 3, 1989      | 57.1 | 78.6 | 79.7 |
| May 19, 1989       | 5.9  | 5.9  | 5.4  |
| June 12, 1989      | 51.0 | 47.4 | 52.3 |
| June 22, 1989      | 16.7 | 22.6 | 21.6 |
| July 23, 1989      | 7.2  | 6.5  | 6.2  |
| July 27, 1989      | 1.5  | 2.2  | 1.9  |
| July 31, 1989      | 8.1  | 21.7 | 30.2 |
| August 5, 1989     | 13.0 | 22.8 | 21.3 |
| August 18, 1989    | 1.5  | 2.4  | 3.6  |
| September 22, 1989 | 24.2 | 39.9 | 57.8 |
| September 30, 1989 | 40.5 | 49.4 | 54.7 |
| October 16, 1989   | 45.3 | 49.0 | 71.2 |
| November 14, 1989  | 54.1 | 71.9 | 78.1 |
| January 29, 1990   | 56.4 | 70.4 | 66.4 |
| February 3, 1990   | 73.6 | 80.9 | 87.9 |
| February 9, 1990   | 55.8 | 73.5 | 77.6 |



Table A.5: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| February 15, 1990  | 56.0 | 59.2 | 63.9 |
| March 16, 1990     | 54.9 | 71.4 | 55.3 |
| April 6, 1990      | 50.0 | 49.1 | 55.0 |
| May 26, 1990       | 1.6  | 1.7  | 1.8  |
| May 28, 1990       | 12.5 | 14.2 | 16.1 |
| June 2, 1990       | 16.3 | 16.7 | 20.7 |
| June 21, 1990      | 3.3  | 2.0  | 3.7  |
| July 30, 1990      | 0.6  | 1.3  | 2.6  |
| August 8, 1990     | 1.0  | 2.3  | 3.4  |
| August 29, 1990    | 1.1  | 1.4  | 2.4  |
| September 9, 1990  | 3.0  | 1.4  | 2.3  |
| September 12, 1990 | 0.9  | 1.6  | 2.1  |
| October 4, 1990    | 1.2  | 4.3  | 4.5  |
| December 2, 1990   | 34.8 | 55.3 | 68.1 |
| December 20, 1990  | 54.6 | 75.3 | 64.6 |
| December 27, 1990  | 24.7 | 33.7 | 35.7 |
| December 30, 1990  | 31.8 | 42.0 | 54.2 |
| January 6, 1991    | 85.8 | 96.5 | 91.1 |
| February 6, 1991   | 73.4 | 80.2 | 68.3 |
| February 13, 1991  | 46.5 | 51.3 | 52.9 |
| February 17, 1991  | 61.9 | 76.8 | 97.6 |
| March 22, 1991     | 55.6 | 60.2 | 58.9 |
| March 29, 1991     | 81.0 | 76.4 | 75.0 |
| April 15, 1991     | 66.9 | 78.7 | 76.7 |
| April 19, 1991     | 63.6 | 62.9 | 60.1 |
| May 9, 1991        | 7.0  | 5.4  | 1.8  |
| May 18, 1991       | 2.2  | 7.8  | 2.3  |
| May 27, 1991       | 2.7  | 4.0  | 3.0  |
| May 29, 1991       | 7.1  | 8.8  | 5.7  |
| June 22, 1991      | 10.7 | 25.4 | 19.1 |
| June 25, 1991      | 40.2 | 42.8 | 37.3 |
| July 10, 1991      | 1.9  | 1.7  | 1.4  |
| July 12, 1991      | 5.5  | 9.2  | 9.0  |
| August 7, 1991     | 4.6  | 5.5  | 4.0  |
| October 5, 1991    | 2.3  | 3.3  | 1.9  |
| October 15, 1991   | 5.5  | 14.4 | 4.7  |
| November 21, 1991  | 15.8 | 18.1 | 14.8 |
| December 13, 1991  | 50.3 | 53.2 | 71.4 |
| December 28, 1991  | 62.2 | 64.0 | 63.5 |
| February 24, 1992  | 65.1 | 76.4 | 74.3 |
| March 30, 1992     | 61.1 | 74.7 | 70.8 |
| May 28, 1992       | 4.6  | 7.0  | 6.6  |

Table A.5: Continued

| Date                        | WSA             | WSB             | WSC             |
|-----------------------------|-----------------|-----------------|-----------------|
| June 17, 1992               | 8.1             | 13.2            | 9.4             |
| June 24, 1992               | 1.9             | 5.2             | 3.8             |
| July 1, 1992                | 44.3            | 31.3            | 32.3            |
| July 14, 1992               | 0.8             | 0.8             | 2.7             |
| July 17, 1992               | 2.6             | 4.9             | 3.3             |
| July 22, 1992               | 2.7             | 3.4             | 2.7             |
| July 24, 1992               | 15.2            | 24.1            | 24.3            |
| August 8, 1992              | 10.4            | 11.0            | 14.5            |
| August 27, 1992             | 9.5             | 8.6             | 11.6            |
| September 4, 1992           | 2.3             | 5.0             | 8.2             |
| September 18, 1992          | 3.4             | 2.8             | 18.5            |
| December 20, 1992           | 37.9            | 37.3            | 27.5            |
| March 29, 1993              | 62.1            | 79.7            | 78.9            |
| April 25, 1993              | 89.5            | 53.4            | 15.8            |
| May 9, 1993                 | 9.4             | 10.1            | 3.9             |
| June 9, 1993                | 1.8             | 3.3             | 4.4             |
| June 21, 1993               | 2.2             | 3.8             | 3.3             |
| July 13, 1993               | 1.5             | 2.7             | 8.8             |
| July 15, 1993               | 8.3             | 11.6            | 34.5            |
| July 26, 1993               | 1.9             | 2.8             | 2.6             |
| September 2, 1993           | 3.1             | 12.1            | 13.0            |
| September 15, 1993          | 3.7             | 7.8             | 8.3             |
| October 18, 1993            | 23.7            | 60.5            | 64.2            |
| Mean $\pm$ Std. Dev. (Pre)  | 23.9 $\pm$ 26.5 | 21.7 $\pm$ 22.8 | 25.5 $\pm$ 29.2 |
| Mean $\pm$ Std. Dev. (Post) | 26.0 $\pm$ 27.5 | 31.7 $\pm$ 29.1 | 34.7 $\pm$ 31.2 |

<sup>1</sup> Final storm before treatment.

Table A.6: WSA, WSB, and WSC discharge volume as % of rainfall (2002-2008).

| Date               | WSA  | WSB  | WSC             |
|--------------------|------|------|-----------------|
| March 17, 2002     | 68.9 | 86.1 | -- <sup>1</sup> |
| April 28, 2002     | 75.7 | 90.8 | --              |
| May 2, 2002        | 75.9 | 86.3 | --              |
| May 13, 2002       | 40.2 | 54.9 | --              |
| May 17, 2002       | 23.0 | 44.3 | --              |
| June 6, 2002       | 4.2  | 7.0  | --              |
| July 19, 2002      | 7.4  | 5.2  | --              |
| September 25, 2002 | 4.0  | 2.3  | --              |
| October 15, 2002   | 5.1  | 2.2  | --              |
| October 29, 2002   | 23.1 | 27.3 | --              |
| November 5, 2002   | 17.6 | 23.0 | --              |
| November 10, 2002  | 17.4 | 65.8 | --              |
| December 10, 2002  | 67.0 | 75.9 | --              |
| December 13, 2002  | 69.0 | 79.0 | --              |
| December 19, 2002  | 36.2 | 36.5 | --              |
| January 1, 2003    | 76.7 | 69.3 | --              |
| February 3, 2003   | 86.6 | 86.8 | --              |
| April 6, 2003      | 46.9 | 52.7 | --              |
| April 17, 2003     | 41.7 | 44.0 | --              |
| May 8, 2003        | 20.3 | 40.4 | --              |
| June 6, 2003       | 17.9 | 27.8 | --              |
| June 11, 2003      | 1.8  | 6.9  | --              |
| July 10, 2003      | 1.9  | 2.4  | --              |
| August 3, 2003     | 1.3  | 2.5  | --              |
| August 17, 2003    | 2.6  | 7.9  | --              |
| September 3, 2003  | 16.8 | 17.9 | --              |
| September 22, 2003 | 2.4  | 0.9  | --              |
| November 12, 2003  | 12.6 | 15.4 | --              |
| November 18, 2003  | 43.6 | 46.7 | --              |
| November 28, 2003  | 48.6 | 47.9 | --              |
| December 10, 2003  | 53.0 | 65.6 | --              |
| January 17, 2004   | 58.2 | 65.2 | --              |
| February 2, 2004   | 51.8 | 65.5 | --              |
| February 5, 2004   | 35.5 | 20.9 | --              |
| March 5, 2004      | 47.2 | 56.9 | --              |
| May 16, 2004       | 3.3  | 2.2  | --              |
| May 24, 2004       | 1.8  | 2.6  | --              |
| May 26, 2004       | 23.3 | 22.9 | --              |
| May 27, 2004       | 57.7 | 71.0 | --              |
| June 15, 2004      | 39.9 | 77.8 | --              |
| June 22, 2004      | 25.9 | 33.7 | --              |

Table A.6: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| June 25, 2004      | 39.1 | 54.0 | --   |
| July 6, 2004       | 5.5  | 6.5  | --   |
| July 26, 2004      | 0.6  | 1.5  | --   |
| July 27, 2004      | 16.1 | 10.1 | --   |
| July 31, 2004      | 5.8  | 2.2  | --   |
| August 5, 2004     | 4.1  | 5.2  | --   |
| September 7, 2004  | 13.8 | 13.8 | --   |
| September 16, 2004 | 44.0 | 52.0 | --   |
| October 13, 2004   | 6.9  | 8.8  | --   |
| October 18, 2004   | 36.2 | 44.7 | --   |
| October 27, 2004   | 62.2 | 61.4 | --   |
| November 4, 2004   | 60.4 | 54.3 | --   |
| November 11, 2004  | 33.4 | 24.2 | --   |
| November 30, 2004  | 77.4 | 85.8 | --   |
| December 9, 2004   | 91.3 | 98.9 | --   |
| December 23, 2004  | 14.5 | 25.0 | --   |
| April 29, 2005     | 61.8 | 80.2 | 99.6 |
| May 19, 2005       | 10.4 | 16.8 | 9.4  |
| May 22, 2005       | 36.8 | 64.8 | 19.5 |
| June 3, 2005       | 77.5 | 90.0 | 31.3 |
| June 20, 2005      | 2.1  | 9.8  | 5.8  |
| July 1, 2005       | 3.1  | 1.9  | 1.7  |
| July 7, 2005       | 6.0  | 0.7  | 21.2 |
| July 17, 2005      | 11.6 | 12.7 | 7.5  |
| July 18, 2005      | 32.1 | 27.3 | 31.6 |
| July 27, 2005      | 3.4  | 0.8  | 6.5  |
| August 5, 2005     | 2.4  | 1.3  | 1.3  |
| August 16, 2005    | 1.7  | 0.5  | 0.6  |
| August 19, 2005    | 3.2  | 1.0  | 1.9  |
| December 3, 2005   | 2.7  | 2.6  | 18.2 |
| December 15, 2005  | 12.1 | 6.4  | 12.8 |
| January 17, 2006   | 21.4 | 34.2 | 25.2 |
| January 23, 2006   | 40.0 | 59.0 | 86.1 |
| March 13, 2006     | 59.5 | 81.9 | 85.4 |
| May 2, 2006        | 5.9  | 10.3 | 42.2 |
| May 25, 2006       | 7.9  | 6.3  | 19.9 |
| May 31, 2006       | 16.6 | 32.4 | 46.6 |
| July 5, 2006       | 20.7 | 32.2 | 45.7 |
| September 23, 2006 | 2.1  | 3.6  | 3.9  |
| September 23, 2006 | 7.7  | 17.7 | 18.0 |
| September 30, 2006 | 3.4  | 8.3  | 7.9  |
| October 26, 2006   | 20.7 | 31.5 | 43.5 |

Table A.6: Continued

| Date                         | WSA             | WSB             | WSC             |
|------------------------------|-----------------|-----------------|-----------------|
| March 28, 2007               | 29.1            | 6.4             | 14.5            |
| May 16, 2007                 | 2.2             | 4.6             | 14.0            |
| June 1, 2007                 | 2.1             | 1.8             | 12.7            |
| June 5, 2007                 | 6.0             | 5.5             | 11.3            |
| June 18, 2007                | 0.6             | 0.1             | 1.4             |
| June 29, 2007                | 5.2             | 1.4             | 6.8             |
| August 3, 2007               | 1.6             | 0.5             | 6.5             |
| November 14, 2007            | 4.7             | 2.3             | 6.5             |
| November 26, 2007            | 3.5             | 6.0             | 18.2            |
| December 10, 2007            | 4.6             | 7.4             | 14.0            |
| December 13, 2007            | 16.3            | 29.8            | 34.6            |
| December 20, 2007            | 3.2             | 17.9            | 27.7            |
| January 29, 2008             | 30.1            | 50.5            | 59.5            |
| February 6, 2008             | 49.9            | 87.4            | 91.9            |
| May 11, 2008                 | 3.9             | 8.6             | 10.0            |
| June 3, 2008                 | 4.5             | 7.2             | 9.5             |
| August 26, 2008              | 1.2             | 0.6             | 1.5             |
| November 14, 2008            | 2.5             | 10.6            | 5.8             |
| December 10, 2008            | 2.1             | 8.9             | 7.0             |
| December 10, 2008            | 7.7             | 33.7            | 16.2            |
| December 16, 2008            | 19.6            | 27.6            | 40.7            |
| Mean $\pm$ Std. Dev. (02-04) | 32.7 $\pm$ 26.4 | 38.4 $\pm$ 29.9 | --              |
| Mean $\pm$ Std. Dev. (05-08) | 14.3 $\pm$ 18.1 | 20.3 $\pm$ 25.3 | 23.5 $\pm$ 25.2 |

<sup>1</sup>No data available.

Table A.7: WSA, WSB, and WSC time to peak (2002-2008).

| Date               | WSA  | WSB  | WSC             |
|--------------------|------|------|-----------------|
| March 17, 2002     | 8.3  | 6.9  | -- <sup>1</sup> |
| April 28, 2002     | 1.7  | 4.6  | --              |
| May 2, 2002        | 3.9  | 5.4  | --              |
| May 13, 2002       | 4.3  | 3.2  | --              |
| May 17, 2002       | 1.3  | 3.3  | --              |
| June 6, 2002       | 9.0  | 2.5  | --              |
| July 19, 2002      | 2.3  | 7.6  | --              |
| September 25, 2002 | 16.3 | 16.1 | --              |
| October 15, 2002   | 1.5  | 13.0 | --              |
| October 29, 2002   | 2.5  | 2.3  | --              |
| November 5, 2002   | 17.0 | 12.2 | --              |
| November 10, 2002  | 17.0 | 23.5 | --              |
| December 10, 2002  | 20.5 | 17.4 | --              |
| December 13, 2002  | 14.0 | 14.7 | --              |
| December 19, 2002  | 21.2 | 8.0  | --              |
| January 1, 2003    | 2.8  | 18.5 | --              |
| February 3, 2003   | 10.0 | 6.2  | --              |
| April 6, 2003      | 15.5 | 17.7 | --              |
| April 17, 2003     | 17.0 | 6.0  | --              |
| May 8, 2003        | 2.1  | 2.5  | --              |
| June 6, 2003       | 6.4  | 6.5  | --              |
| June 11, 2003      | 0.5  | 0.5  | --              |
| July 10, 2003      | 1.0  | 1.9  | --              |
| August 3, 2003     | 1.0  | 9.5  | --              |
| August 17, 2003    | 2.0  | 0.5  | --              |
| September 3, 2003  | 7.5  | 9.6  | --              |
| September 22, 2003 | 2.5  | 4.5  | --              |
| November 12, 2003  | 0.8  | 7.3  | --              |
| November 18, 2003  | 10.0 | 9.9  | --              |
| November 28, 2003  | 23.0 | 17.5 | --              |
| December 10, 2003  | 1.5  | 19.7 | --              |
| January 17, 2004   | 21.5 | 14.8 | --              |
| February 2, 2004   | 13.8 | 14.8 | --              |
| February 5, 2004   | 8.4  | 6.0  | --              |
| March 5, 2004      | 6.5  | 7.0  | --              |
| May 16, 2004       | 2.4  | 4.1  | --              |
| May 24, 2004       | 0.8  | 1.5  | --              |
| May 26, 2004       | 1.9  | 6.0  | --              |
| May 27, 2004       | 1.1  | 3.0  | --              |
| June 15, 2004      | 2.3  | 2.0  | --              |
| June 22, 2004      | 0.5  | 0.5  | --              |

Table A.7: Continued

| Date               | WSA  | WSB  | WSC  |
|--------------------|------|------|------|
| June 25, 2004      | 8.5  | 8.5  | --   |
| July 6, 2004       | 0.8  | 0.6  | --   |
| July 26, 2004      | 1.0  | 1.0  | --   |
| July 27, 2004      | 1.5  | 2.0  | --   |
| July 31, 2004      | 0.5  | 0.4  | --   |
| August 5, 2004     | 3.5  | 3.3  | --   |
| September 7, 2004  | 14.7 | 14.7 | --   |
| September 16, 2004 | 14.6 | 14.9 | --   |
| October 13, 2004   | 0.6  | 1.0  | --   |
| October 18, 2004   | 26.3 | 4.8  | --   |
| October 27, 2004   | 4.8  | 4.3  | --   |
| November 4, 2004   | 6.8  | 11.0 | --   |
| November 11, 2004  | 3.8  | 3.9  | --   |
| November 30, 2004  | 29.8 | 27.9 | --   |
| December 9, 2004   | 6.2  | 7.4  | --   |
| December 23, 2004  | 3.8  | 4.1  | --   |
| April 29, 2005     | 9.9  | 12.2 | 12.4 |
| May 19, 2005       | 21.8 | 22.0 | 22.0 |
| May 22, 2005       | 1.3  | 1.4  | 1.4  |
| June 3, 2005       | 0.6  | 0.7  | 0.5  |
| June 20, 2005      | 0.9  | 0.8  | 1.4  |
| July 1, 2005       | 2.8  | 1.0  | 3.0  |
| July 7, 2005       | 12.5 | 14.5 | 13.1 |
| July 17, 2005      | 0.6  | 0.5  | 0.8  |
| July 18, 2005      | 0.8  | 2.6  | 2.4  |
| July 27, 2005      | 0.6  | 0.4  | 0.5  |
| August 5, 2005     | 1.0  | 1.0  | 0.8  |
| August 16, 2005    | 0.8  | 0.8  | 0.8  |
| August 19, 2005    | 0.4  | 0.2  | 0.2  |
| December 3, 2005   | 5.7  | 5.0  | 6.0  |
| December 15, 2005  | 6.3  | 7.2  | 11.8 |
| January 17, 2006   | 17.2 | 16.7 | 18.4 |
| January 23, 2006   | 8.0  | 4.1  | 7.9  |
| March 13, 2006     | 8.3  | 8.6  | 6.0  |
| May 2, 2006        | 0.8  | 0.9  | 0.6  |
| May 25, 2006       | 1.5  | 2.3  | 1.6  |
| May 31, 2006       | 0.5  | 0.8  | 1.5  |
| July 5, 2006       | 7.6  | 6.6  | 7.7  |
| September 23, 2006 | 2.6  | 9.8  | 9.8  |
| September 23, 2006 | 1.2  | 3.6  | 3.6  |
| September 30, 2006 | 1.2  | 8.5  | 8.3  |
| October 26, 2006   | 16.9 | 23.1 | 16.4 |

Table A.7: Continued

| Date                         | WSA           | WSB           | WSC           |
|------------------------------|---------------|---------------|---------------|
| March 28, 2007               | 1.6           | 13.5          | 0.6           |
| May 16, 2007                 | 1.3           | 1.3           | 0.6           |
| June 1, 2007                 | 0.5           | 0.8           | 0.6           |
| June 5, 2007                 | 2.6           | 0.9           | 1.3           |
| June 18, 2007                | 0.5           | 0.1           | 0.5           |
| June 29, 2007                | 1.4           | 0.5           | 0.6           |
| August 3, 2007               | 0.2           | 0.8           | 0.2           |
| November 14, 2007            | 1.5           | 3.3           | 4.2           |
| November 26, 2007            | 1.5           | 3.0           | 7.1           |
| December 10, 2007            | 18.0          | 16.3          | 16.0          |
| December 13, 2007            | 4.8           | 4.5           | 4.8           |
| December 20, 2007            | 2.3           | 18.0          | 6.5           |
| January 29, 2008             | 10.0          | 6.8           | 10.0          |
| February 6, 2008             | 11.6          | 11.3          | 15.0          |
| May 11, 2008                 | 6.5           | 11.0          | 19.2          |
| June 3, 2008                 | 0.5           | 0.5           | 0.5           |
| August 26, 2008              | 0.9           | 9.1           | 0.6           |
| November 14, 2008            | 27.1          | 6.5           | 6.5           |
| December 10, 2008            | 2.1           | 8.0           | 6.8           |
| December 10, 2008            | 4.3           | 6.3           | 9.6           |
| December 16, 2008            | 7.7           | 4.9           | 5.6           |
| Mean $\pm$ Std. Dev. (02-04) | 7.5 $\pm$ 7.6 | 7.9 $\pm$ 6.5 | --            |
| Mean $\pm$ Std. Dev. (05-08) | 5.1 $\pm$ 6.3 | 6.0 $\pm$ 6.1 | 5.9 $\pm$ 6.0 |

<sup>1</sup>No data available.



Table A.8: Curve numbers for WSA, WSB, and WSC (1982-1993).<sup>1</sup>

| Date                         | CN ( $\lambda=0.2$ ) |     |     | CN ( $\lambda=0.05$ ) |     |     |
|------------------------------|----------------------|-----|-----|-----------------------|-----|-----|
|                              | WSA                  | WSB | WSC | WSA                   | WSB | WSC |
| February 8, 1982             | 96                   | 90  | 90  | 95                    | 86  | 87  |
| February 16, 1982            | 97                   | 94  | 99  | 96                    | 92  | 99  |
| March 15, 1982               | 97                   | 89  | 97  | 97                    | 86  | 96  |
| March 31, 1982               | 92                   | 92  | 96  | 90                    | 89  | 95  |
| May 21, 1982                 | 70                   | 70  | 71  | 59                    | 58  | 60  |
| May 28, 1982                 | 81                   | 82  | 83  | 74                    | 75  | 77  |
| June 4, 1982                 | 83                   | 84  | 85  | 77                    | 78  | 80  |
| July 28, 1982                | 62                   | 62  | 61  | 48                    | 48  | 47  |
| August 5, 1982               | 64                   | 64  | 63  | 51                    | 51  | 50  |
| September 13, 1982           | 53                   | 52  | 50  | 38                    | 37  | 35  |
| September 25, 1982           | 81                   | 81  | 80  | 74                    | 74  | 73  |
| November 20, 1982            | 82                   | 81  | 78  | 75                    | 73  | 70  |
| November 30, 1982            | 94                   | 93  | 92  | 92                    | 91  | 90  |
| December 5, 1982             | 91                   | 91  | 90  | 89                    | 88  | 87  |
| December 15, 1982            | 97                   | 96  | 99  | 97                    | 96  | 99  |
| January 21, 1983             | 92                   | 92  | 91  | 90                    | 90  | 88  |
| February 10, 1983            | 96                   | 96  | 96  | 95                    | 95  | 95  |
| April 14, 1983               | 94                   | 95  | 95  | 93                    | 94  | 93  |
| May 3, 1983                  | 92                   | 92  | 92  | 90                    | 90  | 90  |
| May 13, 1983                 | 78                   | 85  | 84  | 69                    | 79  | 79  |
| May 22, 1983                 | 85                   | 87  | 86  | 79                    | 83  | 81  |
| June 3, 1983                 | 74                   | 73  | 73  | 64                    | 62  | 62  |
| June 4, 1983                 | 95                   | 96  | 96  | 94                    | 95  | 95  |
| July 3, 1983                 | 65                   | 64  | 64  | 52                    | 51  | 51  |
| July 5, 1983                 | 84                   | 85  | 84  | 78                    | 79  | 78  |
| July 18, 1983                | 84                   | 83  | 83  | 78                    | 77  | 77  |
| August 2, 1983               | 70                   | 69  | 69  | 59                    | 58  | 58  |
| August 11, 1983              | 66                   | 66  | 66  | 54                    | 54  | 54  |
| August 27, 1983 <sup>2</sup> | 80                   | 80  | 80  | 72                    | 73  | 72  |
| November 14, 1983            | 74                   | 76  | 74  | 63                    | 67  | 64  |
| December 27, 1983            | 89                   | 88  | 87  | 86                    | 84  | 82  |
| February 27, 1984            | 89                   | 87  | 87  | 85                    | 82  | 83  |
| March 20, 1984               | 97                   | 97  | 98  | 97                    | 96  | 98  |
| March 28, 1984               | 96                   | 95  | 96  | 95                    | 94  | 96  |
| April 4, 1984                | 95                   | 95  | 95  | 94                    | 94  | 94  |
| April 9, 1984                | 92                   | 92  | 96  | 90                    | 90  | 95  |
| April 21, 1984               | 94                   | 95  | 96  | 92                    | 93  | 95  |
| May 4, 1984                  | 98                   | 98  | 98  | 97                    | 98  | 98  |
| May 6, 1984                  | 81                   | 88  | 98  | 75                    | 85  | 98  |
| May 23, 1984                 | 86                   | 87  | 87  | 82                    | 83  | 83  |
| May 28, 1984                 | 80                   | 95  | 97  | 72                    | 94  | 96  |

Table A.8: Continued

| Date               | CN ( $\lambda=0.2$ ) |     |     | CN ( $\lambda=0.05$ ) |     |     |
|--------------------|----------------------|-----|-----|-----------------------|-----|-----|
|                    | WSA                  | WSB | WSC | WSA                   | WSB | WSC |
| July 26, 1984      | 74                   | 75  | 77  | 64                    | 66  | 69  |
| July 27, 1984      | 81                   | 93  | 95  | 73                    | 91  | 94  |
| July 30, 1984      | 84                   | 90  | 93  | 78                    | 87  | 91  |
| August 22, 1984    | 64                   | 71  | 76  | 50                    | 60  | 67  |
| November 4, 1984   | 80                   | 95  | 96  | 73                    | 94  | 95  |
| November 18, 1984  | 71                   | 88  | 90  | 59                    | 84  | 87  |
| November 28, 1984  | 87                   | 94  | 94  | 83                    | 93  | 93  |
| December 20, 1984  | 95                   | 99  | 99  | 95                    | 99  | 99  |
| December 24, 1984  | 95                   | 98  | 100 | 94                    | 98  | 100 |
| January 3, 1985    | 92                   | 97  | 98  | 90                    | 96  | 98  |
| February 11, 1985  | 85                   | 88  | 87  | 79                    | 84  | 83  |
| May 15, 1985       | 78                   | 80  | 83  | 69                    | 72  | 76  |
| June 5, 1985       | 81                   | 86  | 86  | 73                    | 81  | 81  |
| June 10, 1985      | 75                   | 78  | 78  | 65                    | 70  | 69  |
| June 11, 1985      | 83                   | 95  | 94  | 77                    | 93  | 92  |
| July 10, 1985      | 70                   | 73  | 75  | 59                    | 63  | 65  |
| July 30, 1985      | 68                   | 69  | 69  | 56                    | 57  | 57  |
| August 1, 1985     | 65                   | 70  | 70  | 51                    | 59  | 59  |
| August 17, 1985    | 83                   | 88  | 88  | 77                    | 85  | 84  |
| August 25, 1985    | 84                   | 85  | 90  | 78                    | 80  | 87  |
| August 30, 1985    | 65                   | 77  | 79  | 52                    | 68  | 71  |
| September 26, 1985 | 80                   | 84  | 86  | 73                    | 78  | 81  |
| November 2, 1985   | 70                   | 88  | 89  | 59                    | 84  | 86  |
| December 12, 1985  | 93                   | 96  | 96  | 91                    | 95  | 95  |
| February 2, 1986   | 95                   | 98  | 99  | 94                    | 97  | 99  |
| April 28, 1986     | 87                   | 88  | 91  | 82                    | 84  | 88  |
| May 11, 1986       | 75                   | 81  | 79  | 65                    | 73  | 70  |
| July 2, 1986       | 78                   | 82  | 82  | 70                    | 75  | 75  |
| July 20, 1986      | 80                   | 81  | 86  | 73                    | 73  | 81  |
| July 26, 1986      | 78                   | 79  | 83  | 69                    | 71  | 77  |
| October 1, 1986    | 83                   | 83  | 86  | 77                    | 77  | 81  |
| November 5, 1986   | 75                   | 82  | 85  | 65                    | 75  | 79  |
| November 8, 1986   | 70                   | 81  | 83  | 59                    | 73  | 77  |
| November 10, 1986  | 88                   | 92  | 95  | 83                    | 90  | 94  |
| December 8, 1986   | 95                   | 96  | 99  | 94                    | 95  | 99  |
| January 18, 1987   | 92                   | 93  | 97  | 90                    | 92  | 96  |
| February 22, 1987  | 96                   | 97  | 97  | 96                    | 97  | 97  |
| February 26, 1987  | 96                   | 96  | 97  | 95                    | 95  | 97  |
| March 18, 1987     | 99                   | 99  | 98  | 99                    | 99  | 98  |
| March 30, 1987     | 90                   | 91  | 93  | 87                    | 89  | 92  |
| May 12, 1987       | 83                   | 85  | 87  | 77                    | 79  | 82  |
| May 21, 1987       | 80                   | 82  | 82  | 72                    | 75  | 75  |

Table A.8: Continued

| Date               | CN ( $\lambda=0.2$ ) |     |     | CN ( $\lambda=0.05$ ) |     |     |
|--------------------|----------------------|-----|-----|-----------------------|-----|-----|
|                    | WSA                  | WSB | WSC | WSA                   | WSB | WSC |
| May 25, 1987       | 84                   | 86  | 87  | 78                    | 80  | 82  |
| June 16, 1987      | 85                   | 85  | 86  | 79                    | 80  | 81  |
| June 25, 1987      | 68                   | 69  | 70  | 55                    | 58  | 59  |
| July 11, 1987      | 77                   | 83  | 86  | 67                    | 76  | 81  |
| August 22, 1987    | 77                   | 79  | 80  | 68                    | 71  | 73  |
| September 12, 1987 | 81                   | 82  | 83  | 74                    | 75  | 77  |
| November 9, 1987   | 65                   | 65  | 68  | 52                    | 52  | 56  |
| November 16, 1987  | 82                   | 85  | 84  | 75                    | 80  | 79  |
| December 14, 1987  | 86                   | 89  | 89  | 81                    | 85  | 85  |
| December 24, 1987  | 60                   | 63  | 71  | 46                    | 50  | 60  |
| January 18, 1988   | 92                   | 89  | 94  | 90                    | 86  | 92  |
| April 6, 1988      | 97                   | 95  | 98  | 97                    | 93  | 98  |
| May 4, 1988        | 91                   | 80  | 88  | 89                    | 72  | 85  |
| June 2, 1988       | 84                   | 84  | 89  | 79                    | 79  | 85  |
| June 9, 1988       | 83                   | 84  | 84  | 77                    | 78  | 79  |
| August 23, 1988    | 58                   | 60  | 65  | 44                    | 46  | 52  |
| September 16, 1988 | 57                   | 57  | 60  | 42                    | 43  | 46  |
| November 4, 1988   | 73                   | 75  | 77  | 62                    | 65  | 68  |
| November 27, 1988  | 91                   | 94  | 94  | 88                    | 93  | 93  |
| December 21, 1988  | 83                   | 89  | 88  | 77                    | 85  | 84  |
| December 24, 1988  | 96                   | 95  | 96  | 96                    | 94  | 95  |
| January 11, 1989   | 93                   | 93  | 97  | 91                    | 92  | 96  |
| February 3, 1989   | 92                   | 94  | 97  | 90                    | 93  | 96  |
| February 13, 1989  | 90                   | 92  | 94  | 88                    | 90  | 92  |
| February 20, 1989  | 98                   | 94  | 95  | 97                    | 92  | 94  |
| March 20, 1989     | 90                   | 92  | 93  | 87                    | 90  | 92  |
| April 3, 1989      | 95                   | 98  | 98  | 94                    | 98  | 98  |
| May 19, 1989       | 86                   | 86  | 86  | 81                    | 81  | 81  |
| June 12, 1989      | 80                   | 78  | 80  | 72                    | 69  | 73  |
| June 22, 1989      | 91                   | 92  | 92  | 88                    | 90  | 90  |
| July 23, 1989      | 79                   | 79  | 78  | 71                    | 71  | 70  |
| July 27, 1989      | 77                   | 78  | 77  | 68                    | 69  | 69  |
| July 31, 1989      | 79                   | 86  | 89  | 71                    | 82  | 86  |
| August 5, 1989     | 80                   | 85  | 84  | 72                    | 79  | 78  |
| August 18, 1989    | 64                   | 66  | 68  | 51                    | 53  | 56  |
| September 22, 1989 | 76                   | 84  | 90  | 67                    | 78  | 87  |
| September 30, 1989 | 89                   | 91  | 93  | 85                    | 89  | 91  |
| October 16, 1989   | 77                   | 79  | 89  | 68                    | 71  | 86  |
| November 14, 1989  | 93                   | 96  | 97  | 91                    | 95  | 97  |
| January 29, 1990   | 95                   | 97  | 96  | 94                    | 97  | 96  |
| February 3, 1990   | 96                   | 97  | 98  | 96                    | 97  | 98  |
| February 9, 1990   | 93                   | 96  | 97  | 91                    | 96  | 97  |

Table A.8: Continued

| Date               | CN ( $\lambda=0.2$ ) |     |     | CN ( $\lambda=0.05$ ) |     |     |
|--------------------|----------------------|-----|-----|-----------------------|-----|-----|
|                    | WSA                  | WSB | WSC | WSA                   | WSB | WSC |
| February 15, 1990  | 94                   | 95  | 95  | 93                    | 93  | 94  |
| March 16, 1990     | 92                   | 96  | 92  | 90                    | 95  | 90  |
| April 6, 1990      | 94                   | 94  | 95  | 92                    | 92  | 94  |
| May 26, 1990       | 82                   | 83  | 83  | 76                    | 76  | 76  |
| May 28, 1990       | 81                   | 82  | 82  | 73                    | 75  | 76  |
| June 2, 1990       | 80                   | 81  | 83  | 73                    | 73  | 76  |
| June 21, 1990      | 80                   | 78  | 80  | 72                    | 70  | 72  |
| July 30, 1990      | 84                   | 85  | 87  | 78                    | 80  | 82  |
| August 8, 1990     | 77                   | 79  | 80  | 68                    | 71  | 73  |
| August 29, 1990    | 79                   | 80  | 81  | 72                    | 72  | 74  |
| September 9, 1990  | 82                   | 80  | 81  | 75                    | 73  | 74  |
| September 12, 1990 | 85                   | 86  | 86  | 79                    | 81  | 81  |
| October 4, 1990    | 70                   | 75  | 75  | 59                    | 65  | 65  |
| December 2, 1990   | 88                   | 93  | 96  | 84                    | 92  | 95  |
| December 20, 1990  | 90                   | 95  | 93  | 87                    | 94  | 91  |
| December 27, 1990  | 83                   | 86  | 87  | 76                    | 82  | 83  |
| December 30, 1990  | 88                   | 91  | 94  | 84                    | 88  | 92  |
| January 6, 1991    | 98                   | 100 | 99  | 98                    | 100 | 99  |
| February 6, 1991   | 98                   | 98  | 97  | 97                    | 98  | 97  |
| February 13, 1991  | 92                   | 93  | 94  | 90                    | 92  | 92  |
| February 17, 1991  | 90                   | 95  | 99  | 88                    | 94  | 100 |
| March 22, 1991     | 93                   | 94  | 94  | 92                    | 93  | 93  |
| March 29, 1991     | 98                   | 97  | 97  | 97                    | 97  | 96  |
| April 15, 1991     | 96                   | 98  | 98  | 96                    | 98  | 97  |
| April 19, 1991     | 96                   | 95  | 95  | 95                    | 95  | 94  |
| May 9, 1991        | 84                   | 83  | 80  | 79                    | 77  | 72  |
| May 18, 1991       | 79                   | 84  | 80  | 72                    | 79  | 72  |
| May 27, 1991       | 82                   | 83  | 82  | 75                    | 77  | 75  |
| May 29, 1991       | 77                   | 78  | 75  | 68                    | 70  | 66  |
| June 22, 1991      | 79                   | 86  | 83  | 70                    | 81  | 77  |
| June 25, 1991      | 89                   | 90  | 88  | 86                    | 87  | 84  |
| July 10, 1991      | 80                   | 80  | 79  | 73                    | 72  | 72  |
| July 12, 1991      | 74                   | 78  | 77  | 64                    | 69  | 69  |
| August 7, 1991     | 73                   | 74  | 72  | 63                    | 64  | 62  |
| October 5, 1991    | 65                   | 67  | 65  | 53                    | 55  | 52  |
| October 15, 1991   | 80                   | 86  | 80  | 73                    | 81  | 72  |
| November 21, 1991  | 70                   | 71  | 69  | 58                    | 60  | 57  |
| December 13, 1991  | 94                   | 95  | 97  | 93                    | 94  | 97  |
| December 28, 1991  | 94                   | 94  | 94  | 92                    | 93  | 92  |
| February 24, 1992  | 94                   | 96  | 96  | 93                    | 96  | 95  |
| March 30, 1992     | 94                   | 97  | 96  | 93                    | 96  | 95  |
| May 28, 1992       | 68                   | 71  | 71  | 56                    | 60  | 59  |

Table A.8: Continued

| Date                        | CN ( $\lambda=0.2$ ) |             |             | CN ( $\lambda=0.05$ ) |             |             |
|-----------------------------|----------------------|-------------|-------------|-----------------------|-------------|-------------|
|                             | WSA                  | WSB         | WSC         | WSA                   | WSB         | WSC         |
| June 17, 1992               | 87                   | 89          | 87          | 82                    | 86          | 83          |
| June 24, 1992               | 71                   | 75          | 74          | 60                    | 66          | 64          |
| July 1, 1992                | 80                   | 73          | 74          | 73                    | 63          | 64          |
| July 14, 1992               | 72                   | 72          | 75          | 61                    | 61          | 66          |
| July 17, 1992               | 87                   | 88          | 87          | 82                    | 85          | 83          |
| July 22, 1992               | 81                   | 82          | 81          | 74                    | 75          | 74          |
| July 24, 1992               | 70                   | 76          | 76          | 58                    | 66          | 66          |
| August 8, 1992              | 83                   | 83          | 85          | 77                    | 77          | 80          |
| August 27, 1992             | 68                   | 67          | 70          | 56                    | 55          | 59          |
| September 4, 1992           | 86                   | 88          | 90          | 81                    | 84          | 86          |
| September 18, 1992          | 76                   | 75          | 86          | 66                    | 65          | 81          |
| December 20, 1992           | 90                   | 90          | 87          | 87                    | 87          | 83          |
| March 29, 1993              | 94                   | 97          | 97          | 92                    | 97          | 96          |
| April 25, 1993              | 99                   | 95          | 86          | 99                    | 94          | 82          |
| May 9, 1993                 | 82                   | 82          | 77          | 75                    | 75          | 68          |
| June 9, 1993                | 76                   | 78          | 79          | 67                    | 70          | 71          |
| June 21, 1993               | 77                   | 79          | 78          | 68                    | 71          | 70          |
| July 13, 1993               | 54                   | 56          | 64          | 39                    | 41          | 51          |
| July 15, 1993               | 86                   | 87          | 94          | 81                    | 83          | 92          |
| July 26, 1993               | 66                   | 67          | 67          | 53                    | 55          | 54          |
| September 2, 1993           | 71                   | 79          | 80          | 60                    | 72          | 72          |
| September 15, 1993          | 58                   | 63          | 64          | 43                    | 50          | 50          |
| October 18, 1993            | 83                   | 94          | 95          | 77                    | 92          | 93          |
| Mean $\pm$ Std. Dev. (Pre)  | 82 $\pm$ 12          | 82 $\pm$ 12 | 82 $\pm$ 13 | 76 $\pm$ 17           | 76 $\pm$ 17 | 76 $\pm$ 18 |
| Mean $\pm$ Std. Dev. (Post) | 83 $\pm$ 10          | 85 $\pm$ 10 | 87 $\pm$ 9  | 77 $\pm$ 14           | 80 $\pm$ 14 | 82 $\pm$ 13 |

<sup>1</sup> AMC II<sup>2</sup> Final storm before treatment.

Table A.9: Curve numbers for WSA, WSB, and WSC (2002-2008).<sup>1</sup>

| Date               | CN ( $\lambda=0.2$ ) |     |                 | CN ( $\lambda=0.05$ ) |     |                 |
|--------------------|----------------------|-----|-----------------|-----------------------|-----|-----------------|
|                    | WSA                  | WSB | WSC             | WSA                   | WSB | WSC             |
| March 17, 2002     | 97                   | 99  | -- <sup>2</sup> | 97                    | 99  | -- <sup>2</sup> |
| April 28, 2002     | 99                   | 100 | --              | 99                    | 100 | --              |
| May 2, 2002        | 95                   | 97  | --              | 94                    | 97  | --              |
| May 13, 2002       | 87                   | 92  | --              | 83                    | 89  | --              |
| May 17, 2002       | 85                   | 92  | --              | 80                    | 90  | --              |
| June 6, 2002       | 75                   | 78  | --              | 65                    | 69  | --              |
| July 19, 2002      | 84                   | 83  | --              | 78                    | 76  | --              |
| September 25, 2002 | 69                   | 66  | --              | 57                    | 53  | --              |
| October 15, 2002   | 85                   | 82  | --              | 79                    | 75  | --              |
| October 29, 2002   | 92                   | 93  | --              | 90                    | 92  | --              |
| November 5, 2002   | 89                   | 91  | --              | 86                    | 88  | --              |
| November 10, 2002  | 90                   | 98  | --              | 87                    | 98  | --              |
| December 10, 2002  | 98                   | 99  | --              | 98                    | 99  | --              |
| December 13, 2002  | 97                   | 98  | --              | 96                    | 98  | --              |
| December 19, 2002  | 95                   | 95  | --              | 93                    | 94  | --              |
| January 1, 2003    | 99                   | 99  | --              | 99                    | 99  | --              |
| February 3, 2003   | 99                   | 99  | --              | 99                    | 99  | --              |
| April 6, 2003      | 93                   | 94  | --              | 91                    | 92  | --              |
| April 17, 2003     | 92                   | 93  | --              | 90                    | 91  | --              |
| May 8, 2003        | 89                   | 94  | --              | 85                    | 92  | --              |
| June 6, 2003       | 83                   | 87  | --              | 77                    | 82  | --              |
| June 11, 2003      | 85                   | 89  | --              | 80                    | 85  | --              |
| July 10, 2003      | 76                   | 76  | --              | 66                    | 67  | --              |
| August 3, 2003     | 81                   | 82  | --              | 73                    | 75  | --              |
| August 17, 2003    | 82                   | 86  | --              | 75                    | 81  | --              |
| September 3, 2003  | 69                   | 70  | --              | 57                    | 58  | --              |
| September 22, 2003 | 70                   | 67  | --              | 59                    | 55  | --              |
| November 12, 2003  | 89                   | 90  | --              | 85                    | 87  | --              |
| November 18, 2003  | 87                   | 88  | --              | 83                    | 85  | --              |
| November 28, 2003  | 96                   | 96  | --              | 95                    | 95  | --              |
| December 10, 2003  | 97                   | 98  | --              | 96                    | 98  | --              |
| January 17, 2004   | 95                   | 96  | --              | 94                    | 95  | --              |
| February 2, 2004   | 96                   | 98  | --              | 96                    | 98  | --              |
| February 5, 2004   | 88                   | 83  | --              | 84                    | 76  | --              |
| March 5, 2004      | 89                   | 92  | --              | 85                    | 89  | --              |
| May 16, 2004       | 85                   | 84  | --              | 79                    | 78  | --              |
| May 24, 2004       | 74                   | 75  | --              | 63                    | 65  | --              |
| May 26, 2004       | 82                   | 82  | --              | 75                    | 75  | --              |
| May 27, 2004       | 94                   | 96  | --              | 93                    | 96  | --              |
| June 15, 2004      | 89                   | 97  | --              | 85                    | 97  | --              |
| June 22, 2004      | 95                   | 96  | --              | 94                    | 95  | --              |
| June 25, 2004      | 92                   | 95  | --              | 90                    | 94  | --              |

Table A.9: Continued

| Date               | CN ( $\lambda=0.2$ ) |     |     | CN ( $\lambda=0.05$ ) |     |     |
|--------------------|----------------------|-----|-----|-----------------------|-----|-----|
|                    | WSA                  | WSB | WSC | WSA                   | WSB | WSC |
| July 6, 2004       | 87                   | 87  | --  | 82                    | 83  | --  |
| July 26, 2004      | 70                   | 72  | --  | 59                    | 62  | --  |
| July 27, 2004      | 91                   | 89  | --  | 89                    | 86  | --  |
| July 31, 2004      | 87                   | 85  | --  | 83                    | 79  | --  |
| August 5, 2004     | 79                   | 80  | --  | 70                    | 72  | --  |
| September 7, 2004  | 58                   | 58  | --  | 43                    | 43  | --  |
| September 16, 2004 | 81                   | 84  | --  | 73                    | 78  | --  |
| October 13, 2004   | 87                   | 88  | --  | 82                    | 84  | --  |
| October 18, 2004   | 88                   | 91  | --  | 85                    | 88  | --  |
| October 27, 2004   | 95                   | 95  | --  | 94                    | 94  | --  |
| November 4, 2004   | 94                   | 92  | --  | 92                    | 90  | --  |
| November 11, 2004  | 94                   | 92  | --  | 92                    | 90  | --  |
| November 30, 2004  | 96                   | 98  | --  | 96                    | 98  | --  |
| December 9, 2004   | 99                   | 100 | --  | 99                    | 100 | --  |
| December 23, 2004  | 90                   | 93  | --  | 87                    | 91  | --  |
| April 29, 2005     | 92                   | 96  | 100 | 90                    | 96  | 100 |
| May 19, 2005       | 78                   | 82  | 78  | 70                    | 76  | 69  |
| May 22, 2005       | 95                   | 98  | 92  | 94                    | 98  | 90  |
| June 3, 2005       | 99                   | 100 | 94  | 99                    | 100 | 93  |
| June 20, 2005      | 70                   | 78  | 75  | 59                    | 70  | 65  |
| July 1, 2005       | 76                   | 75  | 74  | 67                    | 65  | 64  |
| July 7, 2005       | 86                   | 81  | 92  | 82                    | 74  | 90  |
| July 17, 2005      | 83                   | 84  | 81  | 77                    | 78  | 74  |
| July 18, 2005      | 96                   | 95  | 96  | 95                    | 94  | 95  |
| July 27, 2005      | 85                   | 82  | 87  | 80                    | 75  | 83  |
| August 5, 2005     | 80                   | 78  | 78  | 72                    | 70  | 70  |
| August 16, 2005    | 82                   | 80  | 80  | 76                    | 73  | 73  |
| August 19, 2005    | 82                   | 80  | 81  | 76                    | 72  | 74  |
| December 3, 2005   | 86                   | 86  | 93  | 81                    | 81  | 91  |
| December 15, 2005  | 85                   | 82  | 86  | 80                    | 75  | 81  |
| January 17, 2006   | 87                   | 91  | 88  | 82                    | 88  | 84  |
| January 23, 2006   | 90                   | 94  | 98  | 87                    | 93  | 98  |
| March 13, 2006     | 97                   | 99  | 99  | 96                    | 99  | 99  |
| May 2, 2006        | 88                   | 84  | 90  | 86                    | 96  | 95  |
| May 25, 2006       | 81                   | 74  | 80  | 73                    | 87  | 83  |
| May 31, 2006       | 89                   | 85  | 93  | 91                    | 95  | 94  |
| July 5, 2006       | 84                   | 78  | 88  | 84                    | 92  | 90  |
| September 23, 2006 | 75                   | 65  | 77  | 68                    | 77  | 68  |
| September 23, 2006 | 88                   | 85  | 92  | 90                    | 92  | 90  |
| September 30, 2006 | 86                   | 82  | 89  | 86                    | 89  | 85  |
| October 26, 2006   | 72                   | 62  | 78  | 70                    | 84  | 78  |
| March 28, 2007     | 94                   | 93  | 88  | 84                    | 91  | 89  |

Table A.9: Continued

| Date                         | CN ( $\lambda=0.2$ ) |             |             | CN ( $\lambda=0.05$ ) |            |             |
|------------------------------|----------------------|-------------|-------------|-----------------------|------------|-------------|
|                              | WSA                  | WSB         | WSC         | WSA                   | WSB        | WSC         |
| May 16, 2007                 | 79                   | 71          | 82          | 75                    | 87         | 83          |
| June 1, 2007                 | 68                   | 56          | 68          | 55                    | 78         | 70          |
| June 5, 2007                 | 82                   | 75          | 82          | 75                    | 85         | 80          |
| June 18, 2007                | 77                   | 68          | 75          | 66                    | 79         | 71          |
| June 29, 2007                | 87                   | 82          | 83          | 77                    | 88         | 83          |
| August 3, 2007               | 82                   | 75          | 80          | 72                    | 86         | 81          |
| November 14, 2007            | 72                   | 61          | 68          | 57                    | 74         | 64          |
| November 26, 2007            | 87                   | 82          | 88          | 84                    | 92         | 90          |
| December 10, 2007            | 76                   | 66          | 78          | 70                    | 82         | 76          |
| December 13, 2007            | 90                   | 88          | 94          | 92                    | 94         | 93          |
| December 20, 2007            | 84                   | 78          | 91          | 88                    | 93         | 91          |
| January 29, 2008             | 89                   | 86          | 94          | 92                    | 95         | 94          |
| February 6, 2008             | 96                   | 96          | 99          | 99                    | 100        | 100         |
| May 11, 2008                 | 85                   | 80          | 88          | 84                    | 89         | 85          |
| June 3, 2008                 | 87                   | 83          | 89          | 85                    | 90         | 86          |
| August 26, 2008              | 69                   | 57          | 67          | 55                    | 70         | 58          |
| November 14, 2008            | 76                   | 66          | 83          | 76                    | 79         | 71          |
| December 10, 2008            | 68                   | 56          | 76          | 66                    | 74         | 64          |
| December 10, 2008            | 81                   | 74          | 91          | 89                    | 86         | 81          |
| December 16, 2008            | 92                   | 90          | 94          | 93                    | 96         | 95          |
| Mean $\pm$ Std. Dev. (02-04) | 88 $\pm$ 9           | 83 $\pm$ 13 | 89 $\pm$ 10 | 85 $\pm$ 13           | --         | --          |
| Mean $\pm$ Std. Dev. (05-08) | 84 $\pm$ 8           | 78 $\pm$ 11 | 85 $\pm$ 8  | 80 $\pm$ 12           | 87 $\pm$ 8 | 83 $\pm$ 11 |

<sup>1</sup>AMC II<sup>2</sup>No data available.



## APPENDIX B: FWMC TABLES

Table B.1: Growing, non-growing, and yearly Sulfate FWMC  $\pm$  Std. Dev. (1982-1993).

|          | SO <sub>4</sub> (mg L <sup>-1</sup> ) |                |                | Sample Size (n) |         |             |
|----------|---------------------------------------|----------------|----------------|-----------------|---------|-------------|
|          | Year                                  | Growing        | Non-Growing    | Year            | Growing | Non-Growing |
| WSA      | 13.3 $\pm$ 1.7                        | 12.3 $\pm$ 1.5 | 13.4 $\pm$ 1.0 | 44              | 25      | 19          |
| WSB 1982 | 15.4 $\pm$ 2.6                        | 13.6 $\pm$ 1.4 | 15.7 $\pm$ 3.4 | 45              | 25      | 20          |
| WSC      | 13.8 $\pm$ 2.4                        | 11.0 $\pm$ 2.0 | 14.3 $\pm$ 2.0 | 42              | 22      | 20          |
| WSA      | 15.1 $\pm$ 2.7                        | 16.0 $\pm$ 2.6 | 12.7 $\pm$ 2.6 | 47              | 22      | 25          |
| WSB 1983 | 13.8 $\pm$ 3.8                        | 14.5 $\pm$ 3.4 | 12.1 $\pm$ 4.0 | 48              | 23      | 25          |
| WSC      | 11.8 $\pm$ 3.7                        | 12.2 $\pm$ 4.2 | 10.7 $\pm$ 3.3 | 42              | 17      | 25          |
| WSA      | 9.4 $\pm$ 2.0                         | 9.3 $\pm$ 2.4  | 9.5 $\pm$ 1.5  | 52              | 26      | 26          |
| WSB 1984 | 6.3 $\pm$ 2.9                         | 9.7 $\pm$ 1.8  | 5.4 $\pm$ 3.5  | 52              | 26      | 26          |
| WSC      | 10.8 $\pm$ 2.0                        | 9.3 $\pm$ 1.9  | 11.4 $\pm$ 2.0 | 52              | 26      | 26          |
| WSA      | 10.7 $\pm$ 1.3                        | 11.1 $\pm$ 1.4 | 10.6 $\pm$ 1.3 | 48              | 23      | 25          |
| WSB 1985 | 11.1 $\pm$ 1.2                        | 11.1 $\pm$ 1.4 | 11.1 $\pm$ 0.8 | 52              | 27      | 25          |
| WSC      | 10.5 $\pm$ 1.5                        | 10.2 $\pm$ 1.3 | 10.6 $\pm$ 1.7 | 52              | 27      | 25          |
| WSA      | 11.1 $\pm$ 2.6                        | 9.3 $\pm$ 2.0  | 11.4 $\pm$ 1.6 | 48              | 24      | 24          |
| WSB 1986 | 14.0 $\pm$ 2.3                        | 11.8 $\pm$ 1.6 | 14.4 $\pm$ 2.5 | 43              | 19      | 24          |
| WSC      | 10.0 $\pm$ 4.4                        | 5.1 $\pm$ 3.7  | 11.1 $\pm$ 2.3 | 50              | 25      | 25          |
| WSA      | 10.3 $\pm$ 1.6                        | 9.0 $\pm$ 1.7  | 10.8 $\pm$ 1.2 | 44              | 21      | 23          |
| WSB 1987 | 11.8 $\pm$ 2.1                        | 11.8 $\pm$ 2.5 | 11.8 $\pm$ 1.6 | 42              | 20      | 22          |
| WSC      | 9.0 $\pm$ 3.1                         | 8.3 $\pm$ 3.8  | 9.3 $\pm$ 1.9  | 46              | 24      | 22          |
| WSA      | 6.6 $\pm$ 2.7                         | 8.4 $\pm$ 2.8  | 5.9 $\pm$ 1.3  | 41              | 16      | 25          |
| WSB 1988 | 7.4 $\pm$ 2.7                         | 8.8 $\pm$ 2.4  | 7.0 $\pm$ 2.1  | 38              | 14      | 24          |
| WSC      | 8.3 $\pm$ 5.1                         | 11.6 $\pm$ 6.2 | 7.0 $\pm$ 2.9  | 43              | 18      | 25          |
| WSA      | 9.6 $\pm$ 2.4                         | 11.1 $\pm$ 1.3 | 8.7 $\pm$ 2.8  | 52              | 27      | 25          |
| WSB 1989 | 11.8 $\pm$ 2.8                        | 12.9 $\pm$ 2.3 | 11.0 $\pm$ 3.0 | 50              | 27      | 23          |
| WSC      | 10.4 $\pm$ 2.8                        | 11.6 $\pm$ 2.9 | 9.5 $\pm$ 2.1  | 51              | 27      | 24          |
| WSA      | 8.0 $\pm$ 1.4                         | 7.0 $\pm$ 1.1  | 8.5 $\pm$ 0.7  | 52              | 27      | 25          |
| WSB 1990 | 9.1 $\pm$ 1.3                         | 8.0 $\pm$ 1.0  | 9.4 $\pm$ 1.3  | 51              | 27      | 24          |
| WSC      | 7.6 $\pm$ 2.2                         | 6.1 $\pm$ 1.9  | 8.3 $\pm$ 1.4  | 51              | 27      | 24          |
| WSA      | 9.4 $\pm$ 1.8                         | 8.4 $\pm$ 0.6  | 9.9 $\pm$ 2.1  | 52              | 27      | 25          |
| WSB 1991 | 11.2 $\pm$ 2.0                        | 9.4 $\pm$ 1.4  | 11.6 $\pm$ 1.8 | 53              | 27      | 26          |
| WSC      | 8.1 $\pm$ 1.0                         | 8.1 $\pm$ 0.9  | 8.1 $\pm$ 1.0  | 34              | 18      | 16          |
| WSA      | 11.8 $\pm$ 1.2                        | 11.9 $\pm$ 1.2 | 11.7 $\pm$ 1.1 | 52              | 27      | 25          |
| WSB 1992 | 12.6 $\pm$ 0.9                        | 12.6 $\pm$ 0.7 | 12.6 $\pm$ 1.1 | 51              | 28      | 23          |
| WSC      | 10.7 $\pm$ 2.2                        | 9.9 $\pm$ 2.5  | 11.2 $\pm$ 1.5 | 51              | 28      | 23          |
| WSA      | 6.5 $\pm$ 2.0                         | 6.5 $\pm$ 2.2  | 6.5 $\pm$ 1.6  | 44              | 27      | 17          |
| WSB 1993 | 7.0 $\pm$ 2.4                         | 5.4 $\pm$ 2.6  | 8.0 $\pm$ 1.9  | 43              | 27      | 16          |
| WSC      | 6.2 $\pm$ 2.7                         | 4.0 $\pm$ 2.2  | 7.6 $\pm$ 2.7  | 43              | 27      | 16          |

Table B.2: Growing, non-growing, and yearly Sulfate FWMC  $\pm$  Std. Dev. (2002-2008).

|          | SO <sub>4</sub> (mg L <sup>-1</sup> ) |                |                | Sample Size (n) |         |             |
|----------|---------------------------------------|----------------|----------------|-----------------|---------|-------------|
|          | Year                                  | Growing        | Non-Growing    | Year            | Growing | Non-Growing |
| WSA      | 10.8 $\pm$ 1.9                        | 9.8 $\pm$ 1.4  | 11.8 $\pm$ 1.8 | 47              | 23      | 24          |
| WSB 2002 | 11.6 $\pm$ 1.8                        | 10.2 $\pm$ 1.3 | 12.7 $\pm$ 1.6 | 46              | 21      | 25          |
| WSC      | 10.3 $\pm$ 2.5                        | 9.4 $\pm$ 2.7  | 10.8 $\pm$ 1.3 | 48              | 24      | 24          |
| WSA      | 10.5 $\pm$ 1.8                        | 10.0 $\pm$ 2.0 | 10.7 $\pm$ 1.5 | 49              | 25      | 24          |
| WSB 2003 | 12.0 $\pm$ 1.8                        | 13.1 $\pm$ 2.2 | 11.3 $\pm$ 1.3 | 49              | 25      | 24          |
| WSC      | 10.0 $\pm$ 2.2                        | 9.9 $\pm$ 2.4  | 10.1 $\pm$ 1.7 | 49              | 25      | 24          |
| WSA      | 9.1 $\pm$ 0.8                         | 8.9 $\pm$ 0.8  | 9.4 $\pm$ 0.7  | 51              | 28      | 23          |
| WSB 2004 | 9.5 $\pm$ 0.9                         | 9.3 $\pm$ 0.8  | 9.8 $\pm$ 0.7  | 51              | 28      | 23          |
| WSC      | 8.3 $\pm$ 0.9                         | 8.1 $\pm$ 1.0  | 8.5 $\pm$ 0.6  | 51              | 28      | 23          |
| WSA      | 8.5 $\pm$ 1.0                         | 8.7 $\pm$ 1.0  | 8.4 $\pm$ 1.0  | 43              | 22      | 21          |
| WSB 2005 | 8.9 $\pm$ 0.8                         | 8.6 $\pm$ 0.4  | 9.0 $\pm$ 0.9  | 41              | 20      | 21          |
| WSC      | 7.8 $\pm$ 1.7                         | 7.9 $\pm$ 1.6  | 7.8 $\pm$ 1.7  | 40              | 19      | 21          |
| WSA      | 8.7 $\pm$ 2.0                         | 7.6 $\pm$ 2.4  | 9.2 $\pm$ 1.5  | 48              | 24      | 24          |
| WSB 2006 | 8.8 $\pm$ 1.7                         | 7.8 $\pm$ 1.7  | 9.2 $\pm$ 1.7  | 47              | 23      | 24          |
| WSC      | 8.6 $\pm$ 2.1                         | 8.5 $\pm$ 2.3  | 8.6 $\pm$ 1.6  | 49              | 25      | 24          |
| WSA      | 8.6 $\pm$ 1.1                         | 8.6 $\pm$ 1.6  | 8.7 $\pm$ 0.7  | 35              | 14      | 21          |
| WSB 2007 | 0.0 $\pm$ 0.8                         |                |                | 0               | 0       | 0           |
| WSC      | 8.5 $\pm$ 4.9                         | 7.0 $\pm$ 1.7  | 9.1 $\pm$ 5.4  | 36              | 14      | 22          |
| WSA      | 8.2 $\pm$ 1.8                         | 7.7 $\pm$ 0.7  | 8.4 $\pm$ 2.1  | 35              | 13      | 22          |
| WSB 2008 | 8.7 $\pm$ 0.0                         | 8.5 $\pm$ 0.5  | 8.8 $\pm$ 0.9  | 35              | 13      | 22          |
| WSC      | 8.0 $\pm$ 1.3                         | 7.5 $\pm$ 0.5  | 8.1 $\pm$ 1.5  | 36              | 14      | 22          |

Table B.3: Growing, Non-Growing, and Yearly Magnesium FWMC  $\pm$  Std. Dev. (1982-1993).

| Table 2: Soil Growing, Non-Growing, and Ready Magnesium Values (mg L <sup>-1</sup> and Sample Size (n) for 1982-1993). |      |                          |         |                 |      |         |             |
|--|------|--------------------------|---------|-----------------|------|---------|-------------|
|  |      | Mg (mg L <sup>-1</sup> ) |         | Sample Size (n) |      |         |             |
|  |      | Year                     | Growing | Non-Growing     | Year | Growing | Non-Growing |
| WSA  | 1982 | 1.6±0.5                  | 1.7±0.4 | 1.6±0.4         | 45   | 25      | 20          |
| WSB  |      | 1.9±0.4                  | 2.0±0.3 | 1.9±0.5         | 45   | 25      | 20          |
| WSC  |      | 1.6±0.3                  | 1.7±0.3 | 1.6±0.2         | 42   | 22      | 20          |
| WSA  | 1983 | 1.6±0.4                  | 1.6±0.5 | 1.6±0.2         | 47   | 22      | 25          |
| WSB  |      | 1.7±0.4                  | 1.7±0.4 | 1.9±0.2         | 48   | 23      | 25          |
| WSC  |      | 1.5±1.0                  | 1.5±1.6 | 1.6±0.2         | 42   | 17      | 25          |
| WSA  | 1984 | 1.5±0.4                  | 1.4±0.4 | 1.5±0.1         | 52   | 26      | 26          |
| WSB  |      | 2.0±0.1                  | 1.8±0.2 | 2.1±0.1         | 52   | 26      | 26          |
| WSC  |      | 1.9±0.3                  | 1.8±0.3 | 2.0±0.3         | 52   | 26      | 26          |
| WSA  | 1985 | 1.6±0.4                  | 2.0±0.3 | 1.5±0.2         | 48   | 23      | 25          |
| WSB  |      | 2.1±0.2                  | 2.2±0.3 | 2.1±0.1         | 52   | 27      | 25          |
| WSC  |      | 2.1±0.5                  | 2.4±0.6 | 2.0±0.1         | 52   | 27      | 25          |
| WSA  | 1986 | 1.8±0.3                  | 2.1±0.3 | 1.8±0.1         | 48   | 24      | 24          |
| WSB  |      | 2.1±0.1                  | 2.1±0.1 | 2.0±0.1         | 43   | 19      | 24          |
| WSC  |      | 2.2±0.5                  | 2.6±0.5 | 2.1±0.3         | 50   | 25      | 25          |
| WSA  | 1987 | 1.7±0.5                  | 1.9±0.4 | 1.6±0.5         | 44   | 21      | 23          |
| WSB  |      | 1.9±0.3                  | 2.0±0.2 | 1.9±0.4         | 42   | 20      | 22          |
| WSC  |      | 2.0±0.6                  | 2.3±0.6 | 1.9±0.5         | 46   | 24      | 22          |
| WSA  | 1988 | 1.9±0.8                  | 2.2±1.0 | 1.8±0.3         | 41   | 16      | 25          |
| WSB  |      | 2.1±0.7                  | 2.5±1.0 | 2.0±0.2         | 38   | 14      | 24          |
| WSC  |      | 2.4±1.1                  | 3.2±1.3 | 2.0±0.3         | 43   | 18      | 25          |
| WSA  | 1989 | 1.5±0.1                  | 1.5±0.2 | 1.5±0.1         | 52   | 27      | 25          |
| WSB  |      | 1.8±0.1                  | 1.8±0.1 | 1.8±0.2         | 50   | 27      | 23          |
| WSC  |      | 1.7±0.2                  | 1.8±0.3 | 1.7±0.1         | 51   | 27      | 24          |
| WSA  | 1990 | 1.5±0.2                  | 1.6±0.2 | 1.5±0.1         | 52   | 27      | 25          |
| WSB  |      | 1.8±0.1                  | 1.8±0.1 | 1.8±0.1         | 51   | 27      | 24          |
| WSC  |      | 1.8±0.4                  | 1.9±0.4 | 1.7±0.1         | 51   | 27      | 24          |
| WSA  | 1991 | 1.5±0.3                  | 1.6±0.3 | 1.5±0.3         | 52   | 27      | 25          |
| WSB  |      | 1.7±0.1                  | 1.8±0.1 | 1.7±0.1         | 53   | 27      | 26          |
| WSC  |      | 1.7±0.3                  | 1.8±0.3 | 1.6±0.1         | 35   | 19      | 16          |
| WSA  | 1992 | 1.6±0.1                  | 1.7±0.1 | 1.6±0.2         | 52   | 27      | 25          |
| WSB  |      | 1.9±0.1                  | 1.9±0.1 | 1.8±0.1         | 51   | 28      | 23          |
| WSC  |      | 1.8±0.2                  | 2.0±0.1 | 1.7±0.2         | 51   | 28      | 23          |
| WSA  | 1993 | 1.5±0.2                  | 1.7±0.1 | 1.5±0.1         | 44   | 27      | 17          |
| WSB  |      | 1.8±0.1                  | 1.9±0.1 | 1.7±0.1         | 43   | 27      | 16          |
| WSC  |      | 1.8±0.3                  | 2.2±0.3 | 1.6±0.1         | 43   | 27      | 16          |

Table B.4: Growing, non-growing, and yearly Magnesium FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | Mg (mg L <sup>-1</sup> ) |               |               | Sample Size (n) |         |             |
|-----|------|--------------------------|---------------|---------------|-----------------|---------|-------------|
|     |      | Year                     | Growing       | Non-Growing   | Year            | Growing | Non-Growing |
| WSA | 2002 | 1.4 $\pm$ 0.2            | 1.7 $\pm$ 1.1 | 1.9 $\pm$ 1.2 | 47              | 23      | 24          |
| WSB |      | 1.6 $\pm$ 0.2            | 1.7 $\pm$ 1.1 | 1.8 $\pm$ 1.3 | 46              | 21      | 25          |
| WSC |      | 1.5 $\pm$ 0.2            | 1.9 $\pm$ 0.7 | 1.9 $\pm$ 0.8 | 48              | 24      | 24          |
| WSA | 2003 | 1.4 $\pm$ 0.2            | 1.5 $\pm$ 0.5 | 1.8 $\pm$ 0.6 | 51              | 27      | 24          |
| WSB |      | 1.5 $\pm$ 0.1            | 1.6 $\pm$ 0.5 | 1.7 $\pm$ 0.6 | 51              | 27      | 24          |
| WSC |      | 1.4 $\pm$ 0.3            | 1.5 $\pm$ 0.3 | 1.6 $\pm$ 0.4 | 51              | 27      | 24          |
| WSA | 2004 | 1.3 $\pm$ 0.1            | 1.2 $\pm$ 0.7 | 1.2 $\pm$ 0.8 | 51              | 28      | 23          |
| WSB |      | 1.4 $\pm$ 0.1            | 1.1 $\pm$ 0.3 | 1.1 $\pm$ 0.3 | 51              | 28      | 23          |
| WSC |      | 1.4 $\pm$ 0.1            | 1.2 $\pm$ 0.2 | 1.2 $\pm$ 0.2 | 51              | 28      | 23          |
| WSA | 2005 | 1.4 $\pm$ 0.2            | 2.1 $\pm$ 1.2 | 2.5 $\pm$ 1.2 | 43              | 22      | 21          |
| WSB |      | 1.5 $\pm$ 0.2            | 1.8 $\pm$ 0.7 | 2.1 $\pm$ 0.3 | 41              | 20      | 21          |
| WSC |      | 1.4 $\pm$ 0.2            | 1.6 $\pm$ 0.5 | 1.9 $\pm$ 0.2 | 40              | 19      | 21          |
| WSA | 2006 | 1.2 $\pm$ 0.2            | 1.5 $\pm$ 0.5 | 1.4 $\pm$ 0.6 | 48              | 24      | 24          |
| WSB |      | 1.3 $\pm$ 0.1            | 1.5 $\pm$ 0.4 | 1.6 $\pm$ 0.5 | 47              | 23      | 24          |
| WSC |      | 1.3 $\pm$ 0.2            | 1.6 $\pm$ 0.3 | 1.6 $\pm$ 0.4 | 49              | 25      | 24          |
| WSA | 2007 | 1.5 $\pm$ 0.3            | 1.5 $\pm$ 1.2 | 2.3 $\pm$ 1.3 | 35              | 14      | 21          |
| WSB |      | 1.7 $\pm$ 0.3            | 1.5 $\pm$ 1.1 | 1.8 $\pm$ 1.3 | 33              | 13      | 20          |
| WSC |      | 1.6 $\pm$ 0.4            | 1.6 $\pm$ 0.9 | 2.0 $\pm$ 1.0 | 36              | 14      | 22          |
| WSA | 2008 | 1.4 $\pm$ 0.5            | 1.1 $\pm$ 0.6 | 1.2 $\pm$ 0.5 | 35              | 13      | 22          |
| WSB |      | 1.4 $\pm$ 0.4            | 1.0 $\pm$ 0.6 | 1.0 $\pm$ 0.4 | 35              | 13      | 22          |
| WSC |      | 1.3 $\pm$ 0.4            | 1.0 $\pm$ 0.6 | 0.9 $\pm$ 0.2 | 36              | 14      | 22          |

Table B.5: Growing, non-growing, and yearly Calcium FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | Ca (mg L <sup>-1</sup> ) |               |               | Sample Size (n) |         |             |
|-----|------|--------------------------|---------------|---------------|-----------------|---------|-------------|
|     |      | Year                     | Growing       | Non-Growing   | Year            | Growing | Non-Growing |
| WSA | 1982 | 1.9 $\pm$ 0.6            | 2.2 $\pm$ 0.5 | 1.8 $\pm$ 0.5 | 45              | 25      | 20          |
| WSB |      | 2.2 $\pm$ 1.1            | 2.6 $\pm$ 1.1 | 2.1 $\pm$ 0.8 | 45              | 25      | 20          |
| WSC |      | 2.0 $\pm$ 1.3            | 2.5 $\pm$ 1.4 | 1.8 $\pm$ 0.7 | 42              | 22      | 20          |
| WSA | 1983 | 2.2 $\pm$ 0.6            | 2.4 $\pm$ 0.6 | 1.9 $\pm$ 0.4 | 47              | 22      | 25          |
| WSB |      | 2.1 $\pm$ 0.7            | 2.0 $\pm$ 0.8 | 2.3 $\pm$ 0.5 | 48              | 23      | 25          |
| WSC |      | 1.9 $\pm$ 0.9            | 1.9 $\pm$ 1.2 | 2.1 $\pm$ 0.6 | 42              | 17      | 25          |
| WSA | 1984 | 2.0 $\pm$ 0.7            | 1.9 $\pm$ 0.7 | 2.1 $\pm$ 0.5 | 52              | 26      | 26          |
| WSB |      | 2.6 $\pm$ 0.4            | 2.3 $\pm$ 0.2 | 2.7 $\pm$ 0.4 | 52              | 26      | 26          |
| WSC |      | 2.5 $\pm$ 0.4            | 2.4 $\pm$ 0.3 | 2.6 $\pm$ 0.4 | 52              | 26      | 26          |
| WSA | 1985 | 2.1 $\pm$ 0.9            | 3.1 $\pm$ 0.9 | 2.0 $\pm$ 0.5 | 48              | 23      | 25          |
| WSB |      | 2.7 $\pm$ 0.4            | 2.9 $\pm$ 0.5 | 2.6 $\pm$ 0.3 | 52              | 27      | 25          |
| WSC |      | 2.7 $\pm$ 0.7            | 3.1 $\pm$ 0.7 | 2.6 $\pm$ 0.4 | 52              | 27      | 25          |
| WSA | 1986 | 2.6 $\pm$ 0.9            | 3.6 $\pm$ 0.7 | 2.4 $\pm$ 0.4 | 48              | 24      | 24          |
| WSB |      | 2.4 $\pm$ 0.6            | 3.2 $\pm$ 0.6 | 2.3 $\pm$ 0.3 | 43              | 19      | 24          |
| WSC |      | 2.7 $\pm$ 1.2            | 4.2 $\pm$ 1.1 | 2.4 $\pm$ 0.6 | 50              | 25      | 25          |
| WSA | 1987 | 2.4 $\pm$ 1.4            | 3.0 $\pm$ 1.3 | 2.1 $\pm$ 1.4 | 44              | 21      | 23          |
| WSB |      | 2.6 $\pm$ 1.3            | 2.8 $\pm$ 1.0 | 2.5 $\pm$ 1.5 | 42              | 20      | 22          |
| WSC |      | 2.8 $\pm$ 1.8            | 3.8 $\pm$ 1.8 | 2.5 $\pm$ 1.5 | 46              | 24      | 22          |
| WSA | 1988 | 2.7 $\pm$ 1.5            | 3.3 $\pm$ 1.7 | 2.5 $\pm$ 0.6 | 41              | 16      | 25          |
| WSB |      | 3.2 $\pm$ 2.2            | 4.7 $\pm$ 2.9 | 2.7 $\pm$ 0.3 | 38              | 14      | 24          |
| WSC |      | 3.6 $\pm$ 2.2            | 4.8 $\pm$ 2.5 | 3.1 $\pm$ 0.7 | 43              | 18      | 25          |
| WSA | 1989 | 2.1 $\pm$ 0.4            | 2.1 $\pm$ 0.4 | 2.1 $\pm$ 0.4 | 52              | 27      | 25          |
| WSB |      | 2.3 $\pm$ 0.4            | 2.0 $\pm$ 0.2 | 2.4 $\pm$ 0.5 | 50              | 27      | 23          |
| WSC |      | 2.5 $\pm$ 0.5            | 2.5 $\pm$ 0.6 | 2.5 $\pm$ 0.3 | 51              | 27      | 24          |
| WSA | 1990 | 1.8 $\pm$ 0.4            | 2.1 $\pm$ 0.4 | 1.7 $\pm$ 0.2 | 52              | 27      | 25          |
| WSB |      | 2.1 $\pm$ 0.3            | 2.1 $\pm$ 0.4 | 2.0 $\pm$ 0.2 | 51              | 27      | 24          |
| WSC |      | 2.4 $\pm$ 0.9            | 2.6 $\pm$ 0.9 | 2.3 $\pm$ 0.4 | 51              | 27      | 24          |
| WSA | 1991 | 1.8 $\pm$ 1.0            | 2.1 $\pm$ 1.0 | 1.7 $\pm$ 0.7 | 52              | 27      | 25          |
| WSB |      | 2.0 $\pm$ 0.4            | 2.2 $\pm$ 0.4 | 2.0 $\pm$ 0.4 | 53              | 27      | 26          |
| WSC |      | 2.1 $\pm$ 0.6            | 2.5 $\pm$ 0.6 | 1.9 $\pm$ 0.1 | 35              | 19      | 16          |
| WSA | 1992 | 1.9 $\pm$ 0.4            | 2.1 $\pm$ 0.3 | 1.7 $\pm$ 0.4 | 52              | 27      | 25          |
| WSB |      | 2.0 $\pm$ 0.3            | 2.2 $\pm$ 0.2 | 2.0 $\pm$ 0.3 | 51              | 28      | 23          |
| WSC |      | 2.3 $\pm$ 0.6            | 2.7 $\pm$ 0.4 | 2.1 $\pm$ 0.6 | 51              | 28      | 23          |
| WSA | 1993 | 2.0 $\pm$ 0.8            | 2.4 $\pm$ 0.7 | 1.8 $\pm$ 0.2 | 44              | 27      | 17          |
| WSB |      | 2.6 $\pm$ 0.7            | 3.3 $\pm$ 0.6 | 2.1 $\pm$ 0.2 | 43              | 27      | 16          |
| WSC |      | 2.8 $\pm$ 1.0            | 4.0 $\pm$ 0.8 | 2.2 $\pm$ 0.2 | 43              | 27      | 16          |

Table B.6: Growing, non-growing, and yearly Calcium FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | Ca (mg L <sup>-1</sup> ) |               |               | Sample Size (n) |         |             |
|-----|------|--------------------------|---------------|---------------|-----------------|---------|-------------|
|     |      | Year                     | Growing       | Non-Growing   | Year            | Growing | Non-Growing |
| WSA | 2002 | 1.7 $\pm$ 1.1            | 1.9 $\pm$ 1.2 | 1.5 $\pm$ 0.3 | 47              | 23      | 24          |
| WSB |      | 1.7 $\pm$ 1.1            | 1.8 $\pm$ 1.3 | 1.7 $\pm$ 0.3 | 46              | 21      | 25          |
| WSC |      | 1.9 $\pm$ 0.7            | 1.9 $\pm$ 0.8 | 1.8 $\pm$ 0.3 | 48              | 24      | 24          |
| WSA | 2003 | 1.5 $\pm$ 0.5            | 1.8 $\pm$ 0.6 | 1.4 $\pm$ 0.2 | 51              | 27      | 24          |
| WSB |      | 1.6 $\pm$ 0.5            | 1.7 $\pm$ 0.6 | 1.5 $\pm$ 0.1 | 51              | 27      | 24          |
| WSC |      | 1.5 $\pm$ 0.3            | 1.6 $\pm$ 0.4 | 1.5 $\pm$ 0.2 | 51              | 27      | 24          |
| WSA | 2004 | 1.2 $\pm$ 0.7            | 1.2 $\pm$ 0.8 | 1.1 $\pm$ 0.1 | 51              | 28      | 23          |
| WSB |      | 1.1 $\pm$ 0.3            | 1.1 $\pm$ 0.3 | 1.1 $\pm$ 0.2 | 51              | 28      | 23          |
| WSC |      | 1.2 $\pm$ 0.2            | 1.2 $\pm$ 0.2 | 1.2 $\pm$ 0.2 | 51              | 28      | 23          |
| WSA | 2005 | 2.1 $\pm$ 1.2            | 2.5 $\pm$ 1.2 | 1.8 $\pm$ 1.0 | 43              | 22      | 21          |
| WSB |      | 1.8 $\pm$ 0.7            | 2.1 $\pm$ 0.3 | 1.6 $\pm$ 1.0 | 41              | 20      | 21          |
| WSC |      | 1.6 $\pm$ 0.5            | 1.9 $\pm$ 0.2 | 1.5 $\pm$ 0.7 | 40              | 19      | 21          |
| WSA | 2006 | 1.5 $\pm$ 0.5            | 1.4 $\pm$ 0.6 | 1.5 $\pm$ 0.2 | 48              | 24      | 24          |
| WSB |      | 1.5 $\pm$ 0.4            | 1.6 $\pm$ 0.5 | 1.5 $\pm$ 0.2 | 47              | 23      | 24          |
| WSC |      | 1.6 $\pm$ 0.3            | 1.6 $\pm$ 0.4 | 1.6 $\pm$ 0.2 | 49              | 25      | 24          |
| WSA | 2007 | 1.5 $\pm$ 1.2            | 2.3 $\pm$ 1.3 | 1.2 $\pm$ 0.4 | 35              | 14      | 21          |
| WSB |      | 1.5 $\pm$ 1.1            | 1.8 $\pm$ 1.3 | 1.4 $\pm$ 0.3 | 33              | 13      | 20          |
| WSC |      | 1.6 $\pm$ 0.9            | 2.0 $\pm$ 1.0 | 1.5 $\pm$ 0.7 | 36              | 14      | 22          |
| WSA | 2008 | 1.1 $\pm$ 0.6            | 1.2 $\pm$ 0.5 | 1.1 $\pm$ 0.7 | 35              | 13      | 22          |
| WSB |      | 1.0 $\pm$ 0.6            | 1.0 $\pm$ 0.4 | 1.0 $\pm$ 0.7 | 35              | 13      | 22          |
| WSC |      | 1.0 $\pm$ 0.6            | 0.9 $\pm$ 0.2 | 1.0 $\pm$ 0.8 | 36              | 14      | 22          |

Table B.7: Growing, non-growing, and yearly Potassium FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | K (mg L <sup>-1</sup> ) |               |               | Sample Size (n) |         |             |
|-----|------|-------------------------|---------------|---------------|-----------------|---------|-------------|
|     |      | Year                    | Growing       | Non-Growing   | Year            | Growing | Non-Growing |
| WSA | 1982 | 1.1 $\pm$ 0.4           | 1.3 $\pm$ 0.4 | 1.1 $\pm$ 0.4 | 44              | 25      | 19          |
| WSB |      | 1.3 $\pm$ 0.7           | 1.7 $\pm$ 0.7 | 1.2 $\pm$ 0.4 | 45              | 25      | 20          |
| WSC |      | 1.2 $\pm$ 0.6           | 1.5 $\pm$ 0.6 | 1.2 $\pm$ 0.3 | 42              | 22      | 20          |
| WSA | 1983 | 1.1 $\pm$ 0.6           | 1.1 $\pm$ 0.7 | 1.1 $\pm$ 0.5 | 47              | 22      | 25          |
| WSB |      | 1.2 $\pm$ 0.8           | 1.3 $\pm$ 1.0 | 1.2 $\pm$ 0.5 | 48              | 23      | 25          |
| WSC |      | 1.3 $\pm$ 3.0           | 1.3 $\pm$ 4.4 | 1.3 $\pm$ 0.1 | 42              | 17      | 25          |
| WSA | 1984 | 1.1 $\pm$ 0.8           | 1.1 $\pm$ 0.8 | 1.2 $\pm$ 0.6 | 52              | 26      | 26          |
| WSB |      | 1.7 $\pm$ 0.5           | 2.1 $\pm$ 0.5 | 1.6 $\pm$ 0.4 | 52              | 26      | 26          |
| WSC |      | 1.8 $\pm$ 0.5           | 1.6 $\pm$ 0.4 | 1.8 $\pm$ 0.1 | 52              | 26      | 26          |
| WSA | 1985 | 1.1 $\pm$ 0.5           | 1.6 $\pm$ 0.5 | 1.1 $\pm$ 0.4 | 48              | 23      | 25          |
| WSB |      | 1.7 $\pm$ 0.7           | 2.0 $\pm$ 0.9 | 1.6 $\pm$ 0.4 | 52              | 27      | 25          |
| WSC |      | 1.9 $\pm$ 0.6           | 2.1 $\pm$ 0.6 | 1.9 $\pm$ 0.2 | 52              | 27      | 25          |
| WSA | 1986 | 1.9 $\pm$ 0.6           | 1.8 $\pm$ 0.6 | 1.9 $\pm$ 0.6 | 48              | 24      | 24          |
| WSB |      | 2.3 $\pm$ 0.9           | 2.0 $\pm$ 0.6 | 2.3 $\pm$ 1.0 | 43              | 19      | 24          |
| WSC |      | 2.8 $\pm$ 1.1           | 2.9 $\pm$ 1.2 | 2.7 $\pm$ 0.1 | 50              | 25      | 25          |
| WSA | 1987 | 1.2 $\pm$ 0.7           | 1.4 $\pm$ 0.3 | 1.1 $\pm$ 0.9 | 44              | 21      | 23          |
| WSB |      | 1.4 $\pm$ 0.5           | 1.8 $\pm$ 0.2 | 1.3 $\pm$ 0.6 | 42              | 20      | 22          |
| WSC |      | 1.7 $\pm$ 0.8           | 2.0 $\pm$ 0.6 | 1.5 $\pm$ 0.3 | 46              | 24      | 22          |
| WSA | 1988 | 1.5 $\pm$ 0.7           | 1.5 $\pm$ 0.9 | 1.5 $\pm$ 0.4 | 41              | 16      | 25          |
| WSB |      | 1.6 $\pm$ 0.9           | 2.0 $\pm$ 1.2 | 1.5 $\pm$ 0.3 | 38              | 14      | 24          |
| WSC |      | 2.1 $\pm$ 1.1           | 2.6 $\pm$ 1.3 | 1.9 $\pm$ 0.4 | 43              | 18      | 25          |
| WSA | 1989 | 1.1 $\pm$ 0.3           | 1.1 $\pm$ 0.2 | 1.0 $\pm$ 0.3 | 52              | 27      | 25          |
| WSB |      | 1.3 $\pm$ 0.5           | 1.4 $\pm$ 0.2 | 1.3 $\pm$ 0.7 | 50              | 27      | 23          |
| WSC |      | 1.6 $\pm$ 0.5           | 1.8 $\pm$ 0.5 | 1.4 $\pm$ 0.2 | 51              | 27      | 24          |
| WSA | 1990 | 1.1 $\pm$ 0.4           | 1.2 $\pm$ 0.3 | 1.0 $\pm$ 0.5 | 52              | 27      | 25          |
| WSB |      | 1.3 $\pm$ 0.3           | 1.5 $\pm$ 0.3 | 1.3 $\pm$ 0.3 | 51              | 27      | 24          |
| WSC |      | 1.6 $\pm$ 0.5           | 1.7 $\pm$ 0.5 | 1.6 $\pm$ 0.1 | 51              | 27      | 24          |
| WSA | 1991 | 1.1 $\pm$ 0.4           | 1.1 $\pm$ 0.2 | 1.0 $\pm$ 0.5 | 52              | 27      | 25          |
| WSB |      | 1.5 $\pm$ 0.2           | 1.4 $\pm$ 0.1 | 1.5 $\pm$ 0.2 | 53              | 27      | 26          |
| WSC |      | 1.5 $\pm$ 0.2           | 1.6 $\pm$ 0.2 | 1.4 $\pm$ 0.0 | 35              | 19      | 16          |
| WSA | 1992 | 1.2 $\pm$ 0.4           | 1.1 $\pm$ 0.3 | 1.2 $\pm$ 0.5 | 52              | 27      | 25          |
| WSB |      | 1.3 $\pm$ 0.3           | 1.4 $\pm$ 0.3 | 1.2 $\pm$ 0.2 | 51              | 28      | 23          |
| WSC |      | 1.5 $\pm$ 0.3           | 1.5 $\pm$ 0.3 | 1.5 $\pm$ 0.1 | 51              | 28      | 23          |
| WSA | 1993 | 1.0 $\pm$ 0.4           | 1.2 $\pm$ 0.4 | 0.9 $\pm$ 0.2 | 44              | 27      | 17          |
| WSB |      | 1.3 $\pm$ 0.3           | 1.5 $\pm$ 0.4 | 1.2 $\pm$ 0.2 | 43              | 27      | 16          |
| WSC |      | 1.5 $\pm$ 0.4           | 1.8 $\pm$ 0.4 | 1.3 $\pm$ 0.1 | 43              | 27      | 16          |



Table B.8: Growing, non-growing, and yearly Potassium FVMC  $\pm$  Std. Dev. (2002-2008).

|     |      | K (mg L <sup>-1</sup> ) |               |               | Sample Size (n) |         |             |
|-----|------|-------------------------|---------------|---------------|-----------------|---------|-------------|
|     |      | Year                    | Growing       | Non-Growing   | Year            | Growing | Non-Growing |
| WSA | 2002 | 1.6 $\pm$ 0.7           | 1.7 $\pm$ 0.7 | 1.6 $\pm$ 0.4 | 47              | 23      | 24          |
| WSB |      | 1.7 $\pm$ 0.6           | 1.9 $\pm$ 0.6 | 1.7 $\pm$ 0.4 | 46              | 21      | 25          |
| WSC |      | 2.0 $\pm$ 0.8           | 2.1 $\pm$ 0.9 | 1.8 $\pm$ 0.1 | 48              | 24      | 24          |
| WSA | 2003 | 1.4 $\pm$ 0.5           | 1.6 $\pm$ 0.4 | 1.3 $\pm$ 0.4 | 51              | 27      | 24          |
| WSB |      | 1.5 $\pm$ 0.3           | 1.7 $\pm$ 0.3 | 1.4 $\pm$ 0.3 | 51              | 27      | 24          |
| WSC |      | 1.7 $\pm$ 0.5           | 1.8 $\pm$ 0.5 | 1.6 $\pm$ 0.2 | 51              | 27      | 24          |
| WSA | 2004 | 1.3 $\pm$ 0.2           | 1.4 $\pm$ 0.2 | 1.2 $\pm$ 0.1 | 51              | 28      | 23          |
| WSB |      | 1.5 $\pm$ 0.2           | 1.5 $\pm$ 0.2 | 1.4 $\pm$ 0.1 | 51              | 28      | 23          |
| WSC |      | 1.6 $\pm$ 0.2           | 1.6 $\pm$ 0.2 | 1.5 $\pm$ 0.3 | 51              | 28      | 23          |
| WSA | 2005 | 1.6 $\pm$ 0.5           | 1.9 $\pm$ 0.4 | 1.3 $\pm$ 0.4 | 43              | 22      | 21          |
| WSB |      | 1.7 $\pm$ 0.4           | 2.0 $\pm$ 0.4 | 1.4 $\pm$ 0.3 | 41              | 20      | 21          |
| WSC |      | 1.7 $\pm$ 0.6           | 2.1 $\pm$ 0.5 | 1.5 $\pm$ 0.2 | 40              | 19      | 21          |
| WSA | 2006 | 1.6 $\pm$ 0.4           | 2.0 $\pm$ 0.3 | 1.4 $\pm$ 0.4 | 48              | 24      | 24          |
| WSB |      | 1.6 $\pm$ 0.4           | 2.0 $\pm$ 0.3 | 1.4 $\pm$ 0.4 | 47              | 23      | 24          |
| WSC |      | 1.7 $\pm$ 0.4           | 2.0 $\pm$ 0.3 | 1.5 $\pm$ 0.2 | 49              | 25      | 24          |
| WSA | 2007 | 1.0 $\pm$ 0.6           | 1.2 $\pm$ 0.5 | 0.9 $\pm$ 0.6 | 35              | 14      | 21          |
| WSB |      | 0.9 $\pm$ 0.3           | 0.9 $\pm$ 0.3 | 0.9 $\pm$ 0.3 | 33              | 13      | 20          |
| WSC |      | 1.2 $\pm$ 0.8           | 1.0 $\pm$ 0.4 | 1.2 $\pm$ 0.1 | 36              | 14      | 22          |
| WSA | 2008 | 1.2 $\pm$ 0.6           | 1.4 $\pm$ 0.5 | 1.2 $\pm$ 0.6 | 35              | 13      | 22          |
| WSB |      | 1.1 $\pm$ 0.4           | 1.3 $\pm$ 0.4 | 1.1 $\pm$ 0.3 | 35              | 13      | 22          |
| WSC |      | 1.3 $\pm$ 0.4           | 1.5 $\pm$ 0.4 | 1.3 $\pm$ 0.3 | 36              | 14      | 22          |

Table B.9: Growing, non-growing, and yearly Sodium FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | Na (mg L <sup>-1</sup> ) |         |             | Sample Size (n) |         |             |
|-----|------|--------------------------|---------|-------------|-----------------|---------|-------------|
|     |      | Year                     | Growing | Non-Growing | Year            | Growing | Non-Growing |
| WSA | 1982 | 1.1±0.3                  | 1.2±0.2 | 1.1±0.3     | 44              | 25      | 19          |
| WSB |      | 1.3±0.4                  | 1.5±0.3 | 1.3±0.5     | 45              | 25      | 20          |
| WSC |      | 1.1±0.3                  | 1.3±0.3 | 1.1±0.3     | 42              | 22      | 20          |
| WSA | 1983 | 0.7±0.1                  | 0.6±0.1 | 0.8±0.1     | 47              | 22      | 25          |
| WSB |      | 1.0±0.2                  | 1.0±0.2 | 0.9±0.3     | 48              | 23      | 25          |
| WSC |      | 0.6±0.2                  | 0.6±0.4 | 0.8±0.1     | 42              | 17      | 25          |
| WSA | 1984 | 0.7±0.1                  | 0.7±0.1 | 0.7±0.1     | 52              | 26      | 26          |
| WSB |      | 0.8±0.2                  | 0.9±0.2 | 0.8±0.2     | 52              | 26      | 26          |
| WSC |      | 0.8±0.1                  | 0.8±0.2 | 0.8±0.1     | 52              | 26      | 26          |
| WSA | 1985 | 0.8±0.2                  | 0.9±0.1 | 0.8±0.2     | 48              | 23      | 25          |
| WSB |      | 1.1±0.4                  | 1.4±0.3 | 1.0±0.2     | 52              | 27      | 25          |
| WSC |      | 1.0±0.2                  | 1.1±0.1 | 1.0±0.2     | 52              | 27      | 25          |
| WSA | 1986 | 0.9±0.1                  | 1.0±0.1 | 0.8±0.1     | 48              | 24      | 24          |
| WSB |      | 1.0±0.4                  | 1.5±0.3 | 0.9±0.3     | 43              | 19      | 24          |
| WSC |      | 0.9±0.3                  | 1.3±0.3 | 0.8±0.1     | 50              | 25      | 25          |
| WSA | 1987 | 0.8±0.2                  | 0.9±0.2 | 0.8±0.2     | 44              | 21      | 23          |
| WSB |      | 0.9±0.4                  | 1.0±0.3 | 0.9±0.4     | 42              | 20      | 22          |
| WSC |      | 0.9±0.3                  | 1.1±0.4 | 0.8±0.3     | 46              | 24      | 22          |
| WSA | 1988 | 1.2±0.4                  | 1.3±0.4 | 1.2±0.3     | 41              | 16      | 25          |
| WSB |      | 1.4±0.5                  | 1.5±0.5 | 1.4±0.6     | 38              | 14      | 24          |
| WSC |      | 1.4±0.5                  | 1.8±0.5 | 1.3±0.4     | 43              | 18      | 25          |
| WSA | 1989 | 0.8±0.2                  | 0.8±0.2 | 0.9±0.2     | 52              | 27      | 25          |
| WSB |      | 1.0±0.5                  | 1.1±0.6 | 1.0±0.4     | 50              | 27      | 23          |
| WSC |      | 0.9±0.2                  | 0.8±0.2 | 0.9±0.2     | 51              | 27      | 24          |
| WSA | 1990 | 0.7±0.1                  | 0.8±0.1 | 0.7±0.1     | 52              | 27      | 25          |
| WSB |      | 0.8±0.2                  | 0.9±0.2 | 0.8±0.2     | 51              | 27      | 24          |
| WSC |      | 0.7±0.1                  | 0.8±0.1 | 0.7±0.1     | 51              | 27      | 24          |
| WSA | 1991 | 0.7±0.1                  | 0.8±0.1 | 0.7±0.1     | 52              | 27      | 25          |
| WSB |      | 0.8±0.2                  | 0.9±0.2 | 0.8±0.2     | 53              | 27      | 26          |
| WSC |      | 0.7±0.1                  | 0.7±0.1 | 0.7±0.0     | 35              | 19      | 16          |
| WSA | 1992 | 0.7±0.1                  | 0.7±0.2 | 0.7±0.1     | 52              | 27      | 25          |
| WSB |      | 0.8±0.2                  | 0.9±0.2 | 0.8±0.2     | 51              | 28      | 23          |
| WSC |      | 0.8±0.2                  | 0.8±0.2 | 0.7±0.1     | 51              | 28      | 23          |
| WSA | 1993 | 0.8±0.1                  | 0.9±0.1 | 0.7±0.1     | 44              | 27      | 17          |
| WSB |      | 0.9±0.2                  | 1.1±0.2 | 0.8±0.1     | 43              | 27      | 16          |
| WSC |      | 0.8±0.1                  | 0.9±0.1 | 0.7±0.1     | 43              | 27      | 16          |

Table B.10: Growing, non-growing, and yearly Sodium FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | Na (mg L <sup>-1</sup> ) |         |             | Sample Size (n) |         |             |
|-----|------|--------------------------|---------|-------------|-----------------|---------|-------------|
|     |      | Year                     | Growing | Non-Growing | Year            | Growing | Non-Growing |
| WSA | 2002 | 1.0±0.2                  | 1.1±0.2 | 0.9±0.1     | 47              | 23      | 24          |
| WSB |      | 1.1±0.3                  | 1.1±0.3 | 1.1±0.2     | 46              | 21      | 25          |
| WSC |      | 1.0±0.1                  | 1.0±0.1 | 1.0±0.1     | 48              | 24      | 24          |
| WSA | 2003 | 0.9±0.2                  | 1.1±0.1 | 0.9±0.2     | 51              | 27      | 24          |
| WSB |      | 1.0±0.3                  | 1.0±0.2 | 0.9±0.3     | 51              | 27      | 24          |
| WSC |      | 0.9±0.2                  | 1.0±0.1 | 0.9±0.2     | 49              | 27      | 24          |
| WSA | 2004 | 1.0±0.2                  | 0.9±0.1 | 1.2±0.3     | 51              | 28      | 23          |
| WSB |      | 1.0±0.2                  | 0.9±0.2 | 1.1±0.3     | 51              | 28      | 23          |
| WSC |      | 1.1±0.2                  | 1.0±0.1 | 1.2±0.3     | 51              | 28      | 23          |
| WSA | 2005 | 1.2±0.4                  | 1.4±0.4 | 1.0±0.2     | 43              | 22      | 21          |
| WSB |      | 1.1±0.3                  | 1.4±0.2 | 0.9±0.3     | 41              | 20      | 21          |
| WSC |      | 1.0±0.3                  | 1.3±0.1 | 0.9±0.2     | 40              | 19      | 21          |
| WSA | 2006 | 1.1±0.2                  | 1.2±0.2 | 1.1±0.2     | 48              | 24      | 24          |
| WSB |      | 1.3±0.2                  | 1.4±0.2 | 1.3±0.2     | 47              | 23      | 24          |
| WSC |      | 1.2±0.1                  | 1.2±0.1 | 1.2±0.2     | 49              | 25      | 24          |
| WSA | 2007 | 1.1±0.3                  | 1.2±0.3 | 1.1±0.2     | 35              | 14      | 21          |
| WSB |      | 1.1±0.3                  | 1.2±0.3 | 1.1±0.2     | 33              | 13      | 20          |
| WSC |      | 1.1±0.2                  | 1.1±0.2 | 1.1±0.1     | 36              | 14      | 22          |
| WSA | 2008 | 1.0±0.3                  | 1.2±0.1 | 0.9±0.2     | 35              | 13      | 22          |
| WSB |      | 1.0±0.3                  | 1.2±0.2 | 0.9±0.3     | 35              | 13      | 22          |
| WSC |      | 0.9±0.3                  | 1.1±0.2 | 0.9±0.3     | 36              | 14      | 22          |

Table B.11: Growing, non-growing, and yearly Alkalinity FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | ALK (mg L <sup>-1</sup> CaCO <sub>3</sub> ) |                 |                 | Sample Size (n) |         |             |
|-----|------|---|-----------------|-----------------|-----------------|---------|-------------|
|     |      | Year  | Growing         | Non-Growing     | Year            | Growing | Non-Growing |
| WSA | 1982 | 7.7 $\pm$ 4.5                               | 10.8 $\pm$ 3.6  | 7.3 $\pm$ 2.0   | 44              | 25      | 19          |
| WSB |      | 8.2 $\pm$ 6.2                               | 12.1 $\pm$ 5.6  | 7.5 $\pm$ 4.0   | 43              | 25      | 18          |
| WSC |      | 8.8 $\pm$ 5.6                               | 11.6 $\pm$ 5.9  | 8.3 $\pm$ 2.4   | 41              | 22      | 19          |
| WSA | 1983 | 12.2 $\pm$ 10.4                             | 13.7 $\pm$ 10.1 | 8.2 $\pm$ 6.0   | 47              | 22      | 25          |
| WSB |      | 13.2 $\pm$ 11.3                             | 14.0 $\pm$ 12.7 | 11.0 $\pm$ 6.4  | 47              | 23      | 24          |
| WSC |      | 15.0 $\pm$ 26.4                             | 16.1 $\pm$ 37.1 | 12.1 $\pm$ 7.9  | 41              | 17      | 24          |
| WSA | 1984 | 13.2 $\pm$ 9.9                              | 12.1 $\pm$ 10.5 | 15.6 $\pm$ 4.6  | 52              | 26      | 26          |
| WSB |      | 25.5 $\pm$ 8.0                              | 16.7 $\pm$ 7.4  | 28.0 $\pm$ 8.1  | 52              | 26      | 26          |
| WSC |      | 21.4 $\pm$ 7.9                              | 20.7 $\pm$ 7.7  | 21.8 $\pm$ 6.2  | 52              | 26      | 26          |
| WSA | 1985 | 22.6 $\pm$ 6.9                              | 27.4 $\pm$ 7.7  | 22.1 $\pm$ 5.0  | 48              | 23      | 25          |
| WSB |      | 25.2 $\pm$ 5.3                              | 25.4 $\pm$ 6.2  | 25.1 $\pm$ 4.0  | 52              | 27      | 25          |
| WSC |      | 26.4 $\pm$ 8.4                              | 30.5 $\pm$ 8.7  | 25.1 $\pm$ 5.9  | 52              | 27      | 25          |
| WSA | 1986 | 17.7 $\pm$ 4.9                              | 22.0 $\pm$ 4.0  | 17.1 $\pm$ 3.9  | 48              | 24      | 24          |
| WSB |      | 17.5 $\pm$ 5.7                              | 20.3 $\pm$ 5.2  | 17.0 $\pm$ 5.4  | 43              | 19      | 24          |
| WSC |      | 21.2 $\pm$ 10.5                             | 31.8 $\pm$ 9.8  | 18.8 $\pm$ 5.3  | 50              | 25      | 25          |
| WSA | 1987 | 14.1 $\pm$ 4.1                              | 15.9 $\pm$ 4.4  | 13.4 $\pm$ 1.2  | 44              | 21      | 23          |
| WSB |      | 14.8 $\pm$ 3.9                              | 16.7 $\pm$ 4.5  | 14.3 $\pm$ 2.2  | 42              | 20      | 22          |
| WSC |      | 16.3 $\pm$ 5.5                              | 20.6 $\pm$ 5.3  | 14.7 $\pm$ 2.2  | 46              | 24      | 22          |
| WSA | 1988 | 11.9 $\pm$ 6.3                              | 9.6 $\pm$ 5.3   | 12.7 $\pm$ 6.9  | 41              | 16      | 25          |
| WSB |      | 12.8 $\pm$ 6.8                              | 14.2 $\pm$ 6.8  | 12.3 $\pm$ 6.8  | 38              | 14      | 24          |
| WSC |      | 14.1 $\pm$ 8.9                              | 16.6 $\pm$ 10.4 | 13.1 $\pm$ 6.6  | 43              | 18      | 25          |
| WSA | 1989 | 19.5 $\pm$ 3.4                              | 20.0 $\pm$ 4.1  | 19.1 $\pm$ 2.4  | 52              | 27      | 25          |
| WSB |      | 19.4 $\pm$ 3.1                              | 20.1 $\pm$ 3.2  | 18.8 $\pm$ 2.8  | 50              | 27      | 23          |
| WSC |      | 21.4 $\pm$ 4.9                              | 23.3 $\pm$ 5.0  | 19.9 $\pm$ 3.5  | 51              | 27      | 24          |
| WSA | 1990 | 24.5 $\pm$ 10.6                             | 33.3 $\pm$ 8.6  | 20.8 $\pm$ 7.0  | 52              | 27      | 25          |
| WSB |      | 25.5 $\pm$ 11.0                             | 33.2 $\pm$ 8.2  | 23.0 $\pm$ 7.5  | 51              | 27      | 24          |
| WSC |      | 32.7 $\pm$ 19.1                             | 40.6 $\pm$ 17.7 | 29.5 $\pm$ 12.0 | 51              | 27      | 24          |
| WSA | 1991 | 34.1 $\pm$ 10.3                             | 32.5 $\pm$ 8.6  | 34.8 $\pm$ 11.8 | 52              | 27      | 25          |
| WSB |      | 44.1 $\pm$ 11.7                             | 34.9 $\pm$ 8.1  | 46.2 $\pm$ 14.5 | 53              | 27      | 26          |
| WSC |      | 31.8 $\pm$ 8.0                              | 37.5 $\pm$ 6.0  | 29.0 $\pm$ 5.9  | 35              | 19      | 16          |
| WSA | 1992 | 38.3 $\pm$ 21.9                             | 25.3 $\pm$ 3.9  | 46.3 $\pm$ 29.2 | 52              | 27      | 25          |
| WSB |      | 43.6 $\pm$ 20.8                             | 27.4 $\pm$ 3.8  | 50.8 $\pm$ 27.4 | 51              | 28      | 23          |
| WSC |      | 41.1 $\pm$ 17.9                             | 33.1 $\pm$ 5.9  | 45.7 $\pm$ 24.3 | 51              | 28      | 23          |
| WSA | 1993 | 13.3 $\pm$ 4.6                              | 15.5 $\pm$ 4.2  | 12.2 $\pm$ 1.6  | 44              | 27      | 17          |
| WSB |      | 16.2 $\pm$ 4.2                              | 20.1 $\pm$ 4.6  | 13.8 $\pm$ 1.6  | 42              | 26      | 16          |
| WSC |      | 18.9 $\pm$ 7.3                              | 26.3 $\pm$ 6.7  | 14.5 $\pm$ 2.9  | 43              | 27      | 16          |

Table B.12: Growing, non-growing, and yearly Alkalinity FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | ALK (mg L <sup>-1</sup> CaCO <sub>3</sub> ) |                 |                  | Sample Size (n) |         |             |
|-----|------|---|-----------------|------------------|-----------------|---------|-------------|
|     |      | Year  | Growing         | Non-Growing      | Year            | Growing | Non-Growing |
| WSA | 2002 | 28.3 $\pm$ 16.1                             | 31.1 $\pm$ 16.2 | 25.5 $\pm$ 9.7   | 47              | 23      | 24          |
| WSB |      | 27.6 $\pm$ 16.4                             | 29.7 $\pm$ 16.9 | 25.9 $\pm$ 9.1   | 46              | 21      | 25          |
| WSC |      | 32.5 $\pm$ 15.6                             | 35.6 $\pm$ 15.3 | 30.4 $\pm$ 11.7  | 48              | 24      | 24          |
| WSA | 2003 | 15.2 $\pm$ 14.6                             | 24.6 $\pm$ 15.3 | 10.6 $\pm$ 6.2   | 51              | 27      | 24          |
| WSB |      | 13.9 $\pm$ 12.1                             | 19.5 $\pm$ 12.2 | 10.4 $\pm$ 5.6   | 51              | 27      | 24          |
| WSC |      | 17.7 $\pm$ 14.8                             | 24.6 $\pm$ 15.2 | 13.9 $\pm$ 9.7   | 49              | 27      | 24          |
| WSA | 2004 | 13.8 $\pm$ 13.7                             | 17.4 $\pm$ 15.3 | 9.9 $\pm$ 1.6    | 51              | 28      | 23          |
| WSB |      | 13.5 $\pm$ 8.6                              | 15.7 $\pm$ 9.0  | 10.5 $\pm$ 1.6   | 51              | 28      | 23          |
| WSC |      | 34.0 $\pm$ 191.9                            | 22.3 $\pm$ 7.7  | 46.7 $\pm$ 280.1 | 51              | 28      | 23          |
| WSA | 2005 | 19.5 $\pm$ 23.8                             | 27.0 $\pm$ 26.5 | 12.9 $\pm$ 10.2  | 43              | 22      | 21          |
| WSB |      | 15.3 $\pm$ 12.3                             | 20.6 $\pm$ 11.0 | 11.5 $\pm$ 9.2   | 41              | 20      | 21          |
| WSC |      | 17.3 $\pm$ 14.5                             | 28.0 $\pm$ 14.9 | 12.8 $\pm$ 5.4   | 40              | 19      | 21          |
| WSA | 2006 | 11.6 $\pm$ 5.2                              | 13.0 $\pm$ 5.9  | 10.9 $\pm$ 2.4   | 48              | 24      | 24          |
| WSB |      | 12.2 $\pm$ 5.4                              | 15.4 $\pm$ 6.1  | 10.8 $\pm$ 2.0   | 47              | 23      | 24          |
| WSC |      | 13.1 $\pm$ 6.9                              | 14.9 $\pm$ 8.0  | 12.2 $\pm$ 3.1   | 49              | 25      | 24          |
| WSA | 2007 | 13.6 $\pm$ 18.6                             | 21.2 $\pm$ 23.0 | 10.8 $\pm$ 5.5   | 35              | 14      | 21          |
| WSB |      | 12.3 $\pm$ 20.1                             | 16.7 $\pm$ 25.4 | 11.1 $\pm$ 3.4   | 33              | 13      | 20          |
| WSC |      | 18.6 $\pm$ 18.0                             | 28.1 $\pm$ 21.2 | 15.1 $\pm$ 10.4  | 36              | 14      | 22          |
| WSA | 2008 | 18.0 $\pm$ 15.5                             | 30.1 $\pm$ 16.3 | 14.9 $\pm$ 9.4   | 35              | 13      | 22          |
| WSB |      | 14.0 $\pm$ 16.7                             | 19.7 $\pm$ 16.7 | 12.5 $\pm$ 14.3  | 35              | 13      | 22          |
| WSC |      | 15.4 $\pm$ 12.8                             | 19.9 $\pm$ 9.8  | 13.7 $\pm$ 13.3  | 36              | 14      | 22          |

Table B.13: Growing, non-growing, and yearly Nitrate FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | NO <sub>3</sub> (mg L <sup>-1</sup> ) |                |                | Sample Size (n) |         |             |
|-----|------|---------------------------------------|----------------|----------------|-----------------|---------|-------------|
|     |      | Year                                  | Growing        | Non-Growing    | Year            | Growing | Non-Growing |
| WSA | 1982 | 0.1 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.1 $\pm$ 0.1  | 44              | 25      | 19          |
| WSB |      | 0.1 $\pm$ 0.2                         | 0.2 $\pm$ 0.2  | 0.1 $\pm$ 0.1  | 45              | 25      | 20          |
| WSC |      | 0.1 $\pm$ 0.04                        | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.0  | 42              | 22      | 20          |
| WSA | 1983 | 0.1 $\pm$ 0.2                         | 0.0 $\pm$ 0.2  | 0.2 $\pm$ 0.2  | 47              | 22      | 25          |
| WSB |      | 0.1 $\pm$ 1.7                         | 0.0 $\pm$ 1.8  | 0.4 $\pm$ 1.5  | 48              | 23      | 25          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.1  | 42              | 17      | 25          |
| WSA | 1984 | 0.2 $\pm$ 0.2                         | 0.1 $\pm$ 0.3  | 0.3 $\pm$ 0.2  | 52              | 26      | 26          |
| WSB |      | 1.2 $\pm$ 0.6                         | 0.4 $\pm$ 0.4  | 1.5 $\pm$ 0.6  | 52              | 26      | 26          |
| WSC |      | 1.3 $\pm$ 0.8                         | 0.3 $\pm$ 0.4  | 1.7 $\pm$ 0.9  | 52              | 26      | 26          |
| WSA | 1985 | 0.2 $\pm$ 0.1                         | 0.4 $\pm$ 0.1  | 0.2 $\pm$ 0.1  | 48              | 23      | 25          |
| WSB |      | 0.9 $\pm$ 0.3                         | 0.6 $\pm$ 0.2  | 1.0 $\pm$ 0.2  | 52              | 27      | 25          |
| WSC |      | 1.1 $\pm$ 0.4                         | 0.6 $\pm$ 0.3  | 1.3 $\pm$ 0.2  | 52              | 27      | 25          |
| WSA | 1986 | 0.1 $\pm$ 0.1                         | 0.2 $\pm$ 0.2  | 0.1 $\pm$ 0.1  | 48              | 24      | 24          |
| WSB |      | 0.6 $\pm$ 0.2                         | 0.3 $\pm$ 0.1  | 0.7 $\pm$ 0.3  | 43              | 19      | 24          |
| WSC |      | 0.8 $\pm$ 0.4                         | 0.1 $\pm$ 0.1  | 1.0 $\pm$ 0.3  | 50              | 25      | 25          |
| WSA | 1987 | 0.1 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.0 $\pm$ 0.1  | 44              | 21      | 23          |
| WSB |      | 0.2 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.2 $\pm$ 0.1  | 42              | 20      | 22          |
| WSC |      | 0.4 $\pm$ 0.2                         | 0.2 $\pm$ 0.1  | 0.5 $\pm$ 0.2  | 46              | 24      | 22          |
| WSA | 1988 | 0.2 $\pm$ 0.3                         | 0.2 $\pm$ 0.4  | 0.1 $\pm$ 0.1  | 41              | 16      | 25          |
| WSB |      | 0.2 $\pm$ 0.3                         | 0.3 $\pm$ 0.4  | 0.1 $\pm$ 0.1  | 38              | 14      | 24          |
| WSC |      | 0.3 $\pm$ 0.2                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.2  | 43              | 18      | 25          |
| WSA | 1989 | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.0  | 52              | 27      | 25          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03 | 0.1 $\pm$ 0.04 | 50              | 27      | 23          |
| WSC |      | 0.1 $\pm$ 0.05                        | 0.1 $\pm$ 0.03 | 0.1 $\pm$ 0.0  | 51              | 27      | 24          |
| WSA | 1990 | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.1 $\pm$ 0.1  | 52              | 27      | 25          |
| WSB |      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.2 $\pm$ 0.1  | 50              | 27      | 23          |
| WSC |      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.1  | 51              | 27      | 24          |
| WSA | 1991 | 1.0 $\pm$ 1.0                         | 1.2 $\pm$ 0.7  | 1.0 $\pm$ 1.2  | 52              | 27      | 25          |
| WSB |      | 3.0 $\pm$ 1.0                         | 1.3 $\pm$ 0.4  | 3.4 $\pm$ 1.4  | 53              | 27      | 26          |
| WSC |      | 1.6 $\pm$ 0.8                         | 1.5 $\pm$ 0.6  | 1.7 $\pm$ 0.9  | 35              | 19      | 16          |
| WSA | 1992 | 1.3 $\pm$ 0.6                         | 1.6 $\pm$ 0.5  | 1.1 $\pm$ 0.5  | 48              | 27      | 21          |
| WSB |      | 2.0 $\pm$ 0.8                         | 1.3 $\pm$ 0.5  | 2.4 $\pm$ 0.8  | 47              | 27      | 20          |
| WSC |      | 2.2 $\pm$ 1.2                         | 1.3 $\pm$ 0.7  | 2.8 $\pm$ 1.4  | 50              | 28      | 22          |
| WSA | 1993 | 0.2 $\pm$ 0.4                         | 0.3 $\pm$ 0.4  | 0.2 $\pm$ 0.2  | 35              | 18      | 17          |
| WSB |      | 0.4 $\pm$ 0.2                         | 0.5 $\pm$ 0.3  | 0.3 $\pm$ 0.1  | 34              | 18      | 16          |
| WSC |      | 0.5 $\pm$ 0.2                         | 0.5 $\pm$ 0.1  | 0.5 $\pm$ 0.2  | 34              | 18      | 16          |

Table B.14: Growing, non-growing, and yearly Nitrate FWMC  $\pm$  Std. Dev. (2002-2008).

|          | NO <sub>3</sub> (mg L <sup>-1</sup> ) |                |               | Sample Size (n) |         |             |
|----------|---------------------------------------|----------------|---------------|-----------------|---------|-------------|
|          | Year                                  | Growing        | Non-Growing   | Year            | Growing | Non-Growing |
| WSA      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.1 | 47              | 23      | 24          |
| WSB 2002 | 0.3 $\pm$ 0.2                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.2 | 46              | 21      | 25          |
| WSC      | 0.3 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.1 | 48              | 24      | 24          |
| WSA      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.1 | 51              | 27      | 24          |
| WSB 2003 | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.1 | 51              | 27      | 24          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.1 | 49              | 25      | 24          |
| WSA      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.1 $\pm$ 0.0 | 51              | 28      | 23          |
| WSB 2004 | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.2 $\pm$ 0.1 | 51              | 28      | 23          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.1 $\pm$ 0.0  | 0.2 $\pm$ 0.1 | 51              | 28      | 23          |
| WSA      | 0.2 $\pm$ 0.3                         | 0.2 $\pm$ 0.2  | 0.1 $\pm$ 0.3 | 43              | 22      | 21          |
| WSB 2005 | 0.3 $\pm$ 0.2                         | 0.2 $\pm$ 0.1  | 0.3 $\pm$ 0.3 | 41              | 20      | 21          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1  | 0.2 $\pm$ 0.1 | 40              | 19      | 21          |
| WSA      | 0.1 $\pm$ 0.2                         | 0.2 $\pm$ 0.2  | 0.1 $\pm$ 0.1 | 48              | 24      | 24          |
| WSB 2006 | 0.2 $\pm$ 0.1                         | 0.3 $\pm$ 0.1  | 0.2 $\pm$ 0.2 | 47              | 23      | 24          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.2 $\pm$ 0.1 | 49              | 25      | 24          |
| WSA      | 0.2 $\pm$ 0.2                         | 0.2 $\pm$ 0.2  | 0.2 $\pm$ 0.2 | 35              | 14      | 21          |
| WSB 2007 | 0.3 $\pm$ 0.2                         | 0.2 $\pm$ 0.1  | 0.4 $\pm$ 0.2 | 33              | 13      | 20          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.1 $\pm$ 0.1  | 0.3 $\pm$ 0.2 | 36              | 14      | 22          |
| WSA      | 0.2 $\pm$ 0.4                         | 0.1 $\pm$ 0.1  | 0.2 $\pm$ 0.5 | 35              | 13      | 22          |
| WSB 2008 | 0.2 $\pm$ 0.5                         | 0.2 $\pm$ 0.1  | 0.2 $\pm$ 0.5 | 35              | 13      | 22          |
| WSC      | 0.2 $\pm$ 0.1                         | 0.1 $\pm$ 0.05 | 0.2 $\pm$ 0.1 | 36              | 14      | 22          |

Table B.15: Growing, non-growing, and yearly Phosphate FWMC  $\pm$  Std. Dev. (1982-1993).

|     |      | PO <sub>4</sub> (mg L <sup>-1</sup> ) |                 |                 | Sample Size (n) |         |             |
|-----|------|---------------------------------------|-----------------|-----------------|-----------------|---------|-------------|
|     |      | Year                                  | Growing         | Non-Growing     | Year            | Growing | Non-Growing |
| WSA | 1982 | 0.0 $\pm$ 0.1                         | 0.05 $\pm$ 0.1  | 0.01 $\pm$ 0.03 | 44              | 25      | 19          |
| WSB |      | 0.0 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.01 $\pm$ 0.04 | 45              | 25      | 20          |
| WSC |      | 0.0 $\pm$ 0.1                         | 0.03 $\pm$ 0.2  | 0.01 $\pm$ 0.1  | 43              | 22      | 21          |
| WSA | 1983 | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.04  | 47              | 22      | 25          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.05 $\pm$ 0.02 | 0.07 $\pm$ 0.02 | 48              | 23      | 25          |
| WSC |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.03  | 42              | 17      | 25          |
| WSA | 1984 | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.02  | 52              | 26      | 26          |
| WSB |      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1   | 0.2 $\pm$ 0.05  | 52              | 26      | 26          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.2 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 52              | 26      | 26          |
| WSA | 1985 | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.05  | 48              | 23      | 25          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.04  | 0.1 $\pm$ 0.05  | 52              | 27      | 25          |
| WSC |      | 0.2 $\pm$ 0.1                         | 0.2 $\pm$ 0.1   | 0.2 $\pm$ 0.1   | 52              | 27      | 25          |
| WSA | 1986 | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.05  | 48              | 24      | 24          |
| WSB |      | 0.1 $\pm$ 0.1                         | 0.2 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 43              | 19      | 24          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 50              | 25      | 25          |
| WSA | 1987 | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.08  | 44              | 21      | 23          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.05  | 42              | 20      | 22          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.04  | 0.1 $\pm$ 0.1   | 46              | 24      | 22          |
| WSA | 1988 | 0.2 $\pm$ 0.2                         | 0.2 $\pm$ 0.2   | 0.2 $\pm$ 0.2   | 41              | 16      | 25          |
| WSB |      | 0.3 $\pm$ 0.3                         | 0.3 $\pm$ 0.4   | 0.3 $\pm$ 0.3   | 38              | 14      | 24          |
| WSC |      | 0.3 $\pm$ 0.2                         | 0.3 $\pm$ 0.2   | 0.3 $\pm$ 0.2   | 43              | 18      | 25          |
| WSA | 1989 | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 52              | 27      | 25          |
| WSB |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 50              | 27      | 23          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 51              | 27      | 24          |
| WSA | 1990 | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.04  | 52              | 27      | 25          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.06  | 51              | 27      | 24          |
| WSC |      | 0.1 $\pm$ 0.04                        | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.04  | 51              | 27      | 24          |
| WSA | 1991 | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.03  | 52              | 27      | 25          |
| WSB |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.08  | 53              | 27      | 26          |
| WSC |      | 0.1 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.1 $\pm$ 0.03  | 35              | 19      | 16          |
| WSA | 1992 | 0.0 $\pm$ 0.1                         | 0.1 $\pm$ 0.1   | 0.04 $\pm$ 0.02 | 52              | 27      | 25          |
| WSB |      | 0.0 $\pm$ 0.0                         | 0.04 $\pm$ 0.01 | 0.03 $\pm$ 0.01 | 51              | 28      | 23          |
| WSC |      | 0.0 $\pm$ 0.01                        | 0.04 $\pm$ 0.01 | 0.03 $\pm$ 0.01 | 51              | 28      | 23          |
| WSA | 1993 | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.02  | 44              | 27      | 17          |
| WSB |      | 0.1 $\pm$ 0.0                         | 0.1 $\pm$ 0.03  | 0.1 $\pm$ 0.02  | 42              | 26      | 16          |
| WSC |      | 0.1 $\pm$ 0.03                        | 0.1 $\pm$ 0.04  | 0.1 $\pm$ 0.02  | 43              | 27      | 16          |



Table B.16: Growing, non-growing, and yearly Chlorine FWMC Std. Dev. (2002-2008).

|     |      | Cl (mg L <sup>-1</sup> ) |          |             | Sample Size (n) |         |             |
|-----|------|--------------------------|----------|-------------|-----------------|---------|-------------|
|     |      | Year                     | Growing  | Non-Growing | Year            | Growing | Non-Growing |
| WSA | 2002 | 0.8±0.1                  | 0.8±0.1  | 0.8±0.1     | 47              | 23      | 24          |
| WSB |      | 0.8±0.1                  | 0.8±0.1  | 0.8±0.1     | 46              | 21      | 25          |
| WSC |      | 0.7±0.1                  | 0.8±0.1  | 0.7±0.1     | 48              | 24      | 24          |
| WSA | 2003 | 0.9±0.6                  | 1.2±0.7  | 0.7±0.2     | 51              | 25      | 24          |
| WSB |      | 0.9±0.8                  | 1.2±1.0  | 0.7±0.2     | 51              | 25      | 24          |
| WSC |      | 0.8±0.5                  | 1.1±0.6  | 0.7±0.1     | 49              | 25      | 24          |
| WSA | 2004 | 0.6±0.1                  | 0.6±0.1  | 0.6±0.1     | 51              | 28      | 23          |
| WSB |      | 0.6±0.5                  | 0.6±0.7  | 0.6±0.1     | 51              | 28      | 23          |
| WSC |      | 0.6±0.1                  | 0.6±0.1  | 0.6±0.1     | 51              | 28      | 23          |
| WSA | 2005 | 0.5±0.1                  | 0.5±0.1  | 0.6±0.1     | 43              | 22      | 21          |
| WSB |      | 0.5±0.1                  | 0.4±0.1  | 0.6±0.1     | 41              | 20      | 21          |
| WSC |      | 0.5±0.2                  | 0.4±0.1  | 0.5±0.2     | 40              | 19      | 21          |
| WSA | 2006 | 1.2±0.5                  | 0.8±0.5  | 1.4±0.5     | 48              | 24      | 24          |
| WSB |      | 1.2±0.6                  | 0.7±0.5  | 1.4±0.6     | 47              | 23      | 24          |
| WSC |      | 1.0±0.5                  | 0.9±0.5  | 1.0±0.6     | 49              | 25      | 24          |
| WSA | 2007 | 1.0±0.4                  | 1.1±0.4  | 1.0±0.3     | 35              | 14      | 21          |
| WSB |      | 0.9±0.4                  | 1.1±0.5  | 0.9±0.3     | 33              | 13      | 20          |
| WSC |      | 1.0±0.7                  | 0.9±0.4  | 1.1±0.8     | 36              | 14      | 22          |
| WSA | 2008 | 1.1±0.4                  | 0.8±0.3  | 1.1±0.4     | 35              | 13      | 22          |
| WSB |      | 1.0±0.3                  | 0.9±0.2  | 1.0±0.3     | 35              | 13      | 22          |
| WSC |      | 0.9±0.3                  | 0.9±0.27 | 1.0±0.3     | 36              | 14      | 22          |

Table B.17: Growing, non-growing, and yearly Ammonia FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | NH <sub>4</sub> -N (mg L <sup>-1</sup> ) |                 |                 | Sample Size (n) |         |             |
|-----|------|--|-----------------|-----------------|-----------------|---------|-------------|
|     |      | Year                                     | Growing         | Non-Growing     | Year            | Growing | Non-Growing |
| WSA | 2002 | 0.04 $\pm$ 0.04                          | 0.05 $\pm$ 0.05 | 0.03 $\pm$ 0.02 | 37              | 23      | 14          |
| WSB |      | 0.03 $\pm$ 0.05                          | 0.04 $\pm$ 0.05 | 0.02 $\pm$ 0.03 | 36              | 21      | 15          |
| WSC |      | 0.03 $\pm$ 0.05                          | 0.04 $\pm$ 0.05 | 0.02 $\pm$ 0.03 | 38              | 24      | 14          |
| WSA | 2003 | 0.02 $\pm$ 0.02                          | 0.01 $\pm$ 0.01 | 0.03 $\pm$ 0.02 | 51              | 27      | 24          |
| WSB |      | 0.02 $\pm$ 0.02                          | 0.01 $\pm$ 0.03 | 0.03 $\pm$ 0.02 | 51              | 27      | 24          |
| WSC |      | 0.02 $\pm$ 0.02                          | 0.02 $\pm$ 0.02 | 0.03 $\pm$ 0.02 | 51              | 27      | 24          |
| WSA | 2004 | 0.02 $\pm$ 0.02                          | 0.03 $\pm$ 0.02 | 0.01 $\pm$ 0.02 | 38              | 20      | 18          |
| WSB |      | 0.05 $\pm$ 0.02                          | 0.1 $\pm$ 0.02  | 0.04 $\pm$ 0.02 | 28              | 17      | 11          |
| WSC |      | 0.03 $\pm$ 0.04                          | 0.03 $\pm$ 0.04 | 0.02 $\pm$ 0.02 | 32              | 21      | 11          |
| WSA | 2005 | 0.03 $\pm$ 0.02                          | 0.04 $\pm$ 0.02 | 0.03 $\pm$ 0.01 | 14              | 7       | 7           |
| WSB |      | 0.03 $\pm$ 0.01                          | 0.04 $\pm$ 0.01 | 0.02 $\pm$ 0.01 | 16              | 9       | 7           |
| WSC |      | 0.02 $\pm$ 0.01                          | 0.03 $\pm$ 0.01 | 0.02 $\pm$ 0.02 | 14              | 6       | 8           |
| WSA | 2006 | 0.05 $\pm$ 0.1                           | 0.04 $\pm$ 0.1  | 0.05 $\pm$ 0.04 | 20              | 10      | 10          |
| WSB |      | 0.05 $\pm$ 0.03                          | 0.04 $\pm$ 0.04 | 0.05 $\pm$ 0.02 | 20              | 9       | 11          |
| WSC |      | 0.04 $\pm$ 0.03                          | 0.1 $\pm$ 0.03  | 0.04 $\pm$ 0.01 | 20              | 10      | 10          |
| WSA | 2007 | 0.1 $\pm$ 0.1                            | 0.2 $\pm$ 0.1   | 0.1 $\pm$ 0.1   | 19              | 6       | 13          |
| WSB |      | 0.1 $\pm$ 0.04                           | 0.1 $\pm$ 0.02  | 0.1 $\pm$ 0.04  | 18              | 6       | 12          |
| WSC |      | 0.1 $\pm$ 0.1                            | 0.1 $\pm$ 0.04  | 0.1 $\pm$ 0.1   | 23              | 6       | 17          |
| WSA | 2008 | 0.1 $\pm$ 0.04                           | 0.1 $\pm$ 0.05  | 0.1 $\pm$ 0.02  | 7               | 2       | 5           |
| WSB |      | 0.04 $\pm$ 0.1                           | 0.03 $\pm$ 0.0  | 0.07 $\pm$ 0.1  | 6               | 2       | 4           |
| WSC |      | 0.1 $\pm$ 0.04                           | 0.1 $\pm$ 0.0   | 0.1 $\pm$ 0.04  | 11              | 1       | 10          |

Table B.18: Growing, non-growing, and yearly TOC FWMC  $\pm$  Std. Dev. (2002-2008).

|     |      | TOC (mg L <sup>-1</sup> ) |                |                | Sample Size (n) |         |             |
|-----|------|---------------------------|----------------|----------------|-----------------|---------|-------------|
|     |      | Year                      | Growing        | Non-Growing    | Year            | Growing | Non-Growing |
| WSA | 2002 | 4.3 $\pm$ 3.7             | 4.4 $\pm$ 4.4  | 4.3 $\pm$ 1.2  | 47              | 23      | 24          |
| WSB |      | 3.8 $\pm$ 2.8             | 3.7 $\pm$ 3.3  | 3.9 $\pm$ 1.1  | 46              | 21      | 25          |
| WSC |      | 5.2 $\pm$ 4.1             | 5.3 $\pm$ 5.0  | 5.2 $\pm$ 1.4  | 48              | 24      | 24          |
| WSA | 2003 | 3.4 $\pm$ 2.1             | 4.6 $\pm$ 2.1  | 2.8 $\pm$ 1.0  | 51              | 27      | 24          |
| WSB |      | 3.0 $\pm$ 1.4             | 3.6 $\pm$ 1.5  | 2.7 $\pm$ 0.6  | 51              | 27      | 24          |
| WSC |      | 3.8 $\pm$ 2.3             | 4.7 $\pm$ 2.5  | 3.3 $\pm$ 1.3  | 51              | 27      | 24          |
| WSA | 2004 | 3.5 $\pm$ 1.6             | 4.3 $\pm$ 1.6  | 2.6 $\pm$ 0.5  | 51              | 28      | 23          |
| WSB |      | 3.4 $\pm$ 1.4             | 4.0 $\pm$ 1.3  | 2.5 $\pm$ 0.7  | 51              | 28      | 23          |
| WSC |      | 3.9 $\pm$ 1.6             | 4.8 $\pm$ 1.6  | 2.9 $\pm$ 0.7  | 51              | 28      | 23          |
| WSA | 2005 | 3.9 $\pm$ 2.9             | 4.6 $\pm$ 3.1  | 3.3 $\pm$ 2.1  | 43              | 22      | 21          |
| WSB |      | 3.0 $\pm$ 1.6             | 3.5 $\pm$ 1.4  | 2.7 $\pm$ 1.4  | 41              | 20      | 21          |
| WSC |      | 3.4 $\pm$ 2.4             | 4.2 $\pm$ 1.8  | 3.0 $\pm$ 2.7  | 40              | 19      | 21          |
| WSA | 2006 | 3.8 $\pm$ 1.6             | 4.9 $\pm$ 1.3  | 3.2 $\pm$ 1.7  | 46              | 24      | 22          |
| WSB |      | 3.2 $\pm$ 1.3             | 4.0 $\pm$ 1.1  | 2.8 $\pm$ 1.5  | 45              | 23      | 22          |
| WSC |      | 3.6 $\pm$ 1.4             | 4.2 $\pm$ 1.4  | 3.3 $\pm$ 1.2  | 47              | 25      | 22          |
| WSA | 2007 | 4.0 $\pm$ 2.9             | 4.2 $\pm$ 2.1  | 3.9 $\pm$ 3.4  | 35              | 14      | 21          |
| WSB |      | 4.6 $\pm$ 2.5             | 4.8 $\pm$ 1.7  | 4.6 $\pm$ 2.8  | 29              | 12      | 17          |
| WSC |      | 4.4 $\pm$ 2.2             | 5.1 $\pm$ 1.6  | 4.1 $\pm$ 2.6  | 30              | 14      | 16          |
| WSA | 2008 | 0.4 $\pm$ 1.3             | 0.5 $\pm$ 1.6  | 0.4 $\pm$ 1.0  | 35              | 13      | 22          |
| WSB |      | 2.1 $\pm$ 0.7             | 2.0 $\pm$ 0.4  | 2.2 $\pm$ 0.8  | 8               | 5       | 3           |
| WSC |      | 3.7 $\pm$ 0.6             | 2.9 $\pm$ 0.31 | 4.2 $\pm$ 0.01 | 7               | 5       | 2           |

## REFERENCES

- Adams, M., Edwards, P., Kochenderfer, J., & Wood, F. (2004). Fifty years of watershed research on the Fernow Experimental Forest, West Virginia: Effects of forest management and air pollution on hardwood forests.
- Adams, T. O., Hook, D. D., & Floyd, M. A. (1995). Effectiveness monitoring of silvicultural best management practices in South Carolina. *Southern Journal of Applied Forestry*, 19(4), 170-176.
- United States Environmental Protection Agency Agency (USEPA). (2013). Water: Rivers & Streams.
- United States Environmental Protection Agency (USEPA). (2015). *Clean Water Rule: Definition of "Waters of the United States"*. Federal Register.
- National Council for Air and Stream Improvement (NCASI). (1999a). *Assessing effects of timber harvest on riparian zone features and functions for aquatic and wildlife habitat*.
- National Council for Air and Stream Improvement (NCASI). (1999b). *Scale Considerations and the Detectability of Sedimentary Cumulative Watershed Effects*.
- Aneja, V. P., Nelson, D. R., Roelle, P. A., Walker, J. T., & Battye, W. (2003). Agricultural ammonia emissions and ammonium concentrations associated with aerosols and precipitation in the southeast United States. *Journal of Geophysical Research: Atmospheres*, 108(D4).
- Arthur, M., Coltharp, G., & Brown, D. (1998). Effects of Best Management Practices on Forest Streamwater Quality in Eastern Kentucky: Wiley Online Library.
- Aubertin, G., & Patric, J. (1974). Water quality after clearcutting a small watershed in West Virginia. *Journal of Environmental Quality*, 3(3), 243-249.
- Aust, W. M., Carroll, M. B., Bolding, M. C., & Dolloff, C. A. (2011). Operational forest stream crossings effects on water quality in the Virginia Piedmont. *Southern Journal of Applied Forestry*, 35(3), 123-130.
- Barton, C., Karathanasis, A., & Chalfant, G. (2002). Influence of acidic

- atmospheric deposition on soil solution composition in the Daniel Boone National Forest, Kentucky, USA. *Environmental Geology*, 41(6), 672-682.
- BAUER, S. W. (1974). A modified Horton equation for infiltration during intermittent rainfall. *Hydrological Sciences Journal*, 19(2), 219-225.
- Benda, L., Hassan, M. A., Church, M., & May, C. L. (2005). Geomorphology Of steep-land headwaters: The transition from hillslopes to channels. *Journal of the American Water Resources Association*, 41(4), 835.
- Binkley, D., & Brown, T. C. (1993). Forest practices as nonpoint sources of pollution in North America. *Water resources bulletin*, 29(5), 729-740.
- Binkley, D., & Fisher, R. (2012). *Ecology and management of forest soils*: John Wiley & Sons.
- Binkley, D., Son, Y., & Valentine, D. W. (2000). Do forests receive occult inputs of nitrogen. *Ecosystems*, 3(4), 321-331.
- Blackburn-Lynch, W. C. (2015). *Development of techniques for assessing and restoring streams on surface mined lands*. Thesis (Ph. D.)--University of Kentucky, 2015.
- Boggs, J. (2016). Effects of timber harvest on water quantity and quality in small watersheds in the piedmont of North Carolina. *Journal of Forestry*, 114(1), 27-40.
- Bormann, F. H., & Likens, G. E. (1979). Catastrophic disturbance and the steady state in northern hardwood forests: A new look at the role of disturbance in the development of forest ecosystems suggests important implications for land-use policies. *American Scientist*, 67(6), 660-669.
- Bosch, J. M., & Hewlett, J. D. (1982). A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration. *Journal of Hydrology*, 55(1), 3-23.
- Brooks, K. N., Ffolliott, P. F., & Magner, J. A. (2012). *Hydrology and the management of watersheds*: John Wiley & Sons.
- Brooks, R. T., & Colburn, E. A. (2011). Extent and channel morphology of unmapped headwater stream segments of the Quabbin Watershed, Massachusetts: Wiley Online Library.

- Burns, D. A., & Murdoch, P. S. (2005). Effects of a clearcut on the net rates of nitrification and N-mineralization in a northern hardwood forest, Catskill Mountains, New York, USA. *Biogeochemistry*, 72(1), 123-146.
- Burt, T., Howden, N., McDonnell, J., Jones, J., & Hancock, G. (2015). Seeing the climate through the trees: observing climate and forestry impacts on streamflow using a 60- year record. *Hydrological processes*, 29(3), 473-480.
- Carr, A. E., Loague, K., & VanderKwaak, J. E. (2014). Hydrologic- response simulations for the North Fork of Caspar Creek: second- growth, clear-cut, new- growth, and cumulative watershed effect scenarios. *Hydrological processes*, 28(3), 1476-1494.
- Chamuris, G. P. (2016). Hiker's Guide to the Trees, Shrubs, and Woody Vines of Ricketts Glen State Park.
- Chen, Y., & Lin, L.S. (2009). Responses of streams in central Appalachian Mountain region to reduced acidic deposition—comparisons with other regions in North America and Europe. *Science of the total environment*, 407(7), 2285-2295.
- Cherry, M. A. (2006). *Hydrochemical characterization of ten headwater catchments in eastern Kentucky*: University of Kentucky.
- Clark, G. M., Mueller, D. K., & Mast, M. A. (2000). Nutrient concentrations and yields in undeveloped stream basins of the United States: Wiley Online Library.
- Collins, R. F. (1976). *A History of the Daniel Boone National Forest, 1770-1970*. Washington: U.S. Dept. of Agriculture, Forest Service, Southern Region.
- Colson, T., Gregory, J., Dorney, J., & Russell, P. (2008). Topographic and soil maps do not accurately depict headwater stream networks. *National Wetlands Newsletter*, 30(3), 25-28.
- Cristan, R., Aust, W. M., Bolding, M. C., Barrett, S. M., Munsell, J. F., & Schilling, E. (2016). Effectiveness of forestry best management practices in the United States: Literature review. *Forest Ecology and Management*, 360, 133-151.

- Dissmeyer, G. E. (1994). Evaluating the effectiveness of forestry best management practices in meeting water quality goals or standards. Miscellaneous pub: Forest Service, Atlanta, GA (United States). Southern Region.
- Douglass, J. E. (1983). The potential for water yield augmentation from forest management in the eastern United States: Wiley Online Library.
- Douglass, J. E., & Swank, W. T. (1972). Streamflow modification through management of eastern forests.
- Douglass, J. E., & Swank, W. T. (1976). *Multiple use in southern Appalachian hardwoods—a 10-year case history*. Paper presented at the proceedings of the 16th International Union of Forestry Research Organizations world congress.
- Eckhardt, K. (2005). How to construct recursive digital filters for baseflow separation. *Hydrological processes*, 19(2), 507-515.
- Eisenbies, M., Aust, W., Burger, J., & Adams, M. B. (2007). Forest operations, extreme flooding events, and considerations for hydrologic modeling in the Appalachians—A review. *Forest Ecology and Management*, 242(2), 77-98.
- United States Army Corps of Engineers (USACE). (2007). U.S. Army Corps of Engineers jurisdictional determination form instructional guidebook.
- Ford, C. R., Hubbard, R. M., & Vose, J. M. (2011). Quantifying structural and physiological controls on variation in canopy transpiration among planted pine and hardwood species in the southern Appalachians. *Ecohydrology*, 4(2), 183-195.
- Frady, C., Johnson, S., & Li, J. (2007). Stream macroinvertebrate community responses as legacies of forest harvest at the HJ Andrews Experimental Forest, Oregon. *Forest Science*, 53(2), 281-293.
- Freeman, M. C., Pringle, C. M., & Jackson, C. R. (2007). Hydrologic connectivity and the contribution of stream headwaters to ecological integrity at regional scales<sup>1</sup>: Wiley Online Library.
- Fritz, K. M., Hagenbuch, E., D'Amico, E., Reif, M., Wigington, P. J., Leibowitz, S. G., Nadeau, T. L. (2013). Comparing the extent and permanence of

- headwater streams from two field surveys to values from hydrographic databases and maps. *JAWRA Journal of the American Water Resources Association*, 49(4), 867-882.
- Goldman, D. (2010) *USDA-NRCS PLANTS Database*.
- Gravelle, J. A., Ice, G., Link, T. E., & Cook, D. L. (2009). Nutrient concentration dynamics in an inland Pacific Northwest watershed before and after timber harvest. *Forest Ecology and Management*, 257(8), 1663-1675.
- Hansen, W. F. (2001). Identifying stream types and management implications. *Forest Ecology and Management*, 143(1), 39-46.
- Hawkins, R., Woodward, D., & Jiang, R. (2002). *Investigation of the runoff curve number abstraction ratio*. Paper presented at the USDA-NRCS Hydraulic Engineering Workshop, Tucson, Arizona.
- Hawkins, R. H. (1973). Improved prediction of storm runoff in mountain watersheds. *Journal of the Irrigation and Drainage Division*, 99(4), 519-523.
- Hawkins, R. H. (1993). Asymptotic determination of runoff curve numbers from data. *Journal of Irrigation and Drainage Engineering*, 119(2), 334-345.
- Hayes, R. A. (1998). Soil Survey of Breathitt County, Kentucky.
- Herman, D. E., Stange, C. M., & Quam, V. C. (1996). North Dakota tree handbook. *USDA-NRCS ND State Soil Conservation Committee; NDSU Extension and Western Area Power Administration, Bismarck*.
- Hewlett, J. D., & Hibbert, A. R. (1967). Factors affecting the response of small watersheds to precipitation in humid areas. *Forest hydrology*, 1, 275-290.
- Hewlett, J. D., Lull, H. W., & Reinhart, K. G. (1969). In defense of experimental watersheds. *Water resources research*, 5(1), 306-316.
- Hibbert, A. R., & Gottfried, G. J. (1987). Stormflow responses to forest treatments on two Arizona mixed conifer watersheds. *General technical report RM-Rocky Mountain Forest and Range Experiment Station, US Department of Agriculture, Forest Service (USA)*.
- Hornbeck, J. N., & Edwards, P. J. (1990). *Effectiveness of three streamside management practices in the central Appalachians*. Paper presented at



- the Proceedings of the sixth biennial southern silvicultural research conference. Gen. Tech. Rep. SE-70. Asheville, NC: US Department of Agriculture, Forest Service, Southeastern Forest Experiment Station.
- Hornbeck, J., Adams, M., Corbett, E., Verry, E., & Lynch, J. (1993). Long-term impacts of forest treatments on water yield: a summary for northeastern USA. *Journal of Hydrology*, 150(2), 323-344.
- Hornbeck, J. W., & Leak, W. B. (1992). Ecology and management of northern hardwood forests in New England.
- Hornbeck, J. W., Martin, C., Pierce, R., Bormann, F., Likens, G., & Eaton, J. (1987). The northern hardwood forest ecosystem: ten years of recovery from clearcutting. *USDA Forest Service Report NE-RP-596*.
- Ice, G., & Binkley, D. (2003). Forest streamwater concentrations of nitrogen and phosphorus: A comparison with EPA's proposed water quality criteria. *Journal of Forestry*, 101(1), 21-28.
- Ice, G. G., & Stednick, J. D. (2004). Forest watershed research in the United States. *FOREST*, 17.
- Jewett, K., Daugharty, D., Krause, H. H., & Arp, P. A. (1995). Watershed responses to clear-cutting: effects on soil solutions and stream water discharge in central New Brunswick. *Canadian Journal of Soil Science*, 75(4), 475-490.
- Johnson, A. (2013). Accuracy assessments for headwater stream maps in western North Carolina.
- Kaplan, L., Bott, T., Jackson, J., Newbold, J., & Sweeney, B. (2008). Protecting headwaters: The scientific basis for safeguarding stream and river ecosystems. *Avondale: Stroud Water Research Centre*.
- Keppeler, E., Reid, L., & Lisle, T. (2008). *Long-term patterns of hydrologic response after logging in a coastal redwood forest*. Paper presented at the proceedings of the third interagency conference on research in the watersheds.
- Keppeler, E. T., & Ziemer, R. R. (1990). Logging effects on streamflow: Water yield and summer low flows at Caspar Creek in northwestern California.

- Water resources research*, 26(7), 1669-1679.
- Kochenderfer, J. N., Edwards, P. J., & Wood, F. (1997). Hydrologic impacts of logging an Appalachian watershed using West Virginia's best management practices. *Northern Journal of Applied Forestry*, 14(4), 207-218.
- Kochenderfer, J. N., & Hornbeck, J. W. (1999). *Contrasting timber harvesting operations illustrate the value of BMPs*. Paper presented at the Proceedings, 12th Central Hardwood Forest Conference.
- Lange, H., Lischeid, G., Hoch, R., & Hauhs, M. (1996). Water flow paths and residence times in a small headwater catchment at Gårdsjön, Sweden, during steady state storm flow conditions. *Water resources research*, 32(6), 1689-1698.
- Lautemann, E. (1998). State of the World, 1998. *Library Journal*, 123(18), 48.
- Leopold, L. B. (1964). *Fluvial Processes in Geomorphology*: W. H. Freeman.
- Lu, J., Sun, G., McNulty, S. G., & Amatya, D. M. (2003). Modeling actual evapotranspiration from forested watersheds across the southeastern United States: Wiley Online Library.
- Lynch, J. A., & Corbett, E. S. (1990). Evaluation of best management practices for controlling nonpoint pollution from silvicultural operations. *JAWRA Journal of the American Water Resources Association*, 26(1), 41-52.
- MacDonald, L. H., & Coe, D. (2007). Influence of headwater streams on downstream reaches in forested areas. *Forest Science*, 53(2), 148-168.
- Martin, C. W., Hornbeck, J. W., Likens, G. E., & Buso, D. C. (2000). Impacts of intensive harvesting on hydrology and nutrient dynamics of northern hardwood forests. *Canadian Journal of Fisheries and Aquatic Sciences*, 57(S2), 19-29.
- McClure, J., Kolka, R., & White, A. (2004). Effect of forest harvesting best management practices on coarse woody debris distribution in stream and riparian zones in three Appalachian watersheds. *Water, Air and Soil Pollution: Focus*, 4(1), 245-261.
- McDowell, R. C., Grabowski, G. J., & Moore, S. L. (1981). *Geologic map of*

- Kentucky*. United States Geological Survey.
- Meyer, J. L., Jensen, M. N., & Sutton, D. (2003). *Where rivers are born: the scientific imperative for defending small streams and wetlands*: American Rivers.
- Meyer, J. L., Strayer, D. L., Wallace, J. B., Eggert, S. L., Helfman, G. S., & Leonard, N. E. (2007). The contribution of headwater streams to biodiversity in river networks. *JAWRA Journal of the American Water Resources Association*, 43(1), 86-103.
- Mohlenbrock, R. H. (1991). Southern wetland flora: Field office guide to plant species. *USDA-SCS/South National Technical Center, Fort Worth*.
- Mohlenbrock, R. H. (1995). Northeast wetland flora: Field office guide to plant species. *USDA-NRCS/Northeastern National Technical Center, Chester*.
- Moore, A., Nowacki, G., & Burk, A. (2005). Rapid assessment reference condition model: *Potential natural vegetation group (Mixed Mesophytic Hardwood)*: USDA.
- Natural Resource Conservation Service (NRCS). (2004). NRCS National Engineering Handbook, Chapter 10: Estimation of Direct Runoff from Storm Rainfall.
- Neal, C., Robinson, M., Reynolds, B., Neal, M., Rowland, P., Grant, S., Lawlor, A. (2010). Hydrology and water quality of the headwaters of the River Severn: stream acidity recovery and interactions with plantation forestry under an improving pollution climate. *Science of the total environment*, 408(21), 5035-5051.
- Overstreet, J. (1984). Robinson forest inventory: 1980-1982. *University of Kentucky, College of Agriculture, Department of Forestry, Lexington, KY*.
- Pannill, H., McCoy, J., O'Ney, S., Bare, C., Primrose, N., & Bowen, S. (2000). Evaluating the effectiveness of Maryland's best management practices for forest harvest operations. *Maryland Department of Natural Resources Forest Service and Chesapeake & Coastal Watershed Service. Annapolis, Maryland*.
- Pflugmacher, D., Cohen, W., Kennedy, R., & Lefsky, M. (2008). Regional

- applicability of forest height and aboveground biomass models for the Geoscience Laser Altimeter System. *Forest Science*, 54(6), 647-657.
- Pierce, R. S., Hornbeck, J. W., Martin, C. W., Tritton, L. M., Smith, C. T., Federer, C. A., & Yawney, H. W. (1993). *Whole-tree clearcutting in New England: manager's guide to impacts on soils, streams, and regeneration*. US Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.
- Pritchett, W., & Fisher, R. (1987). Nutrient cycling in forest ecosystems. *Properties and Management of Forest Soils (2nd edition)*. Wiley, New York, 180-204.
- Profant, D. (September 27, 2011). Trees, Shrubs, & Woody Vines of Southeastern Ohio.
- Reeves, C. D. (2012). Effectiveness of Elevated Skid Trail Headwater Stream Crossings in the Cumberland Plateau.
- Restrepo, M., & Waisanen, P. (2004). *Strategies for stream classification using G/S*. Paper presented at the proceedings of the 24th annual ESRI international user conference, San Diego, California.
- Reuss, J. (1983). Implications of the calcium-aluminum exchange system for the effect of acid precipitation on soils. *Journal of Environmental Quality*, 12(4), 591-595.
- Richardson, C. J. (1989). Freshwater wetlands: transformers, filters, or sinks. *freshwater wetlands and wildlife. US DOE Office of Scientific and Technical Information, Oak Ridge, TN*, 25-46.
- Richardson, J. S., & Danehy, R. J. (2007). A synthesis of the ecology of headwater streams and their riparian zones in temperate forests. *Forest Science*, 53(2), 131-147.
- Rickert, D. (1985). Water quality—Compensation for discharge in detecting trends in water quality data. Quality of Water Branch Memorandum No. 85.17. Reston, Va. *US Geological Survey*.
- Ruhlman, M. B. (1999). Effectiveness of forestry best management practices: evaluating water quality from intensely managed watersheds.

- Sanderson, T. M. (2014). Monitoring the influence of acid deposition on soil and implications to forest health in the Daniel Boone National Forest. Thesis (M.S.)--University of Kentucky, 2014.
- Santhi, C., Allen, P., Muttiah, R., Arnold, J., & Tuppad, P. (2008). Regional estimation of base flow for the conterminous United States by hydrologic landscape regions. *Journal of Hydrology*, 351(1), 139-153.
- Service, U. F. (2012). *National best management practices for water quality management on national forest systems land*. Washington, D.C. : United States Department of Agriculture, Forest Service.
- Soil Conservation Service (SCS). (1972). *SCS National Engineering Handbook, Section 4: Hydrology*: The service.
- Stednick, J., & Kern, T. (1992). Long term effects of timber harvesting in the Oregon Coast Range: the New Alsea Watershed Study (NAWS). *Interdisciplinary Approaches to Hydrology and Hydrogeology. American Institute of Hydrology, Smyrna, GA*, 502-510.
- Stednick, J. D. (1996). Monitoring the effects of timber harvest on annual water yield. *Journal of Hydrology*, 176(1), 79-95.
- Stednick, J. D. (2008). Long-term streamflow changes following timber harvesting. *Hydrological and biological responses to forest practices: The Alsea Watershed Study* (pp. 139-155). New York, NY: Springer New York.
- Stephen M. Seiberling, A. S. W., and Peter S. White. (2006). Common trees of the North Carolina Piedmont: Identification, descriptions, illustrations, and glossary.
- Stoy, P. C., Katul, G. G., Siqueira, M., Juang, J. Y., Novick, K. A., McCarthy, H. R., Oren, R. (2006). Separating the effects of climate and vegetation on evapotranspiration along a successional chronosequence in the southeastern US. *Global Change Biology*, 12(11), 2115-2135.
- Stringer, J. W., & Perkins, C. (2001). *Kentucky forest practice guidelines for water quality management*. Cooperative Extension Service, University of Kentucky, College of Agriculture.
- Stuart, G., & Dunshie, D. (1976). Effects of timber harvest on water chemistry.

- Hydrol. Pap. [Milwaukee, WI]: U.S. Department of Agriculture, Forest Service, Eastern Region. 34 p.*
- Stuart, G. W., & Edwards, P. J. (2006). Concepts about forests and water. *Northern Journal of Applied Forestry*, 23(1), 11-19.
- Sun, G., McNulty, S. G., Lu, J., Amatya, D. M., Liang, Y., & Kolka, R. (2005). Regional annual water yield from forest lands and its response to potential deforestation across the southeastern United States. *Journal of Hydrology*, 308(1), 258-268.
- Swank, W. T., & DA Jr, C. (2012). *Forest hydrology and ecology at Coweeta* (Vol. 66): Springer Science & Business Media.
- Swank, W. T., Vose, J., & Elliott, K. (2001). Long-term hydrologic and water quality responses following commercial clearcutting of mixed hardwoods on a southern Appalachian catchment. *Forest Ecology and Management*, 143(1), 163-178.
- Taylor, T., Agouridis, C., Warner, R., & Barton, C. (2009). Runoff curve numbers for loose-dumped spoil in the Cumberland Plateau of eastern Kentucky. *International Journal of Mining, Reclamation and Environment*, 23(2), 103-120.
- Troendle, C., & Olsen, W. (1994). Potential effects of timber harvest and water management on streamflow dynamics and sediment transport.
- United States Energy Information Administration (USEIA). (2012). Form EIA-923, Power Plant Operations Report. Table 3.7 Net Generation from Coal.
- United States Environmental Protection Agency (USEPA). 1990. Clean Air Act (CAA) amendments. 42 U.S.C. 7401-7671q.
- United States Environmental Protection Agency (USEPA). 2012. Clean Air Act (CAA) amendments.
- United States Environmental Protection Agency (USEPA). (1992). *Managing nonpoint source pollution, final report to Congress on Section 319 of the Clean Water Act (1989)*. Washington DC.

- Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., & Cushing, C. E. (1980). The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(1), 130-137.
- Villines, J. A., Agouridis, C. T., Warner, R. C., & Barton, C. D. (2015). Using GIS to delineate headwater stream origins in the Appalachian coalfields of Kentucky. *JAWRA Journal of the American Water Resources Association*, 51(6), 1667-1687.
- Vose, J. M., Sun, G., Ford, C. R., Bredemeier, M., Otsuki, K., Wei, X., Zhang, L. (2011). Forest ecohydrological research in the 21st century: what are the critical needs. *Ecohydrology*, 4(2), 146-158.
- Wallace, L. (1988). Comparative physiology of successional forest trees. *Forest hydrology and ecology at Coweeta* (pp. 181-189): Springer.
- Wang, X., Burns, D. A., Yanai, R. D., Briggs, R. D., & Germain, R. H. (2006). Changes in stream chemistry and nutrient export following a partial harvest in the Catskill Mountains, New York, USA. *Forest Ecology and Management*, 223(1), 103-112.
- Warner, R. C., & Agouridis, C. T. (2010a). Valley fill design and construction to improve ecological performance. *United States Environmental Protection Agency (USEPA)*.
- Warner, R. C., Agouridis, C. T., Vingralek, P. T., & Fogle, A. W. (2010b). Reclaimed mineland curve number response to temporal distribution of rainfall. *JAWRA Journal of the American Water Resources Association*, 46(4), 724-732.
- Weatherford, M. K. (2014). *Storm hydrograph characteristics and curve numbers of loose-dumped mine spoil in eastern Kentucky*. Thesis (M.S.)--University of Kentucky, 2014.
- Wei, X., & Zhang, M. (2010). Quantifying streamflow change caused by forest disturbance at a large spatial scale: a single watershed study. *Water resources research*, 46(12).
- Wilm, H. G. (1949). How long should experimental watersheds be calibrated. *Eos, Transactions American Geophysical Union*, 30(2), 272-278.

- Witt, E. L. (2012). Evaluating streamside management zone effectiveness in forested watersheds of the Cumberland Plateau. Thesis (Ph. D.)-- University of Kentucky, 2012.
- Wullschleger, S. D., Hanson, P., & Todd, D. (2001). Transpiration from a multi-species deciduous forest as estimated by xylem sap flow techniques. *Forest Ecology and Management*, 143(1), 205-213.
- Zhang, L., Hickel, K., Dawes, W., Chiew, F. H., Western, A., & Briggs, P. (2004). A rational function approach for estimating mean annual evapotranspiration. *Water resources research*, 40(2).



## **VITA**

### **PLACE OF BIRTH**

Muncie, IN

### **EDUCATION**

B.S. Agricultural and Biological Engineering, Purdue University West Lafayette, Indiana, May 2014

Graduate Certificate, Stream and Watershed Science, University of Kentucky, Lexington, Kentucky, December 2015

### **PROFESSIONAL EXPERIENCE**

Graduate Research Assistant, Department of Biosystems and Agricultural Engineering, University of Kentucky, Lexington, Kentucky, August 2011 - Present  
Advisor: Dr. Carmen Agouridis

Teaching Assistant, Department of Biosystems and Agricultural Engineering, University of Kentucky, Lexington, Kentucky, January 2015 - May 2015

Lab Instructor, Department of Biosystems and Agricultural Engineering, University of Kentucky, Lexington, Kentucky, January 2015 - May 2015

Research Assistant, Department of Agricultural Engineering, University of Viçosa, Viçosa, Brazil, May 2012 - July 2012

### **LICENSURE AND CERTIFICATION**

Engineer-in-Training (EIT), Indiana (May 2014)

### **PROFESSIONAL ASSOCIATIONS**

Alpha Epsilon